

Appendix II. Summary of the confirmed human-caused mortality and serious injury (SI) events involving baleen whale stocks along the Gulf of Mexico Coast, US East Coast, and adjacent Canadian Maritimes, 2006-2010, with number of events attributed to entanglements or vessel collisions by year.

Stock	Mean annual mortality and SI rate (PBR ¹ for reference)	Entanglements			Vessel Collisions		
		Annual rate (US waters / Canadian waters)	Confirmed mortalities (2005, 2006, 2007, 2008, 2009)	Confirmed SIs (2005, 2006, 2007, 2008, 2009)	Annual rate (US waters / Canadian waters)	Confirmed mortalities (2005, 2006, 2007, 2008, 2009)	Confirmed SIs (2005, 2006, 2007, 2008, 2009)
Western North Atlantic right whale (<i>Eubalaena glacialis</i>)	2.8 (0.9)	1.8 (1.6/0.2)	(1, 1, 0, 0, 2)	(0, 0, 1, 2, 2)	1.2 (0.8 / 0.4)	(4, 0, 0, 0, 1)	(1, 0, 0, 0, 0)
Gulf of Maine humpback whale (<i>Megaptera novaeangliae</i>)	7.8 (2.7)	5.8 (5.6 / 0.2)	(1, 1, 2, 2, 3)	(4, 2, 4, 3, 7)	2.0 (2.0 / 0)	(3, 3, 1, 0, 3)	0
Western North Atlantic fin whale (<i>Balaenoptera physalus</i>)	This report is still under review.						
Nova Scotian sei whale (<i>B. borealis</i>)	This report is still under review.						
Western North Atlantic blue whale (<i>B. musculus</i>)	0 (0.9)	0	0	0	0	0	0
Canadian East Coast minke whale (<i>B. acutorostrata</i>)	This report is still under review.						
Western North Atlantic Bryde's whale (<i>B. edeni</i>)	0.2 (0.1)	0	0	0	0.2 (0.2 / 0)	(0, 0, 0, 1, 0)	0

¹ Potential Biological Removal (PBR)