



## All Species

### Comparison of Differences



## Bluefish


### Comparison of Differences



**Figure 7. The distribution of differences between the standard deviation of the average kept pounds of unobserved and observed trips for SBRM 2009 (July 2007 through June 2008), SBRM 2010 (July 2008 through June 2009), and SBRM 2011 (July 2009 through June 2010), by SBRM species group.**

## Atlantic Herring

### Comparison of Differences




## Red Crab

No graph available

**Figure 7, continued. The distribution of differences between the standard deviation of the average kept pounds of unobserved and observed trips for SBRM 2009 (July 2007 through June 2008), SBRM 2010 (July 2008 through June 2009), and SBRM 2011 (July 2009 through June 2010), by SBRM species group.**

## Fluke-Scup-Black Sea Bass

### Comparison of Differences



## Large Mesh Groundfish

### Comparison of Differences





Figure 7, continued. The distribution of differences between the standard deviation of the average kept pounds of unobserved and observed trips for SBRM 2009 (July 2007 through June 2008), SBRM 2010 (July 2008 through June 2009), and SBRM 2011 (July 2009 through June 2010), by SBRM species group.

## Monkfish

### Comparison of Differences



## Sea Scallop

### Comparison of Differences





Figure 7, continued. The distribution of differences between the standard deviation of the average kept pounds of unobserved and observed trips for SBRM 2009 (July 2007 through June 2008), SBRM 2010 (July 2008 through June 2009), and SBRM 2011 (July 2009 through June 2010), by SBRM species group.

## Skate Complex

### Comparison of Differences



## Small Mesh Groundfish

### Comparison of Differences





Figure 7, continued. The distribution of differences between the standard deviation of the average kept pounds of unobserved and observed trips for SBRM 2009 (July 2007 through June 2008), SBRM 2010 (July 2008 through June 2009), and SBRM 2011 (July 2009 through June 2010), by SBRM species group.

## Spiny Dogfish

### Comparison of Differences




## Surfclam and Ocean Quahog

No graph available

**Figure 7, continued. The distribution of differences between the standard deviation of the average kept pounds of unobserved and observed trips for SBRM 2009 (July 2007 through June 2008), SBRM 2010 (July 2008 through June 2009), and SBRM 2011 (July 2009 through June 2010), by SBRM species group.**

## Squid-Butterfish-Mackerel

### Comparison of Differences



## Tilefish

### Comparison of Differences




Figure 7, continued. The distribution of differences between the standard deviation of the average kept pounds of unobserved and observed trips for SBRM 2009 (July 2007 through June 2008), SBRM 2010 (July 2008 through June 2009), and SBRM 2011 (July 2009 through June 2010), by SBRM species group.