

Stock Assessment Update for Ocean Quahogs (*Arctica islandica*) through 2011

by Antonie Chute¹, Daniel Hennen¹, Robert Russell² and Larry Jacobson¹

¹NOAA's National Marine Fisheries Service, 166 Water St., Woods Hole MA 02543

²Maine Dept. of Natural Resources, 194 McKown Pt. Rd., West Boothbay Harbor, ME 04575

U.S. DEPARTMENT OF COMMERCE
National Oceanic and Atmospheric Administration
National Marine Fisheries Service
Northeast Fisheries Science Center
Woods Hole, Massachusetts

May 2013

Northeast Fisheries Science Center Reference Documents

This series is a secondary scientific series designed to assure the long-term documentation and to enable the timely transmission of research results by Center and/or non-Center researchers, where such results bear upon the research mission of the Center (see the outside back cover for the mission statement). These documents receive internal scientific review, and most receive copy editing. The National Marine Fisheries Service does not endorse any proprietary material, process, or product mentioned in these documents.

All documents issued in this series since April 2001, and several documents issued prior to that date, have been copublished in both paper and electronic versions. To access the electronic version of a document in this series, go to <http://www.nefsc.noaa.gov/publications/>. The electronic version is available in PDF format to permit printing of a paper copy directly from the Internet. If you do not have Internet access, or if a desired document is one of the pre-April 2001 documents available only in the paper version, you can obtain a paper copy by contacting the senior Center author of the desired document. Refer to the title page of the document for the senior Center author's name and mailing address. If there is no Center author, or if there is corporate (*i.e.*, non-individualized) authorship, then contact the Center's Woods Hole Laboratory Library (166 Water St., Woods Hole, MA 02543-1026).

Information Quality Act Compliance: In accordance with section 515 of Public Law 106-554, the Northeast Fisheries Science Center completed both technical and policy reviews for this report. These predissemination reviews are on file at the NEFSC Editorial Office.

This document may be cited as:

Chute A, Hennen D, Russell R, Jacobson L. 2013. Stock assessment update for ocean quahogs (*Arctica islandica*) through 2011. US Dept Commer, Northeast Fish Sci Cent Ref Doc. 13-17; 156 p. Available from: National Marine Fisheries Service, 166 Water Street, Woods Hole, MA 02543-1026, or online at <http://www.nefsc.noaa.gov/publications/>

Contents

EXECUTIVE SUMMARY	iii
INTRODUCTION	1
Entire stock vs. the exploited region.....	1
Fishable stock.....	2
Units of measurement	2
Previous assessments	2
Biological characteristics	2
Recruitment.....	3
COMMERCIAL FISHERY DATA.....	4
Prices.....	4
Fishing effort.....	4
Landings per unit effort (LPUE).....	5
Spatial patterns in fishery data.....	5
Fishery data by ten-minute square (TMSQ)	6
Trends for important TMSQ.....	6
Bycatch and discard.....	7
Commercial size selectivity	7
Commercial size-composition data.....	8
FISHERY INDEPENDENT DATA	8
NEFSC clam survey.....	8
Survey tow distance and gear performance based on sensor data	9
Survey tow distance and gear performance in trend analysis.....	10
Survey gear selectivity.....	10
Survey catch used to make stock and fishable abundance/biomass estimates.....	10
Depletion studies and survey dredge efficiency	11
Patch model.....	11
BIOMASS AND FISHING MORTALITY	12
Survey Results	12
Efficiency-corrected swept area biomass.....	13
ESB-based fishing mortality estimates	14
Uncertainty in ESB and ESB-based fishing mortality estimates	15
KLAMZ model	15

KLAMZ model configurations	16
Parameters estimated	17
Variance estimates	17
KLAMZ results	17
DMV region	18
NJ region.....	18
LI region.....	18
SNE region.....	19
GBK region.....	19
Exploited area and entire stock.....	19
Historical retrospective analyses.....	20
REFERENCE POINTS AND STOCK STATUS.....	21
Biological Reference Points.....	21
B_{target}	21
$B_{\text{threshold}}$	21
F_{target}	21
$F_{\text{threshold}}$	21
STOCK STATUS	22
PROJECTIONS	22
REFERENCES	23
TABLES	27
FIGURES.....	50
APPENDIX 1.....	130
APPENDIX 2.....	150

EXECUTIVE SUMMARY

This assessment update for ocean quahog in the US EEZ is based on fishery-dependent data from 1978 through 2011, NEFSC clam survey data from 1982 through 2011, and biological information. Since this is an assessment update, there are no specific terms of reference addressed. In addition, shell length-meat weight relationships, survey and commercial dredge efficiency, and size selectivity estimates were not updated to include new depletion experiment data collected in 2011, and assessment models were configured in the same way as in the previous assessment (NEFSC 2009). These types of auxiliary information will be updated in the next ocean quahog benchmark assessment.

According to the most recent data, ocean quahogs in the US EEZ are not overfished, and overfishing is not occurring. Total fishable stock biomass (all regions) during 2011 was 2.961 million mt of meats, which is above both the biomass target (1.73 million mt) and biomass threshold (1.39 million mt). The fishing mortality rate during 2011 for the exploited region (all areas but GBK) was $F=0.01\text{ y}^{-1}$, which is below the overfishing threshold $F_{45\%}=0.022\text{ y}^{-1}$.

Landings and status table (weights in thousands of mt meats): Ocean quahog

Year:	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	¹ Min	Max	Mean
² Quotas:													
ITQ fishery	20.4	20.4	22.7	24.2	24.2	24.2	24.2	24.2	24.2	24.2	15.9	27.2	22.2
Maine	0.30	0.30	0.30	0.30	0.30	0.30	0.30	0.30	0.30	0.30	0.30	0.30	0.30
³ Landings:													
ITQ fishery	17.9	18.8	17.7	13.6	14.3	15.6	15.5	15.9	16.1	14.1	12.1	22.4	17.7
Maine	0.39	0.36	0.31	0.30	0.37	0.31	0.20	0.17	0.17	0.20	0.01	0.72	0.36
Total	18.3	19.2	18.0	13.9	14.7	15.9	15.7	16.1	16.3	14.3			
Biomass (whole stock):	3,431	3,378	3,324	3,270	3,221	3,170	3,119	3,067	3,014	2,961	2,961	3,946	3,557
Fishing mortality (exploited area):	0.0102	0.0109	0.0105	0.0083	0.0088	0.0098	0.0101	0.0107	0.0112	0.0101	0.0050	0.0110	0.0090
Fishing mortality (whole stock):	0.0052	0.0055	0.0053	0.0041	0.0044	0.0049	0.0049	0.0052	0.0054	0.0049	0.0030	0.0060	0.0050
⁴ Recruitment:	0.072	0.072	0.072	0.072	0.072	0.072	0.072	0.072	0.072	0.072	0.072	78.74	N/A

¹Min, max & mean: Maine quota & landings = 1990-2011; EEZ quota & landings = 1980-2011; EEZ B, F & recruitment = 1978-2011.

² Quotas, landings, stock biomass and recruitment expressed in thousands of metric tons meats.

³ Landings not adjusted for incidental mortality, which is assumed to be 5% of landings. Discards are very low.

⁴ Recruitment is an estimated average assuming zero recruitment in DMV and SVA. 78.74 was the level estimated before 1993.

ocean quahog stock assessment region	abbreviation
U.S. Exclusive Economic Zone	EEZ
Maine	MNE
Georges Bank	GBK
southern New England	SNE
Long Island	LI
New Jersey	NJ
Delaware/Maryland/Virginia	DMV
southern Virginia and North Carolina	SVA

In 2009, the ocean quahog Stock Assessment Review Committee (SARC) recommended new, more conservative reference points for the biomass threshold, fishing mortality threshold, and target fishing mortality. The proposed reference points were approved by the Mid Atlantic Fishery Management Council (MAFMC) in October 2009 for inclusion in Amendment 15 to the ocean quahog and surfclam Fishery Management Plan (FMP) expected to be implemented in 2013. The newly recommended $B_{Threshold}$ of 1.39 million mt is 40% of the 1978 (considered pre-fishery) biomass and the recommended biomass target is 1.73 million mt, which is one-half of the 1978 biomass. $F_{45\%}$ was recommended as a better F_{MSY} proxy for an exceptionally long-lived species with a very low natural mortality rate like ocean quahogs. The current status of the ocean quahog stock remains not overfished and without overfishing relative to these proposed new reference points.

The current FMP requires comparison of the overfishing threshold reference point to fishing mortality occurring in the exploited portion of the stock only, which currently includes the SVA, DMV, NJ, LI, and SNE regions. The GBK region, which has not been open to ocean quahog fishing since 1990 due to the risk of paralytic shellfish poison (PSP) contamination, contains about 43% of total ocean quahog fishable biomass (2011 NEFSC survey data). However, much of Georges Bank has reopened for both ocean quahog and surfclam, as effective new at-sea PSP testing protocols have been developed. Harvest policies for ocean quahog may need to be reconsidered if a fishery develops on Georges Bank. However, there have never been any appreciable ocean quahog landings from GBK due to the distance from processing plants and difficult fishing conditions there.

The ocean quahog fishery has shifted north over the last three decades as catch rates decreased on the original fishing grounds off Delmarva and New Jersey. In the 1980s, the bulk of the fishing effort was off Delmarva and southern New Jersey. In the 1990s, effort fell by half in Delmarva, as effort moved northward to the Long Island and Southern New England regions. For the past ten years, the majority of fishing effort has been in the Long Island region.

Fishing effort has declined in the ocean quahog fishery from a high of over 40,000 hours per year during the 1990s to about 25,000 hours per year recently. The number of active vessels in the EEZ in 2011 was the lowest on record, but the average size of ocean quahog vessels has also increased over time. LPUE for the EEZ stock as a whole has been fairly stable since 1982, but is currently higher in northern areas (LI and SNE) than in the south (NJ and DMV). Landings have declined since the peak of almost 22,500 mt in 1992 to a little more than 14,000 mt in 2011.

Based on assessment data, the ocean quahog population is an unproductive stock with infrequent recruitment, and thus vulnerable to overfishing. After three decades of fishing at a low F , the stock as a whole is being fished down. In 2011, fishable stock biomass in the southernmost regions of SVA, DMV and NJ was less than half of 1978 pre-fishery levels. (Recommended target biomass for the stock as a whole is 50% of the pre-fishery biomass.) Biomass in the more northern regions of LI and SNE increased after 1978 to due to a recruitment event and growth, but then began to decrease in the early 1990s when recruitment declined, and the fishery gradually began to move north into these regions. Recruitment events appear to be localized and separated by decades, although survey length frequencies show that a low level of recruitment occurs on a continuous basis.

Using updated survey and fishery data, the KLAMZ model was run for the whole stock, the exploited part of the stock only, and each individual assessment region except for SVA. For both the exploited part of the stock (SVA, DMV, NJ, LI, and SNE combined) and the whole stock (exploited regions plus GBK), fishable biomass continued to trend downward. Fishing mortality trended upward from 1978 through 1990 in the exploited region, but has been fairly stable since, as catches and biomass have both fallen. Estimates of fishable biomass are decreasing in all regions. KLAMZ biomass estimates for GBK in the last assessment showed a slight upward trend, but a smaller estimated survey swept-area biomass in 2011 affected the model trends in biomass and surplus production negatively.

The estimates of biomass and fishing mortality for the EEZ stock in this assessment update do not include the Maine mahogany quahog fishery, which started in inshore Maine but now takes place almost entirely in federal waters using 100,000 (Maine) bushels of the EEZ quota. The Maine stock biomass is small (~1% relative to the rest of the EEZ with landings of about 322 mt meats during 2012) and fishing effort is concentrated in a small area: just two beds separated by a few miles. Appendix 1, which was prepared in collaboration with Robert Russell of the Maine Department of Marine Resources, gives updated stock assessment information about ocean quahogs in Maine waters.

INTRODUCTION

Ocean quahog (*Arctica islandica*) in the US EEZ is regarded as a single stock. The EEZ fishery (with landings of about 14,300 mt meats during 2011) is managed under a single individual transferable quota (ITQ) system that was established for ocean quahogs and Atlantic surfclams (*Spisula solidissima*) in 1990. Murawski and Serchuk (1989) and Serchuk and Murawski (1997) provide detailed information about the history and operation of the EEZ fishery, which began around 1975. The estimates of biomass and fishing mortality for the EEZ stock in this assessment update do not include the Maine mahogany quahog fishery, which started in inshore Maine but now takes place almost entirely in federal waters. The Maine stock biomass is small (~1% relative to the rest of the EEZ) and fishing effort is concentrated in a small area: just two beds separated by a few miles. The rest of the EEZ and the Maine ocean quahog populations have different biological characteristics, support fisheries that are managed separately, use different vessels and gear, and provide different products. Updated information for the Maine fishery is presented as Appendix 1. Landings from Maine are included when total EEZ landings are calculated. Annually, 100,000 Maine bushels from the EEZ quota are allocated to the Maine fishery. Since about 1999, the Maine quota has often been exhausted and Maine fishermen have leased ITQ shares in order to exceed the Maine allocation.

The ocean quahog stock is often broken down into smaller regions (listed below) based on biology, fishery characteristics, and history (Figure 1). These designated regions are important in understanding the fishery but have no legal meaning.

Region	Abbreviation
US exclusive economic zone	EEZ
Maine	MNE
Georges Bank	GBK
Southern New England	SNE
Long Island	LI
New Jersey	NJ
Delmarva	DMV
Southern Virginia and North Carolina	SVA
mid-Atlantic Bight	MAB

Entire stock vs. the exploited region

Data and analysis for both the entire stock of ocean quahogs in the EEZ and the “exploited region” are presented in this assessment update. The entire stock refers to ocean quahogs in all six assessment regions: SVA, DMV, NJ, LI, SNE, and GBK. The “exploited region,” by contrast, excludes GBK, as it has been closed to ocean quahog harvesting since 1990 as a precaution against paralytic shellfish poison (PSP). Data from the 2011 NEFSC clam survey indicate that 43% of ocean quahog biomass is found in the GBK region. As of December 2012, much of Georges Bank reopened for both ocean quahog and surfclam fishing as effective new at-

sea PSP testing protocols have been developed and tested. However, as of this assessment update GBK has not been fished for ocean quahogs. The reason for the distinction between the whole stock and the exploited region is that certain management limits apply only to the exploited region, such as the $F_{threshold}$.

Fishable stock

Fishable ocean quahogs are animals large enough to be taken by the fishery based on the size selectivity curve of commercial gear. Stock biomass is always larger than fishable biomass, as it includes the quahogs that are too small to be taken by the fishery.

Units of measurement

Body size in ocean quahogs is measured in terms of shell length (SL), which is the longest distance along the vertical axis of an intact specimen.

Units of measure for ocean quahogs used in this assessment update are described below. Commercial data are reported in industry bushels in logbooks, and often converted to meat weight (which includes all soft tissue within the shell) for use in this assessment update.

Unit	Equivalent
Industry or Mid-Atlantic bushel (Industry bu)	1.88 ft ³
Maine (US standard) bushel (Maine bu)	1.2448 ft ³
Industry bushels x 10	Pounds meat wt
Industry bushels x 4.5359	Kilograms meat wt
Maine bushel	0.662 industry bushels
Cage	32 Industry bushels

Previous assessments

Stock assessments for ocean quahog in the EEZ were completed by the NEFSC in 1995, 1998, 2000, 2004, 2007, and 2009. The last assessment (NEFSC 2009) concluded that the EEZ ocean quahog resource was not overfished and that overfishing was not occurring.

Biological characteristics

Ocean quahogs are found in the north Atlantic, from Spain on the east side, up to Iceland, and down to Cape Hatteras NC on the west side (Theroux and Wigley 1983; Thorarinsdottir and Einarsson 1996; Lewis et al. 2001). They are found at depths of 10-400 m, depending on latitude. Shallower water habitats, such as the waters around Iceland, are utilized in the northern end of their range (Theroux and Wigley 1983; Thompson et al. 1980). The US stock is mostly found at depths of 25-95 m. Dahlgren et al. (2000) found no genetic differences between samples taken along the US coast from Maine to Virginia, based on mitochondrial cytochrome *b* gene frequencies.

The natural mortality rate and longevity of ocean quahogs are uncertain. Ocean quahogs are extremely long-lived. Individual specimens are commonly aged at over 200 years (Jones 1980; Steingrímsson and Thorarinsdottir 1995; Kilada et al. 2007; Strahl et al. 2007). Ageing studies of ocean quahogs off New Jersey and Long Island (Thompson et al. 1980; Murawski et al. 1982)

established that clams ranging in age from 50-100 years are common. Recently, Wanamaker et al. (2008) aged two ocean quahogs at 287 and 405 years, and Butler et al. (2012) aged one at 507 years, possibly the oldest non-colonial animal ever documented. Based on mean longevity estimates of around 200 years, adult ocean quahogs in the EEZ and off Iceland are assumed to die from natural causes at the rate of about 2% annually (instantaneous rate of natural mortality M of 0.02 per year). When $M=0.02$, about 1% of a cohort is expected to survive after 230 years. Estimates of M for two different unexploited populations of ocean quahogs from Canadian waters (Sable Bank and St Mary's Bay) were 0.03 and 0.10 respectively based on age-frequency data for unexploited populations (Kilada et al. 2007), which suggests there is some variation in natural mortality among locations.

Ocean quahogs grow slowly after the first few years of life (Lewis et al. 2001; Kilada et al. 2007) and typically reach a maximum size of about 110 mm SL, although larger specimens are found in small numbers. Individuals large enough to recruit to the fishery grow less than one percent per year in meat weight and less than 1 mm per year in shell length. Growth has been found to be faster on GBK than further south in the mid-Atlantic Bight (MAB). Ocean quahogs reach 73 mm (50% commercial selectivity) at age 13 years in GBK and 28 years in MAB (NEFSC 2009).

Maturity information specifically for ocean quahogs in the US EEZ is limited (see review in Cargnelli et al. 1999), but size and age at maturity appear to be variable. Off Long Island, the smallest mature ocean quahogs found were 36-41 mm SL and 6 years old, but some were also found to be still sexually immature at the age of 14 years (Thompson et al. 1980; Ropes et al. 1984). Females are more common than males among the oldest and largest individuals in the population (Ropes et al. 1984; Fritz 1991).

The SL/maturity relationship used in the KLAMZ model runs for this assessment update (Figure 2) is from Icelandic ocean quahogs (Thorarinsdottir and Jacobson 2005). The curve indicates 50% of female ocean quahogs are mature at 64 mm (19 years, based on the growth curve in Lewis et al. 2001 for MAB) and 90% are mature at 88 mm SL and 61 years old. Based on the commercial selectivity curve (Figure 2), about 50% of ocean quahogs are selected by the fishery at 72 mm SL (28 years, based on the growth curve for MAB) and 90% are selected by 93mm (86 years). The majority of ocean quahogs have likely reached maturity before recruitment to the fishery.

Shell length-meat weight (SLMW) relationships are important for ocean quahogs because survey catches in number are converted to meat weights based on shell length for many analyses. The SLMW relationships used for this assessment update, estimated using a mixture of frozen and fresh meat samples, are region-specific and have been used since the 2004 assessment (Table 1). A large number of fresh meat samples (which may yield a more accurate weight than frozen samples) have been collected during recent surveys, which will allow the SLMW relationships to be re-estimated in the near future.

Recruitment

Ocean quahog recruitment events are regional and infrequent (Powell and Mann 2005; Harding et al. 2008). The regular presence of small ocean quahogs in survey length frequencies indicates that low-level recruitment is continually occurring, particularly in the northern regions. But thirty

years of survey data (1982-2011) show only a few regional-scale recruitment events occurring over this time where there has been a distinct mode of smaller quahogs visible in the survey length frequency (for example, see Figure 32D: SNE survey length frequencies for 1997 and 2005). Because growth is so slow, there is a delay of one to three decades between larval settlement and production of recruits to the fishery. Ocean quahog recruitment patterns at biomass levels reduced by fishing are a major uncertainty for the fishery (NEFSC 2007).

COMMERCIAL FISHERY DATA

There are no recreational landings of ocean quahogs, as they are found far offshore in deep water, require expensive gear and vessels to harvest, and provide essentially the same product recreational fishermen can get from inshore quahogs (*Mercenaria mercenaria*) with less effort. Landings of ocean quahogs from state waters (inshore of three miles) are effectively zero.

Since 1980, ocean quahog and surfclam vessel captains have been required to fill out specialized trip logbooks, and these are the primary source of landings, location, and effort data for ocean quahogs. Landings and quotas for the ITQ fishery are usually reported in industry bushels (each industry bushel holds 1.88 ft³ of quahogs, about 10 lbs of meats), as clams are offloaded in cages that hold 32 bushels each, and these serve as a unit of measure for processors and vessel operators. For this assessment update, biomass and landings have been converted to meat weights unless otherwise noted.

Total EEZ landings were relatively high during the late 1980s through the mid-1990s, with a peak of 22,500 mt meats (or 4.9 million ITQ bushels) during 1992 (Figure 3; Table 2). After 1996, landings began a steady decline that lasted until 2002, when they increased again by about a third and stayed high for three years. It was in 2002 that the majority of ocean quahog landings began to come from the LI region. After 2004, landings dropped off again and have been fairly stable since. Industry sources report that lower landings during the most recent years have been due to reduced market demand. Landings averaged 91% of the EEZ quota during 1990-2000 and 71% of the EEZ quota during 2001-2011.

Prices

Nominal ex-vessel prices for ITQ ocean quahog landings, expressed as dollars per pound of meats (Figure 4; Table 3) have doubled since the mid 1980s from 31 cents to 65 cents. Adjusted for inflation, prices have been stable or increased slightly during this time.

Fishing effort

Total days (24 hours) fished by the ITQ fishery have decreased from a high of more than 1900 days per year in the early 2000s to less than 1000 days per year in 2011 (Figure 5; Table 4), which is roughly the same as the early 1980s. The total number of trips taken by the ITQ fishery has decreased steadily from about 3,500 trips per year in 1989 to about 1,000 trips in 2011 (Figure 6). In 2009, however, there appeared to be an unusual number of trips into the SNE region, which effectively doubled the total number of ocean quahog trips compared to the year before or the year after. As the number of trips per year has declined, trip duration has increased (Figure 7). The number of vessels active in the ITQ ocean quahog fishery has declined since

2003 (Figure 8), but the average ocean quahog vessel has gotten larger (Figure 9) and landings per trip have doubled since the 1980s (Figure 10).

Landings per unit effort (LPUE)

LPUE (measured as bushels landed per hour fished) for the ocean quahog fishery is mostly a measure of fishing success rather than stock abundance, as any negative change in stock abundance can be masked by the transfer of fishing effort to areas where ocean quahog density remains high. In spite of this, LPUE and NEFSC clam survey data are highly correlated for southern areas (DMV and NJ), where significant levels of fishing have occurred over the longest period of time (NEFSC 2007).

Because the fishery can move to new areas when LPUE begins to decline, LPUE for the ITQ fishery as a whole has been basically stable since 1980, between 100 and 150 bushels per hour (Figure 11; Table 5). Since 1980, LPUE has declined in the traditional southern fishing grounds of DMV and NJ to around 60-80 bushels per hour, but LPUE in the LI and SNE regions, where the majority of fishing effort has been since 1990, has been stable or increasing.

The break-even LPUE for ocean quahogs (where variable costs and revenues are the same) was 80 bushels per hour in 2004 (NEFSC 2004). Inflation, increased steaming time to offshore fishing grounds, operation of new larger vessels, increased costs for food, insurance, etc., and especially the variability in the price of fuel, make it difficult to estimate a current break-even LPUE, but it is probably higher than it was in 2004.

For the 2006 assessment (NEFSC 2007), LPUE data were standardized by adjusting for individual vessel, vessel size, and month to see if these variables would have an effect on the trends. Estimated standardized trends were very similar to trends in nominal (unadjusted) LPUE. Following NEFSC (2009), the LPUE data presented in this assessment update have not been standardized.

Spatial patterns in fishery data

Interpreting spatial patterns in fishery data is important, especially for managing sessile and unproductive organisms like ocean quahogs. The ocean quahog stock is a complicated spatial mosaic, with scattered productive and profitable areas where abundance is high and where fishing mortality tends to be concentrated. The size of a productive ocean quahog bed appears to be less than the size of a ten-minute square (TMSQ, 10 minutes longitude x 10 minutes latitude \cong 100 nm²), which is the smallest unit used to report fishing location in logbooks.

Spatial patterns in landings, effort, and LPUE reflect a shift in the distribution of the fishery northward and offshore over time. During the 1980s, nearly all of the landings and fishing effort were from the southern DMV and NJ regions (Figure 12; Tables 6 and 7). Landings are assigned to assessment region by statistical area (Figure 13). As LPUE declined in DMV and NJ, fishing effort and landings shifted to the LI and SNE regions. In 1980, 35% of landings were from DMV, 65% were from NJ, and less than one percent was from LI. In 1990, 18% of landings were from DMV, 74% of landings were from NJ, 4% were from LI, and 4% were from SNE. During 2011, 2% of landings were from DMV, 13% were from NJ, 72% were from LI and 12% were from SNE. Percentage of fishing effort by region mirrors the percentage of landings, as vessels

have adjusted fishing locations fast enough to keep the LPUE for the stock as a whole stable (Table 4).

Most ocean quahog boats have fished in two assessment regions over the course of a year since 2003 (Figure 14), and about 10% of vessels fished in three regions. If they only fished in one region (between 20% and 40% of vessels in any one year), it was most likely to be LI. Vessels that fished in two regions were most likely to be in the LI and SNE regions. Interestingly, in 2011 more vessels fished in NJ and LI than SNE and LI.

Fishery data by ten-minute square (TMSQ)

Vessels that fish for ocean quahogs in the EEZ are required to report landings and fishing effort by TMSQ for each trip in mandatory logbooks. TMSQ are formed by dividing one-degree squares into six columns and six rows that are 10' wide. Columns are numbered 1-6 counting from west to east, and the column number is given in the TMSQ name before the row number. Rows are numbered 1-6 counting from north to south. Thus, TMSQ 436523 is the ten-minute square whose southeast corner is at 43°30' N and 65°40' E.

Landings during 1980-1990 were concentrated in a few TMSQ that were primarily in the south and relatively inshore (Figure 15). Over time, TMSQ with highest landings shifted offshore and north. Landings during 2001-2011 were concentrated in the LI region.

Fishing effort was concentrated in a few southern TMSQ during 1980-1990, with several adjacent TMSQ in the DMV assessment region having effort levels higher than 1,000 hours per year, and appreciable fishing effort south of 38°N (Figure 16). Fishing effort spread into additional offshore and northern TMSQ during 1991-2000. After 1995, there were few or no TMSQ with effort levels above 1000 hours per year.

Like landings and effort, LPUE was high inshore and south during 1980-1990, with at least ten TMSQ that had LPUE \geq 200 ITQ bushels per hour fished (Figure 17). LPUE in the area below 40°S was generally high. During 1991-1995, LPUE declined in the south and fishing effort spread northward above the 40° line into the LI region, where LPUE was still over 200 bushels per hour in many TMSQ. During 1996-2005, the fishery continued to move northward into the SNE region where catches were profitable. By the 2006-2011 time period, LPUE was less than 100 ITQ bushels per hour in all TMSQ below 39°S where fishing was reported by more than three vessels.

Trends for important TMSQ

Trends in landings, fishing effort, and LPUE during 1980-2011 were plotted for individual TMSQ that were considered important to the fishery. "Important" TMSQ include those where landings were greater than 100,000 bushels in one year during the time period, so even if they are no longer fished or if they have not been fished until recently, they were important to the fishery at some time during 1980-2011. These TMSQ plots have been divided up into assessment regions (Figure 18A-D).

Trends in LPUE for individual TMSQ tend to be high during the first years of exploitation and then tend to decline as effort, annual landings, and cumulative landings increase over time.

Unlike LPUE, which is highest in the first years of exploitation, landings and fishing effort tend to peak after 5-10 years of exploitation as LPUE stays high and then begins to decrease as beds are fished down. Decreasing trends in LPUE appear strongest in the DMV and NJ areas with the longest history of exploitation. However, in some important TMSQ where fishing effort and landings have declined over the years (e.g. TMSQ 387314 and 387443 in DMV, and 407336 in NJ), LPUE appears to have increased slightly recently under light fishing effort.

Bycatch and discard

Discard of ocean quahogs occurs at a minimal level, since the bars of the dredge are set to retain quahogs of the desired size and let the smaller ones fall through. Some incidental mortality occurs when the dredge breaks clams while fishing that are not retained. Based on Murawski and Serchuk (1989), it was assumed starting with the 2004 assessment that incidental mortality rates are $\leq 5\%$ for ocean quahogs damaged during fishing but not handled on deck. As in previous assessments, fishing mortality and other stock assessment calculations in this report assume 5% incidental mortality rates (i.e., landings $\times 1.05$ = assumed catch).

A small amount of bycatch of ocean quahog occurs in the Atlantic surfclam fishery, although there is strong incentive not to fish in areas where both species occur: mixed loads of surfclams and ocean quahogs are not acceptable to processors, and it is not practical to sort catches at sea. Fisheries Observers were aboard 16 surfclam trips between 2004 and 2008, and reported discarded ocean quahogs averaged about 100 pounds per surfclam trip. Off DMV and SVA in the southern end of the ocean quahog's range, survey catches including both surfclam and ocean quahog have become more common in recent years, as surfclams have shifted towards deeper water in response to warm water conditions (Weinberg 2005). This may change discard patterns in the future if processors continue to refuse mixed catches.

Bycatch and discard of ocean quahogs in other fisheries are zero. Ocean quahogs are not vulnerable to bottom trawls, scallop dredges (they are too deep in the sediment), gillnets, or hook and line gear.

Data from 30 observed ocean quahog trips indicate bycatch by the ocean quahog fishery includes mainly scallops (mean of 855 pounds per trip), skates (mean of 331 pounds per trip), monkfish (mean of 92 pounds per trip), unclassified snails (mean of 43 pounds per trip), spiny dogfish (mean of 22 pounds per trip), and crabs (mean of 30 pounds per trip). Summer flounder, ocean pout, longhorn sculpin, and smooth dogfish are caught incidentally at a mean of less than five pounds per trip. Fourspot flounder, striped sea robin, windowpane flounder, witch flounder, American lobster, northern sea robin, sea raven, black sea bass, red hake, and silver hake are caught incidentally at a mean of less than one pound per trip.

Commercial size selectivity

The commercial fishery selectivity curve used in this assessment update (Figure 2) is from Thorarindottir and Jacobson (2005), who estimated selectivity of commercial dredges that harvest ocean quahogs off Iceland. Based on this commercial selectivity curve

$$s_L = 1 / (1 + e^{7.63 - 0.105L})$$

where L is shell length in mm, and about 10%, 50%, and 90% of ocean quahogs are available to the fishery at 51, 72, and 93 mm SL (9, 28, and 86 years, based on the growth curve for MAB).

Dredges and towing speeds used in the US fishery are very similar to those used in the selectivity experiments. The dredge used for selectivity experiments was 24 ft (7.35 m) in length, 5 ft (1.5 m) high, and 12 ft (3.65 m) wide. The cutting blade was 10 ft (3.05 m) wide and set to penetrate sediments to a depth of 3 in (8 cm). The dredge was made of steel bars with intervening spaces of 1¼ in (3.5 cm), and was towed at about 2.1 knots (3.9 km h⁻¹). The vessel used for the experiment pumped water to jets on the dredge at about 109 psi (7.5 bars), similar to many vessels in the US fishery. Fishery selectivity curves are used in tracking trends in fishable biomass, estimating fishing mortality, and calculating biological reference points.

Commercial size-composition data

Commercial length composition data are collected by port agents from landed ocean quahogs (Table 8). The data indicate that the size composition of ocean quahogs harvested in all regions can vary from year to year due to both population dynamics of the ocean quahogs and fishing behavior (Figure 19A-D). The largest quahogs have been harvested recently from the DMV and NJ regions; those from the LI and SNE regions are about a centimeter smaller on average.

Smaller ocean quahogs were landed from the SNE and LI regions during 1997-2000 due to vessels targeting specific beds of small ocean quahogs with a high meat yield. When the port sample length frequencies from the LI and SNE regions are plotted together for recent years (Figure 20A-B), it is possible to see the smaller landed ocean quahogs in 1999.

FISHERY INDEPENDENT DATA

NEFSC clam survey

NEFSC clam survey data from 1982 through 2011 were used for this assessment update. From 1978 to 1997, clam surveys were conducted during the summer on an irregular schedule; since 1999 they have been conducted regularly every three years. Clam survey data prior to 1982 were not used for this assessment update because the surveys were done during different seasons, used other sampling equipment, or are otherwise inconsistent with the rest of the time series. Since 1982 the surveys have been conducted from the R/V *Delaware II* using a standard NEFSC hydraulic dredge with a submersible pump, a 152 cm (60 in) blade, and a 5.08 cm (2 in) mesh liner to retain small ocean quahogs and Atlantic surfclams. The survey dredge differs from commercial dredges in that it is smaller (5 ft blade instead of 8 to 12.5 ft), has a small-mesh liner, and the pump is mounted on the dredge instead of the deck of the vessel. The survey dredge retains ocean quahogs as small as 50 mm SL (size selectivity described below). Changes in ship construction, winch design, winch speed, and pump voltage that may have affected survey dredge efficiency are summarized in Table A7 of NEFSC (2004). Each of these factors has been constant since the 2002 survey.

NEFSC clam surveys are organized around NEFSC shellfish strata and stock assessment regions (Figure 1). Most ITQ ocean quahog landings originate from the area covered by the survey. The survey did not cover GBK during 1982, 1983, 1984, or 2005, and individual strata in other areas

were sometimes missed. Strata not sampled during a particular survey are “filled” for assessment purposes by borrowing data from the same stratum in the previous and/or next survey, if data are available (Table 9). Survey data are never borrowed from surveys further back than the previous survey or beyond the next survey.

Surveys follow a stratified random sampling design, allocating a pre-determined number of tows to each stratum. The data used to measure trends in ocean quahog abundance are only from stations that were random or nearly random. A small number of nearly random tows are added to some surveys in a quasi-random fashion to ensure that important areas are sampled well. Non-random stations are occupied for a variety of purposes, such as repeat tows and depletion experiment setups, but are not used to estimate relative trends in ocean quahog abundance.

A standard tow is nominally 0.125 nm (232 m) in length (5 minutes towing at a speed of 1.5 knots). However, sensor data indicate actual tow lengths depend on depth and are generally slightly longer than 0.125 nm (Weinberg et al. 2002).

Occasionally, randomly selected stations are too rocky or rough to tow. Beginning in 1999, these cases trigger a search for fishable ground in the vicinity (0.5 nm) of the original station (NEFSC 2004). If no fishable ground is located, the station is given a special code (SHG=151) and the research vessel moves on to the next station. The proportion of random stations that cannot be fished is an estimate of the proportion of habitat in a stratum or region that is not suitable habitat for ocean quahog. These estimates of the proportion of unsuitable habitat are used for calculating ocean quahog swept-area biomass.

Following all successful survey tows, all ocean quahogs and Atlantic surfclams in the survey dredge are counted, and shell length is measured to the nearest mm. A few very large catches are subsampled. Mean meat weight (kg) per tow is computed with shell length-meat weight (SLMW) equations from NEFSC (2004; Table 1).

Survey tow distance and gear performance based on sensor data

Since the 1997 survey, sensors attached to the survey dredge have been used to monitor depth, temperature, differential pressure, voltage, frequency and amperage of power supplied to the dredge, *x*-tilt (port-starboard angle), and *y*-tilt (fore-aft angle, effectively the “angle of attack” of the dredge) during survey fishing operations. At the same time, sensors on board the ship monitor electrical frequency, GPS position, vessel bearing, and vessel speed. Most of the sensor data are averaged and recorded at 1-second intervals.

Differential pressure, amperage, and *y*-tilt can be particularly important. Differential pressure is the pressure of water pumped through jets in front of the dredge blade to loosen the sediments. Amperage measures the work done by the pump in moving water through the jets. If there is a blockage at the pump entrance, then both amperage and differential pressure will be low. If there is a blockage downstream from the pump, then amperage will be low and differential pressure will be high. As described below, *y*-tilt can be used to determine if the dredge is on the bottom with the blade in the sediment.

A quantitative system for identifying and eliminating tows with poor performance based on sensor data has been used since the 2005 survey. (See Appendix A2 in NEFSC 2007 and Appendix B3 in NEFSC 2009 for details on the development of this system.) Before 2005 (the 1997, 1999, and 2002 surveys), the sensor data were primarily used to get a more accurate estimate of tow distance.

Survey tow distance and gear performance in trend analysis

Sensor data are not used to calculate tow distances for long-term trend analyses because sensor data are not available prior to 1997. For trend analysis, tow distances are based on start and stop locations recorded for each tow instead of sensor data. The catch at each station is then standardized to a nominal tow distance of 0.125 nm. Tows suitable for trend analysis (gear functioned well; no problems at station) are identified using database codes, which are recorded at sea by the watch chief following each tow, based on criteria used consistently since the late 1970s. Sensor data are used, however, to calculate tow distance and monitor gear performance during tows for depletion, repeat station, and other types of experimental studies conducted since 1997.

Survey gear selectivity

A selectivity curve for ocean quahogs by the NEFSC clam dredge was estimated in NEFSC (2004). Catches by a commercial dredge with a chicken-wire mesh liner (theoretically, to retain all but the smallest juvenile ocean quahogs) during 2003, compared with survey dredge catches in the same area during 2002, served as a basis for the curve, which estimates probability of capture based on shell length (Figure 21). The resulting selectivity curve,

$$s_L = 1 / (1 + e^{8.122 - 0.119L})$$

indicates that 50% of ocean quahogs are fully available to the NEFSC dredge at about 68 mm SL, compared to about 72 mm for commercial dredges. The survey dredge tends to retain smaller ocean quahogs than commercial dredges because of the 50 mm mesh liner in the survey dredge.

Survey catch used to make stock and fishable abundance/biomass estimates

The survey catch data for ocean quahogs were adjusted to estimate relative abundance and biomass of both the whole stock and fishable stock. Abundance and length composition of the whole stock were estimated by adjusting the number of quahogs at sizes not fully selected by the survey dredge upward, by applying the survey dredge selectivity curve. Abundance and length composition for the fishable stock (quahogs large enough to be caught in a commercial dredge) were estimated by adjusting the whole stock estimates downward, by applying the commercial dredge selectivity curve. Calculations of whole stock abundance occasionally produce large estimates for very small ocean quahogs, since survey selectivity is near zero, and the number caught is adjusted upward by a large factor. Calculation of fishable abundance and biomass from ocean quahog survey data is not affected by this problem, since fishable clams are larger and the adjustments made for selectivity are small.

Depletion studies and survey dredge efficiency

Survey dredge efficiency estimates are important because they help scale relative trends to actual biomass levels in modeling, and because they can be used to estimate swept-area biomass of ocean quahogs directly. Dredge efficiency is the proportion of all ocean quahogs living in the path of the dredge that are captured (actually brought on board) by the dredge as it passes over them. This proportion serves as a “conversion factor” between the number of ocean quahogs brought up in the dredge and the actual number of ocean quahogs living in the path of the dredge. Depletion study analysis and efficiency estimates only include ocean quahogs 90 mm SL or larger, which have high selectivity (≥ 0.85) in both survey and commercial clam dredges, and reduce the effects of shell length and size selectivity on efficiency estimates.

Depletion experiments are the first step in estimating survey dredge efficiency. Depletion experiments begin with a closely-spaced group of setup tows (usually 4) taken with the survey dredge at a particular site. “Survey density” of ocean quahogs at the site is then calculated for each setup tow by dividing the catch by area swept (dredge width times the distance traveled while the dredge was effectively fishing). Mean survey density for each depletion experiment site was calculated by averaging the survey density from each setup tow. After the setup tows are completed, many tows are made in quick succession along the same path (or as close as possible) perpendicular to the setup tows, slowly depleting the ocean quahogs until a sufficient decline in catch per tow is noted (Figure 22). Vessel position is used as a proxy for dredge position during depletion experiments. One *Delaware II* ocean quahog depletion experiment has been completed in which the research vessel carried out both the setup and depletion tows, and the survey dredge efficiency was estimated directly. For the rest of the depletion experiments, the setup tows were done by the research vessel and the depletion tows were done with a commercial vessel. In these cases the estimated survey dredge efficiency (e) is:

$$e = \frac{d}{D}$$

where D is the estimated density from the Patch model (see below) and d is the mean survey density for the site.

Patch model

The Patch model (Rago et al. 2006; Hennen et al. 2012) is used to estimate three parameters for each depletion experiment: initial ocean quahog density D ; depletion dredge efficiency e ; and a measure of variance k in catch data. The model uses a gridded rectangular plot populated with simulated quahogs, with the depletion tow paths overlaid to determine (1) how many times each grid was crossed by a dredge and (2) what the underlying density would have to be to match the catches from the depletion tows. The Patch model is used in NEFSC stock assessment work for a variety of shellfish and sedentary demersal finfish. The most important characteristic of the Patch model is that the unique path of each tow is used to interpret declines in the catch, so it is not necessary to assume that the ocean quahogs mix randomly across the entire site after each depletion tow.

Some sources of uncertainty in the Patch model are variations in substrate and gear performance. For instance, the area in Long Island where two depletion experiments were conducted has a relatively thin layer of sand on top of peat. The thin layer of sand may concentrate ocean quahogs near the surface, where they are easier to catch (Pers. comm., E. Powell, University of Southern Mississippi). Also, newly replaced electrical cables (or any other gear differences) may cause changes in actual dredge efficiency if pump voltage and pressure change. Dredge efficiency is harder to estimate for ocean quahogs than surfclams because quahogs live in deeper water, where dredge position data are less reliable, and quahogs sometimes burrow out of range of the dredge. For these reasons, it is probably better to view the full set of depletion experiment dredge efficiency estimates as a distribution with an underlying mean and variance. Individual estimates and estimates for a single survey are too imprecise to be used directly in making survey-specific estimates of survey dredge efficiency. Commercial and survey selectivity estimates used in this assessment update are the same as in NEFSC (2009). New data collected during 2011 were not used, as the process required to produce new estimates is improved by the input of a working group, which will be convened for the next benchmark assessment.

Nineteen depletion experiments for ocean quahog were carried out between 1997 and 2008. Based on these experiments, the median NEFSC survey dredge efficiency is 0.169, with 90% confidence intervals of 0.154-0.285. See NEFSC (2007; 2009) for more detailed information on the efficiency estimates and individual depletion experiments, including details of the Patch model procedure, evaluating uncertainty, and sensitivity analyses.

BIOMASS AND FISHING MORTALITY

Mortality and stock biomass estimates for ocean quahog in the US EEZ are based on NEFSC clam surveys, cooperative survey studies (including depletion experiments used to measure survey dredge efficiency), fishery data, and biological data.

Survey Results

The most recent NEFSC clam survey was conducted during the summer of 2011; it was the last clam survey to use the R/V *Delaware II* as a platform before her retirement in 2012. The 2011 survey made 391 random tows and had good coverage of all regions except SVA, which has been inhabited by so few clams recently (both ocean quahogs and surfclams) that resources have been reallocated to increase coverage of other regions. Cooperative work with an industry vessel during the survey included paired tows and depletion experiments. Ocean quahogs were most abundant, based on number per 2011 survey tow, in the LI, SNE and GBK regions (Figure 23). The abundance of recruited (70+ mm SL) ocean quahogs mirrored this distribution (Figure 24), while pre-recruits (less than 70mm SL) were most abundant on GBK (Figure 25). See Appendix 2 for maps of ocean quahogs caught per tow from all surveys since 1982.

By region, survey trends in abundance and biomass of fishable quahogs have been fairly stable since the early 2000s in GBK, SNE, LI, NJ, and DMV (Figures 26 and 27; Table 10). However, out of those regions, all have seen an overall loss of both biomass and numbers between the 1980s and early 2000s except GBK, where ocean quahog fishing has not occurred (Figures 28 and 29). The survey has caught very low numbers of ocean quahogs in the SVA region since the

late 1970s. This region was probably never good habitat for ocean quahogs and was mainly surveyed for surfclams.

Even though the survey dredge does not retain them fully, the presence of ocean quahogs smaller than 40mm SL in a tow can be a good indicator of an area that will experience recruitment into the fishery in future decades. Quahogs less than 40mm will be at least double the size and 60-80 years old when they finally recruit to the fishery. On GKB, the 2011 survey caught large numbers of these small quahogs that had grown enough since 2008 to be captured more efficiently by the survey dredge (Figure 30). Looking back on the occurrence of quahogs less than 40mm in survey tows since the 1970s, the large recruitment event on GBK can be seen, as well as evidence of low-level but consistent recruitment in the LI region, which now supplies the vast majority of landed ocean quahogs (Figure 31 A-C).

Changes in survey length composition over the years provide additional information about recruitment (Figure 32 A-E). In particular, survey length composition data from DMV, NJ, SNE, and GBK show evidence of recent recruitment events in these regions. Even though the overall density of ocean quahogs is low in the DMV region, it appears juveniles can successfully settle there. The smaller mode in the 2011 DMV length composition plot averages around 72 mm SL, which translates to about 30 years old and 50% commercial selectivity. Because ocean quahogs grow so slowly, recruitment events do not become apparent for some years. The survey dredge only begins to retain ocean quahogs when they are already at least ten years old, and even then they are not fully selected.

The spatial distribution of the ocean quahog resource on the east coast of the U.S. does not appear to have changed since the fishery began in the mid 1970s (Figure 33), but over time densities have declined where the bulk of fishing has occurred (NJ and DMV). The recent recruitment of small ocean quahogs on GBK can be seen clearly in the survey data, but any trends in SNE and LI are not as clear. The extent and number of the surveys used for each time period map are not exactly the same, but each survey does attempt to cover the same area, except for some years when there was no (or incomplete) coverage of GBK.

Efficiency-corrected swept area biomass

Efficiency-corrected swept area biomass (ESB) has been estimated since 1997, when NEFSC clam surveys began collecting dredge sensor data, which are needed to estimate area swept as they determine the amount of time the dredge was fishing during a tow (Table 11). Estimates of ESB are used primarily as scaling information in stock assessment models, since they are a direct estimate of biomass. ESB estimates are only made for the fishable ocean quahog population, sizes at which selectivity is high for the survey dredge.

ESB for ocean quahog is calculated:

$$B = \frac{B'}{e}$$

where:

$$B' = \frac{\bar{\chi}A'}{a}(1 + \phi)u$$

In ESB calculations, e is the best estimate of survey dredge efficiency for ocean quahogs, $\bar{\chi}$ is mean catch in kilograms of fishable ocean quahogs per standard tow based on sensor data, A' is habitat area (nm^2), a is the area that would be covered by the 5 ft wide survey dredge during a standard tow of 0.15 nm ($0.00012405 \text{ nm}^2 \text{ tow}^{-1}$), and u converts kilograms to thousand metric tons (10^{-6}). B' is the minimum swept-area biomass prior to correction for survey dredge efficiency.

The term ϕ used in ESB calculations is the fraction of total biomass in deep water strata off LI (strata 32 and 36), SNE (strata 40, 44, 48), and GBK (strata 56, 58, 60, and 62). In 1999, special tows were made in these areas to approximate the ocean quahog biomass and improve the ESB estimates. According to NEFSC (2000), deep water strata accounted for 0%, 2%, and 13% of ocean quahog biomass in the LI, SNE, and GBK regions. Data for these deep water strata that were only sampled once are otherwise omitted in calculations and, in particular, calculation of mean catch per tow $\bar{\chi}$.

Habitat area for ocean quahogs in each region was estimated as

$$A' = Au$$

where u is the proportion of random tows in the region not precluded by rocky or rough ground (ocean quahogs are generally found in smooth sandy habitats), and A is the total area computed by summing GIS area estimates for each survey stratum in the region. Estimates for u in this assessment update are the same as in NEFSC (2007; 2009).

Mean catch per standard tow ($\bar{\chi}$) is the stratified mean catch of fishable ocean quahogs (C_i) for tow i after adjustment to standard tow distance (d) based on tow distance measurements from sensor data (d_s):

$$\chi_i = \frac{C_i d}{d_s}$$

Only random tows were used in calculations of ESB. Tows without sensor data, with gear damage, or with poor pump performance were excluded from ESB calculations. Following NEFSC (2004), and as described above, tow distance was measured for each station assuming that the dredge was fishing when the blade penetrated the substrate to a depth of at least one inch. Thus, the tow distance at each station was the sum of the distance covered while the sensor data show the dredge angle was $\leq 5.16^\circ$.

ESB-based fishing mortality estimates

Fishing mortality rates are estimated directly from the ratio of catch (landings plus an assumed 5% incidental mortality) and ESB data for each region and year (Table 12). The primary purpose for these calculations is to check on model-based fishing mortality estimates. The NEFSC clam survey provides a good approximation of average yearly biomass, because ocean quahog biomass levels change slowly as fishing, growth, and natural mortality rates are low, and the surveys occur approximately halfway through the year.

Uncertainty in ESB and ESB-based fishing mortality estimates

Variance estimates for ESB and ESB-based fishing mortality estimates are important for using and interpreting results. The formulas for estimating ESB and fishing mortality are products and ratios of constants and random variables. The random variables used in the calculations are typically non-negative and can be assumed to be approximately lognormal; therefore, we estimated uncertainty in ESB and related mortality estimates using a formula for independent lognormal variables in products and ratios (Deming 1960):

$$CV\left(\frac{ab}{c}\right) = \sqrt{CV^2(a) + CV^2(b) + CV^2(c)}$$

where $\ln(ab/c)$, $\ln(a)$, $\ln(b)$ and $\ln(c)$ are normally distributed. Distributions of the simulated products and ratios were skewed to the right and appeared lognormal.

CV estimates for terms used in ESB and related estimates were from a variety of sources, and were sometimes just educated guesses. The CV for best estimate of survey dredge efficiency (e) was 0.21, calculated by bootstrapping the median of the set of individual estimates (15,000 iterations). CVs for sensor tow distances (d), area swept per standard tow (a), total area of region (A), percent suitable habitat (u), and catch were all assumed to be 10%. The CV for area swept (a) is understood to include variance due to Doppler distance measurements and variability in fishing power during the tow due, for example, to rocky or muddy ground.

For a more detailed discussion of uncertainty in estimates of ESB and ESB-based fishing mortality, see NEFSC (2009).

KLAMZ model

KLAMZ (for complete technical description see NEFSC 2009, Appendix B6) is a forward projecting stock assessment model based on the Deriso-Schnute delay-difference equation (Deriso 1980; Schnute 1985; Quinn and Deriso 1999). The delay-difference equation is an implicitly age-structured population dynamics model that is mathematically identical to common age-structured models if fishery selectivity is “knife-edged,” somatic growth follows the von Bertalanffy equation, and natural mortality is the same for all individuals in the modeled population. Knife-edge selectivity means that as soon as an ocean quahog reaches a minimum fishable size, it is considered fully available the fishery, rather than having the selectivity increase gradually to one as the quahog grows. All ocean quahogs over the minimum size experience the same fishing mortality in the model. Natural mortality rates and growth parameters can change from year to year in the KLAMZ model, but are assumed to be the same for all individuals alive during each year. The model is implemented in AD Model Builder.

The main assumptions in the KLAMZ model for ocean quahog are: recruitment follows a “step” pattern with the possibility of moving from one level before 1993 and a different level after; fishery selectivity is knife-edged; the natural mortality rate is a constant $0.02y^{-1}$; and growth in weight can be described by a von Bertalanffy growth curve. Recruitment is assumed to follow a simple mean (estimated to be low for ocean quahogs) because no reliable recruitment index

currently exists, recruitment levels appear to be very low based on survey data, and trends in stock dynamics are driven primarily by fishing mortality.

Recruitment to the ocean quahog fishery is not knife-edged and actually occurs over a range of sizes from 50 to 95 mm SL, with the smallest quahogs having the least chance of retention in a commercial dredge (Figure 2). Under these circumstances, KLAMZ can be used to track trends in fishable biomass, not total biomass. The fishable biomass of ocean quahogs is dominated by larger individuals that are readily captured by the survey dredge. Survey data used in the KLAMZ model are mean kg per standard tow of fishable ocean quahogs.

Despite some simplifying assumptions, KLAMZ has proven to be a relatively robust model with little or no retrospective bias, which has been used successfully for a number of stocks. It provides useful estimates of long-term biomass and fishing mortality, performs well with very limited information about age and growth, and when explicitly age-structured models are difficult to apply. One of the chief reasons for the utility of the KLAMZ model is statistical simplicity. The model used for ocean quahog, for example, estimates only 2-4 parameters.

The NEFSC clam survey was changed in 2012. It is now carried out from an industry vessel using modified commercial gear, with one-third to one-half of the ocean quahog range surveyed each year rather than the whole range every three years. Future ocean quahog stock assessment models will have to be configured to handle these multiple survey areas while providing an estimate of total abundance for all areas in the terminal year. This capability already exists in the Stock Synthesis model (SS3), used in the most recent benchmark assessment for Atlantic surfclams (NEFSC 2013). SS3 uses both age (when available) and size data to provide better information about trends in individual year class strength and mortality. A revised version of KLAMZ as well as SS3 may be used for ocean quahogs in the next benchmark assessment, and estimated trends in biomass and mortality may change if it is decided SS3 makes better use of size composition data. However, age data are not available for quahogs, suggesting that changes in estimated trends would be modest.

KLAMZ model configurations

Since this is an assessment update, the KLAMZ model was run with the same configurations used for the last ocean quahog assessment (NEFSC 2009). Only data from the 2011 survey and landings since 2008 were added to the model runs.

Data used in the KLAMZ model for this assessment update were: NEFSC clam survey biomass and associated CVs for 1982-2011; efficiency corrected swept-area biomass estimates for 1997-2011; and catch during 1977-2011 (landings plus a 5% allowance for incidental mortality). LPUE data are included in the model for comparative purposes, but have no effect on model estimates. Catch data for ocean quahogs were assumed accurate and not estimated in the model. Survey kg/tow data provide the trend information for the model estimates, while the efficiency corrected ESB estimates for 1997-2011 are used for scale only (trends are ignored). ESB estimates are calculated using a scaling parameter Q which is derived from the survey dredge efficiency experiments.

NEFSC clam survey and swept-area biomass data for 1994 were omitted, because an inadvertent increase in the voltage to the dredge pump apparently increased dredge efficiency during the 1994 survey (NEFSC 2004). In addition, survey and swept area biomass data for GBK during 1982, 1983, 1984, 1989, 2002, and 2005 were also omitted because of poor survey coverage during those years.

The KLAMZ model assumes ocean quahogs recruit to the fishery (reach 70 mm SL) at about age 26. In reality, growth patterns differ among regions (Lewis et al. 2001), but there is too little age information available to use region-specific growth curves (NEFSC 2000). A growth curve for ocean quahogs from the mid-Atlantic Bight was used for DMV, NJ, LI, and SNE, and a separate GBK curve was used for that region (Lewis et al. 2001; Figure 34). Variance in instantaneous somatic growth rates is used to help estimate the initial age structure of ocean quahogs in the initial years of the model, but this constraint is unimportant because estimated age structures are stable due to assumptions about recruitment and low mortality rates.

Separate KLAMZ model runs were done for ocean quahogs in each of the DMV, NJ, LI, SNE, and GBK regions; for the stock in the exploited region (DMV, NJ, LI, and SNE regions combined); and for the entire stock (DMV, NJ, LI, SNE, and GBK regions combined) during 1977-2011. The model was not used for the SVA region because survey data are noisy and sparse.

Parameters estimated

KLAMZ models for ocean quahogs estimate four parameters by maximum likelihood and numerical optimization. The parameters estimated are logarithms of: (1) biomass at the beginning of 1977; (2) escapement biomass at the beginning of 1978 (biomass in 1977 that survived until the beginning of 1978); and (3) annual recruitment biomass (a constant for each of two time periods). Fishing mortality rates were calculated by solving the catch equation numerically (see NEFSC 2009, Appendix B6). Survey scaling parameters were calculated using a closed form maximum likelihood estimator.

Variance estimates

Variances for biomass, fishing mortality estimates, and model parameters were estimated by the delta method using exact derivatives calculated by AD Model Builder libraries, or by bootstrapping.

KLAMZ results

This portion of the assessment update is intended to be no more than a report of model results using new survey and catch data, so the model configurations used were not substantively changed from the last assessment. The only change made was to allow a second recruitment parameter to be estimated in the models for DMV, NJ, and GBK. Using this configuration, the model could estimate two separate recruitment constants: one for annual recruitment from 1977 until 1993, and another for recruitment from 1994 forward. It was not, however, necessary for the model to estimate two recruitment constants if only one recruitment constant was supported by the data. Therefore, the changes made to the model in this assessment were not considered substantive, since a one-parameter recruitment was a possible outcome of the current model configuration. Despite these changes, which were intended to improve the probability of model

convergence, KLAMZ runs for the NJ and SNE regions did not converge (which also often occurred in the previous assessment).

DMV region

The KLAMZ model for ocean quahogs in the DMV area (Figures 35 and 36) fit NEFSC survey data well, and KLAMZ biomass matched the trends in observed LPUE (LPUE data do not affect model estimates). The estimated survey scaling parameter for ESB data was $Q=0.999$, indicating that the model was able to match the scale of the observed ESB on average during 1997-2011 using the catch data and trends in NEFSC survey data. Recruitment was near zero in runs for DMV. Survey data generally indicate that recruitment has been low in DMV since 1978, although some small ocean quahogs are present.

Based on KLAMZ model results, the biomass of ocean quahogs in DMV has been in decline since 1978, but the decline has not been as steep in recent years. Estimated fishable biomass is approximately 28% of 1978 biomass.

NJ region

KLAMZ model results for ocean quahogs in the NJ area (Figure 37) fit NEFSC survey index data well even though the model did not converge. The estimated survey scaling parameter for ESB data was $Q=0.98$, indicating that the model was able to scale with the observed ESB.

Based on KLAMZ model results, the biomass of ocean quahogs in NJ has been declining steadily since 1978. Estimated fishable biomass in NJ during 2011 was approximately 38% of the estimate for 1978.

LI region

A particularly strong year class recruited to the LI fishery in the years leading up to 1994, so the recruitment step function was influential in the model runs for LI.

The current results of the model (Figure 38) show ocean quahog biomass in LI increasing steadily from 1978 until 1993, when recruitment decreased, fishing mortality increased, and biomass began to drop, following the recruitment pattern. The estimated scaling parameter for ESB data Q was 0.91, indicating that the model was able to match the observed ESB levels on average. The model fit to survey indices was good, as they also show the results of increased recruitment in the years before 1994.

The correlation between LPUE predicted by KLAMZ (based on observed catch/predicted biomass) and observed LPUE is poor, because the fishery has moved into the region fairly recently compared to the regions to the south. The vessels are able to keep the LPUE high by selecting areas of high relative density, even though the exploitation rate for the entire region is increasing.

Based on the most recent KLAMZ model results, biomass of ocean quahogs in LI has decreased since 1978. Estimated fishable biomass in LI during 2011 was approximately 78% of the estimate for 1978.

SNE region

The KLAMZ model output for SNE (Figure 39) did not fit NEFSC survey data well, but the estimated survey scaling parameter for ESB data was $Q=1.04$, indicating the model matched the scale of observed ESB levels on average during 1997-2011. The KLAMZ predicted LPUE based on catch and biomass estimates did not follow the trends of observed LPUE, but this region has a fishery pattern similar to LI, where the vessels are able to move around and keep LPUE high as the region has not been heavily exploited.

Based on KLAMZ model results, biomass of ocean quahogs in the SNE region have been declining steadily since 1978. Estimated fishable biomass in SNE during 2011 was approximately 47% of the estimate for 1978.

GBK region

The KLAMZ model for ocean quahogs in the GBK region fit NEFSC survey kg/tow data reasonably, although only six survey observations were available (Figure 40) and the recruitment step function was influential. Only four ESB observations were available for GBK, and the estimated scaling parameter was $Q=1.01$. Since there has been no ocean quahog fishing on GBK, there is no LPUE or fishing mortality estimated.

Based on KLAMZ model results, biomass of ocean quahogs on GBK has been decreasing slowly since the early 1990s, but the model fit to survey biomass was poor (Figure 41). Estimated fishable biomass during 2011 was about 6% higher than the estimate for 1978.

Exploited area and entire stock

The KLAMZ model for ocean quahogs in the exploited stock area (all regions except for GBK, where the population of ocean quahogs is not fished) fit NEFSC survey kg/tow reasonably well (Figures 42, 43, and 44). The biomass estimates from the exploited area model correlated poorly with LPUE, probably because the area modeled is so large that the fishery can easily shift its distribution within it (for example, from NJ to LI) to maintain high catch rates, whereas the single-region models containing historical ocean quahog fishing grounds (DMV, NJ) observed LPUE tracks biomass estimates well. The estimated survey scaling parameter for ESB data was $Q=1.18$, indicating that the model believes there is somewhat less biomass on average during 1997-2011 than the observed ESB biomass levels indicate.

Based on KLAMZ model results (Figure 42), biomass of ocean quahogs in the exploited area was stable from 1978 to 1993, and then began to decline when recruitment dropped and fishing mortality was relatively high. Estimated fishable biomass for the exploited area during 2011 was approximately 70% of the estimate for 1978.

Biomass estimates from the KLAMZ model for the exploited region were about 4% smaller than the sum of biomass estimates from regional KLAMZ models for DMV, NJ, LI, and SNE. Confidence intervals of the KLAMZ biomass estimates for the exploited regions were slightly narrower than the previous assessment after the addition of the latest survey data, though there is still considerable uncertainty in the estimate (Figure 45, bottom).

The biomass estimate for the whole stock (Figure 45, top) shows almost the same trend as the estimate for the exploited area, but with a more pronounced increase during the earlier period. The KLAMZ model estimates the biomass of the whole stock to be almost double the estimate for the exploited area, although the uncertainty is higher for the whole stock. The whole stock biomass is estimated to be approximately 86% of the biomass in 1978. The fit to the survey data is not as good as the exploited regions model, but the data from GBK are sparser and less consistent spatially (Figures 46, 47, and 48).

KLAMZ estimates of fishing mortality (Figure 49) for both the exploited regions and the whole stock show the same trends but on a different scale, as the model is removing the same catch from the whole stock as it is from about half the stock in the case of the exploited regions. Time-series estimates of fishing mortality for the single-region KLAMZ models, especially the regions more recently exploited like LI and SNE, have undergone significant changes over time. The fishing mortality estimates for the entire exploited area are quite stable as the fishery moves around within a large range – the same reason LPUE is often stable despite declines in biomass.

Historical retrospective analyses

KLAMZ model results from this assessment update and the previous 5 or 6 assessments (for SARC 27, in 1998, the regional biomass estimates were single points and fishing mortality was not estimated) are interesting to compare, and reflect the changes that have been made to the model configurations over the years, as well as changes in the input time series, as data is accumulated over time. Changes in survey dredge efficiency estimates have altered the scale of the biomass and fishing mortality estimates to some degree for all regions. With each passing year, more depletion experiments are done and the survey dredge efficiency estimates improve. The more recent model results have used a lower survey dredge efficiency, which translates into a higher biomass estimate. Recent changes in how recruitment has been modeled in KLAMZ (using a step function) have affected the trend for the first half of the time series for some regions.

For the DMV and NJ regions, historically fished the hardest, KLAMZ biomass estimates have been trending downward for the whole time series in every instance (Figure 50). The estimates for the LI area have been influenced by the change in recruitment assumptions (Figure 51). In the SNE region, the KLAMZ results from this assessment update show a higher biomass at the beginning of the time series and a faster rate of decline, probably due to a combination of recent low survey kg/tow data points and the decrease in dredge efficiency (Figure 51). The trends in biomass for both the exploited area and the whole stock have been affected by the recruitment step function, peaking during the year the model recruitment decreased (Figures 52 and 53).

KLAMZ estimates of fishing mortality have also changed as a result of the newer survey dredge efficiency estimates. As the biomass estimates have scaled higher, the fishing mortality estimates have scaled lower, as the same catch removed from a larger biomass translates into a lower F value. Otherwise, KLAMZ results for fishing mortality have followed the same trends over the last five assessments (Figures 54-56).

REFERENCE POINTS AND STOCK STATUS

Biological Reference Points

Biological reference points (BRPs) for fishing mortality (F) and biomass (B) are required to manage the ocean quahog stock. Targets are BRPs that represent desirable stock conditions, and thresholds are BRPs that identify undesirable stock conditions.

BRPs for US fisheries are generally linked to the theoretical maximum sustainable yield (MSY) of a stock. MSY theory is probably not applicable to ocean quahogs, because their very low productivity may preclude economically viable levels of sustained catch. Productivity is low because recruitment events have been infrequent and localized, growth is slow, and there is a long time between settlement and recruitment to the spawning population or fishable stock. Ocean quahogs have also not shown an increase in productivity from higher recruitment and faster growth as biomass has been fished down, which would be typical for a species operating under the MSY principle. For these reasons, recommended reference points are described as thresholds and targets but not as proxies for F_{MSY} or B_{MSY} related reference points.

During SARC 48 (NEFSC 2009), BRPs for ocean quahogs were revisited and changes were recommended based on the unique population dynamics of very long-lived species with low rates of adult natural mortality. The revised BRPs are peer reviewed and considered the best science, but they are not incorporated in the FMP as an amendment or framework process has not taken place. Details of the reasoning behind the BRP revisions and analysis done to explore the implications of a range of potential BRP values can be found in the last assessment (NEFSC 2009).

B_{target}

The ocean quahog biomass target used to be $B_{Target} = B_{MSY}$, which is assumed to be half the virgin biomass of the whole stock. This was revised to be half of the fishable fraction of the whole stock biomass during 1978 (considered pre-fishery). The revised biomass target value is 1.73 million metric tons of meats.

$B_{threshold}$

The biomass threshold used to be $B_{Threshold} = \frac{1}{2} B_{MSY}$ (or $\frac{1}{4}$ the virgin biomass of the whole stock). The revised $B_{threshold}$ is 40% of the 1978 whole stock biomass, which was judged to be more realistic. The revised biomass threshold value for the whole stock is 1.39 million metric tons of meats.

F_{target}

F_{Target} was originally $F_{0.1}$, and applied to the exploited area only. The revised F_{target} is a value of F below the threshold, but in an ITQ fishery this is determined by the quota set for the year and therefore at the discretion of the managers. By default it applies to the exploited area only.

$F_{threshold}$

$F_{Threshold}$ used to be $F_{25\%}$ (the fishing mortality rate that reduces lifetime egg production to 25% of its potential). It was decided to revise $F_{threshold}$ to $F_{45\%}$ (egg production at 45% of potential) as

it was better suited to unproductive stocks. The revised fishing mortality threshold for the exploited area is 0.022 y^{-1} .

	Revised BRPs	Value
<i>B</i> target	50% 1978 fishable biomass (whole stock)	1.73 million mt
<i>B</i> threshold*	40% 1978 fishable biomass (whole stock)	1.39 million mt
<i>F</i> target	$F_{\text{target}} < F_{\text{threshold}}$ (exploited area)	set by quota
<i>F</i> threshold*	$F_{45\%}$ (exploited area)	0.022 y^{-1}
	Current whole stock fishable biomass	Current exploited area <i>F</i>
	2.96 million mt	0.010 y^{-1}

STOCK STATUS

Ocean quahogs in the US EEZ are not overfished and overfishing is not occurring. Whole stock fishable biomass during 2011 was 2.96 million mt (Table 13), which is above the revised B_{target} of 1.73 million mt and the revised $B_{\text{threshold}}$ of 1.39 million mt. The fishing mortality rate during 2011 for the stock in the exploited region was $F = 0.010 \text{ y}^{-1}$ (Table 14), below the revised $F_{\text{threshold}}$ of 0.022 y^{-1} . Fishing mortality for the exploited area of the stock was also below the previous $F_{\text{threshold}}$ of 0.08 y^{-1} , and whole stock biomass was above the previous $B_{\text{threshold}}$ of 0.89 million mt.

PROJECTIONS

Ocean quahog biomass, fishing mortality, and catch were projected through 2026 using the KLAMZ model and associated bootstrap/projection software (Figures 57A-C). The projections were run for three scenarios with different assumptions about catch. The probabilities of overfishing or overfished status were calculated for each year. Bootstrap KLAMZ model runs were used as starting points for five thousand projections, to accommodate uncertainty about current biomass. For the first “status quo catch” scenario, yearly fishery removals were fixed at 15,400 mt. For the second “quota catch” scenario, yearly fishery removals were fixed at the level of the current quota, 24,190 mt. The third scenario fixed annual fishing mortality at the overfishing threshold, $F_{45\%}$ or 0.022 y^{-1} . Recruitment was constant and derived from KLAMZ base runs for recent years. Status quo catch was determined by averaging ITQ ocean quahog landings from 2006 through 2011. Catches used in simulations were reported landings with 5% added to account for incidental mortality. All three scenarios were run for both the whole stock (all EEZ ocean quahogs with the exception of Maine) and the exploited area (SVA, DMV, NJ, LI, and SNE). All projections were started in 2011, the last year with biomass estimates from stock assessment models for ocean quahogs and complete landings data.

Stochastic models were used as this approach allows confidence intervals to be calculated, and give almost identical median estimates as the single set of estimates from deterministic models.

A single value of natural mortality (0.02) was used for all projections for this update, instead of running three different sets of projections using 0.015, 0.02, and 0.025. The different values of M affect both the starting biomass and the rate of change in the population over the length of the projection, but in the case of ocean quahogs these rates are low and the differences between runs were predictable. For this update, three different fishery scenarios were projected instead of four, as the minimum landings scenario from the last assessment was determined not to be useful. However, the projections made for this assessment update cover a longer period of time (fifteen years versus five).

The current fishing mortality threshold $F_{45\%} = 0.022y^{-1}$ applies to the exploited region only. The current biomass threshold of 1,390 mt (40% of the 1978 biomass) applies to the whole stock. The recent opening of the GBK region to surfclam and potential ocean quahog fishing may necessitate changes to the way biological reference points are applied to determine stock status. Even though there has been no ocean quahog fishing on GBK since the opening, it is possible there may not be an “exploited” subarea of the ocean quahog range in the EEZ in the future.

The status quo catch scenario (Figure 57A) represents the lowest catch level projected. At this catch level there is little chance of the whole stock becoming overfished between 2011 and 2026 (Table 15). However, the bootstrap estimates indicate there is a very small probability that overfishing may occur in the exploited area by 2020, even at this lowest level of catch.

The full quota catch scenario (Figure 57B) would also probably not lead to the stock becoming overfished, but would generate a more rapid decrease in biomass leading to overfishing in the exploited area (the probability of overfishing is estimated to be 50% by 2024).

The overfishing threshold scenario by definition had overfishing occurring in all years (Figure 57C). The probability of the whole stock reaching an overfished state after 15 years of fishing at $0.022y^{-1}$ is still small, but biomass would be reduced considerably and catches would begin to decrease around 2020.

In all three scenarios, ocean quahog biomass is projected to decline continuously through 2026. If recruitment levels remain as low as they have been in recent years and fishing continues at the status quo rate, there is a reasonable chance there will be overfishing in the exploited area and the whole stock will lose 20% of its biomass within the next ten or twelve years.

REFERENCES

- Butler PG, Wanamaker AD Jr, Scourse JD, Richardson A, Reynolds DJ. Variability of marine climate on the North Icelandic Shelf in a 1357-year proxy archive based on growth increments in the bivalve *Arctica islandica*. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* (2012). Doi:10.1016/j.palaeo.2012.01.016 (in press).
- Cargnelli LM, Griesbach SJ, Packer DB, Weissberger E. 1999. Essential fish habitat source document: Ocean quahog, *Arctica islandica*, life history and habitat characteristics. NOAA Technical Memorandum NMFS-NE-148.

- Dahlgren T, Weinberg J, Halanych K. 2000. Phylogeny of the ocean quahog (*Arctica islandica*): influences of paleoclimate on genetic diversity and species range. *Mar. Bio.* 137: 487-495.
- Deming WE. 1960. *Sample design in business research*. New York: Wiley & Sons, Inc.; 517 p.
- Deriso RB. 1980. Harvesting strategies and parameter estimation for an age-structured model. *Can J Fish Aquat Sci.* 37: 268-282.
- Fritz L. 1991. Seasonal change, morphometrics, growth and sex ratio of the ocean quahog, *Arctica islandica* (Linnaeus, 1767) off New Jersey, U.S.A. *J Shellfish Res.* 10:79-88.
- Harding JM, King SE, Powell EN, Mann R. 2008. Decadal trends in age structure and recruitment patterns of quahogs *Arctica islandica* from the mid-Atlantic bight in relation to water temperature. *J Shellfish Res.* 27(4): 667-690.
- Hennen D, Jacobson LD, Tang J. 2012. Accuracy of the patch model used to estimate density and capture efficiency in depletion experiments for sessile invertebrates and fish. *ICES J Mar Sci.* 69:240-249.
- Jones DS. 1980. Annual cycle of shell growth increment formation in two continental shelf bivalves and its paleoecological significance. *Paleobiology.* 6:331-340.
- Kilada RW, Campana SE, Roddick D. 2007. Validated age, growth, and mortality estimates of the ocean quahog (*Arctica islandica*) in the western Atlantic. *ICES J Mar Sci.* 64:31-38.
- Lewis C, Weinberg J, Davis C. 2001. Population structure and recruitment of *Arctica islandica* (Bivalvia) on Georges Bank from 1980-1999. *J Shellfish Res.* 20: 1135-1144.
- Murawski SA, Serchuk FM. 1979. Shell length-meat weight relationships of ocean quahogs, *Arctica islandica*, from the middle Atlantic shelf. *Proc Nat Shell Assoc.* 69:40-46.
- Murawski SA, Ropes JW, Serchuk FM. 1982. Growth of the ocean quahog, *Arctica islandica*, in the middle Atlantic Bight. *Fish Bull.* 80: 21-34.
- Murawski SA, Serchuk FM. 1989. Mechanized shellfish harvesting and its management: the offshore clam fishery of the eastern United States. *In: Caddy JF, ed. Marine Invertebrate Fisheries: Their Assessment and Management*. New York: Wiley & Sons, Inc.; p. 479-506.
- Northeast Fisheries Science Center (NEFSC). 2000. Stock assessment for ocean quahogs. *In: Report of the 31st Northeast Regional Stock Assessment Workshop (31st SAW)*. Northeast Fish Sci Cent Ref Doc. 00-15.
- NEFSC. 2004. Stock assessment for ocean quahogs. *In: Report of the 38th Northeast Regional Stock Assessment Workshop (38th SAW)*. Northeast Fish Sci Cent Ref Doc. 04-03.

- NEFSC. 2007. Stock assessment for ocean quahogs. *In*: Report of the 44th Northeast Regional Stock Assessment Workshop (44th SAW). Northeast Fish Sci Cent Ref Doc. 07-10.
- NEFSC. 2009. 48th Northeast Regional Stock Assessment Workshop (48th SAW) Assessment Report. Northeast Fish Sci Cent Ref Doc. 09-15; 834 p.
- NEFSC. 2013. 56th Northeast Regional Stock Assessment Workshop (56th SAW) Assessment Report. Northeast Fish Sci Cent Ref Doc. In preparation.
- Powell E, Mann R. 2005. Evidence of recent recruitment in the ocean quahog *Arctica islandica* in the Mid-Atlantic Bight. *J Shellfish Res.* 24: 517-530.
- Quinn T, Deriso RB. 1999. Quantitative fish dynamics. Oxford, UK: Oxford University Press; 560 p.
- Rago PJ, Weinberg JR, Weidman C. 2006. A spatial model to estimate gear efficiency and animal density from depletion experiments. *Can J Fish Aquat Sci.* 63(10):2377-2388.
- Ropes JW, Murawski SA, Serchuk FM. 1984. Size, age, sexual maturity and sex ratio in ocean quahogs, *Arctica islandica* Linne, off Long Island, New York. *Fish Bull.* 82: 253-267.
- Schnute J. 1985. A general theory for analysis of catch and effort data. *Can J Fish Aquat Sci.* 42: 414-429.
- Serchuk F, Murawski S. 1997. The offshore molluscan resources of the Northeastern Coast of the United States: surfclams, ocean quahogs, and sea scallops. *In*: MacKenzie CL Jr, Burrell VG Jr, Rosenfield A, Hobart WL, eds. The History, Present Condition, and Future of the Molluscan Fisheries of North and Central America and Europe, Volume 1, Atlantic and Gulf Coasts. NOAA Tech Rep. 127; p 45-62.
- Steingrimsson SA, Thorarinsdottir G. 1995. Age structure, growth and size at sexual maturity in ocean quahog *Arctica islandica* (Mollusca: Bivalvia), off NW-Iceland. *ICES CM* 1995/K: 54
- Strahl S, Philipp E, Brey T, Broeg K, Abele D. 2007. Physiological aging in the Icelandic population of the ocean quahog *Arctica islandica*. *Aquat Biol.* 1:77-83.
- Theroux RB, Wigley RL. 1983. Distribution and abundance of east coast bivalve mollusks based on specimens in the National Marine fisheries Service Woods Hole collection. NOAA Tech Rep NMFS SSRF 768; 172 p.
- Thompson I, Jones DS, Dreibelbis D. 1980. Annual internal growth banding and life history of the ocean quahog *Arctica islandica* (Mollusca: Bivalvia). *Mar Biol.* 57: 25-34.

- Thorarinsdottir GG, Einarsson ST. 1996. Distribution, abundance, population structure and meat yield of the ocean quahog, *Arctica islandica*, in Icelandic waters. *J Mar Biol Assoc UK*. 76: 1107-1114.
- Thorarinsdottir GG, Jacobson LD. 2005. Fishery biology and biological reference points for management of ocean quahogs (*Arctica islandica*) off Iceland. *Fish Res*. 75: 97-106.
- Wanamaker AD, Heinemeier J, Scourse JD, Richardson CA, Butler PG, Eiriksson J, Knudsen KL. 2008. Very long-lived mollusks confirm 17th century AD tephra-based radiocarbon reservoir age for north Atlantic shelf waters. *Radiocarbon*. 50(3):399-412
- Weinberg J. 2005. Bathymetric shift in the distribution of Atlantic surfclams: response to warmer ocean temperature. *ICES J Mar Sci*. 62: 1444-1453.
- Weinberg JR, Rago P, Wakefield W, Keith C. 2002. Estimation of tow distance and spatial heterogeneity using data from inclinometer sensors: an example using a clam survey dredge. *Fish Res*. 55: 49-61.

Table 1. Parameter estimates for the relationship between shell length (L) in millimeters and meat weight (W) in grams for ocean quahogs used in this assessment update. The equation for the relationship is $W=e^{\alpha}L^{\beta}$.

Region	Alpha	Beta
GBK	-8.969073	2.767282
SNE	-9.124283	2.774989
LI	-9.233646	2.822474
NJ	-9.847183	2.94954
DMV	-9.042313	2.787987
SVA	-9.042313	2.787987

Table 2. Annual ocean quahog landings (including Maine) and quotas in thousands of metric tons of meats.

Year	EEZ landings (Logbook)	EEZ Quota	EEZ Landings / Quota (%)
1980	13.41	15.88	0.84
1981	13.10	18.14	0.72
1982	14.23	18.14	0.78
1983	14.59	18.14	0.80
1984	17.98	18.14	0.99
1985	20.73	22.23	0.93
1986	18.90	27.22	0.69
1987	21.51	27.22	0.79
1988	20.27	27.22	0.74
1989	22.36	23.59	0.95
1990	20.97	24.04	0.87
1991	22.06	24.04	0.92
1992	22.48	24.04	0.93
1993	21.88	24.49	0.89
1994	20.99	24.49	0.86
1995	21.11	22.23	0.95
1996	20.06	20.18	0.99
1997	19.63	19.58	1.00
1998	17.90	18.14	0.99
1999	17.38	20.41	0.85
2000	14.72	20.41	0.72
2001	17.07	20.41	0.84
2002	17.95	20.41	0.88
2003	18.82	20.41	0.92
2004	17.66	22.68	0.78
2005	13.64	24.19	0.56
2006	14.27	24.19	0.59
2007	15.57	24.19	0.64
2008	15.74	24.19	0.65
2009	15.87	24.19	0.66
2010	16.27	24.19	0.67
2011	14.34	24.19	0.59

Table 3. Real and nominal prices for ocean quahog based on dealer data. Average price was computed as total revenues divided by total landed meat weight during each year. The consumer price index (CPI) used to convert nominal dollars to 2010 equivalent dollars is from the Bureau of Economic Analysis Table 1.1.9 (www.bea.gov/iTable/iTable.cfm?ReqID=9&step=1).

Year	Conversion factor for 2010 dollars	ITQ fishery, price per lb		Maine fishery, price per lb	
		Nominal price	Real price (2010 dollars)	Nominal price	Real price (2010 dollars)
1982	0.50	0.31	0.62		
1983	0.52	0.31	0.60		
1984	0.54	0.31	0.58	0.78	1.45
1985	0.56	0.31	0.56		
1986	0.57	0.3	0.53	1.75	3.08
1987	0.58	0.29	0.50	2.3	3.94
1988	0.60	0.29	0.48	1.9	3.15
1989	0.63	0.29	0.46	2.72	4.34
1990	0.65	0.32	0.49	2.7	4.15
1991	0.67	0.34	0.50	4.1	6.08
1992	0.69	0.36	0.52	4.07	5.90
1993	0.71	0.4	0.57	3.58	5.08
1994	0.72	0.38	0.53	3.83	5.32
1995	0.74	0.4	0.54	3.46	4.71
1996	0.75	0.41	0.55	3.1	4.14
1997	0.76	0.42	0.55	2.62	3.44
1998	0.77	0.42	0.54	2.5	3.24
1999	0.78	0.42	0.54	2.75	3.51
2000	0.80	0.43	0.54	2.74	3.43
2001	0.82	0.55	0.67	3.23	3.95
2002	0.83	0.54	0.65	3.69	4.44
2003	0.85	0.53	0.62	3.75	4.42
2004	0.87	0.52	0.60	3.79	4.35
2005	0.90	0.51	0.57	3.6	4.00
2006	0.93	0.51	0.55	3.23	3.47
2007	0.96	0.52	0.54	3.16	3.30
2008	0.98	0.54	0.55	3.29	3.36
2009	0.99	0.60	0.61	3.33	3.37
2010	1.00	0.64	0.64	3.29	3.29
2011	1.02	0.65	0.64	3.38	3.31

Table 4. Ocean quahog fishing effort in days (24 hours) fished by stock assessment region in the US EEZ based on logbook data.

YEAR	DMV	LI	NJ	SNE	SVA	UNK	Total ITQ
1980	279.8	1.3	600.6			11.9	893.6
1981	238.6	0.3	656.2		3.0	2.0	900.1
1982	296.4		606.4		0.3	3.0	906.1
1983	297.1	2.1	580.7	63.9		2.3	946.2
1984	462.8		644.9	105.1	0.6	51.3	1,264.7
1985	419.1	3.6	745.4	86.1	8.5	123.1	1,385.9
1986	510.8	15.0	598.3	47.4		42.1	1,213.7
1987	659.1	33.6	612.4	55.8		2.0	1,362.9
1988	795.8	25.6	483.2	68.3	2.7	2.7	1,378.3
1989	505.2	33.2	1,010.9	97.0		2.1	1,648.3
1990	340.2	53.4	1,222.0	76.6	1.0		1,693.3
1991	502.0	76.8	1,266.6	59.7			1,905.1
1992	229.7	547.8	666.6	81.8			1,526.0
1993	192.6	536.8	1,060.7	74.3			1,864.4
1994	94.2	798.5	855.9	86.8		2.4	1,837.7
1995	67.5	667.3	566.6	356.7			1,658.1
1996	63.4	426.6	389.2	494.4		2.2	1,375.8
1997	114.3	345.6	390.9	563.1			1,414.0
1998	134.4	437.9	291.0	443.3		3.3	1,309.8
1999	108.1	380.5	317.6	510.7		3.7	1,320.7
2000	104.9	294.6	330.0	439.3		536.4	1,705.2
2001	88.1	173.5	451.7	470.9		765.7	1,949.9
2002	178.8	483.5	278.4	324.9		702.7	1,968.3
2003	109.1	671.4	447.9	191.7		4.5	1,424.6
2004	104.0	607.6	329.4	276.8			1,317.7
2005	143.5	521.6	82.2	168.4		1.9	917.6
2006	75.5	604.9	57.8	138.1			876.2
2007	14.4	650.6	155.0	178.1			998.1
2008	39.8	634.6	197.4	173.6			1,045.5
2009	32.9	453.5	223.1	704.1		3.3	1,416.8
2010	71.2	533.6	267.3	212.3		0.8	1,085.3
2011	40.4	589.2	199.0	155.5			984.0

Table 5. Ocean quahog landings per unit effort (LPUE, total bushels landed / total hours fished)
based on logbook data for all vessels operating in the US EEZ.

Year	DMV	LI	NJ	SNE	SVA	UNK	Mean ITQ
1980	146.77	40.38	124.98			132.50	111.16
1981	142.13	122.67	130.08		174.23	159.47	145.71
1982	141.85		116.95		178.29	191.38	157.12
1983	151.94	90.85	130.58	91.15		124.76	117.86
1984	138.84		114.23	64.47	89.27	117.42	104.84
1985	157.01	93.06	138.53	68.70	133.82	124.53	119.27
1986	136.96	235.71	143.66	107.19		137.42	152.19
1987	155.96	277.37	135.65	111.80		133.26	162.81
1988	130.04	197.79	143.84	102.12	145.73	83.04	133.76
1989	124.60	155.42	120.76	135.01		121.67	131.49
1990	118.03	139.78	141.08	109.64	119.20		125.55
1991	98.55	153.60	112.84	129.61			123.65
1992	95.53	194.64	117.23	134.10			135.37
1993	105.10	158.24	108.15	128.26			124.94
1994	91.56	145.34	83.71	122.20		80.48	104.66
1995	90.03	138.04	90.76	153.49			118.08
1996	106.12	136.91	113.43	171.40		108.40	127.25
1997	85.04	137.79	97.65	150.84			117.83
1998	101.99	132.81	124.56	147.30		83.11	117.95
1999	100.38	143.62	130.38	123.57		51.38	109.86
2000	89.64	129.60	107.88	111.65		76.87	103.13
2001	81.19	117.70	103.19	99.81		107.85	101.95
2002	99.90	132.90	93.16	101.81		117.00	108.95
2003	77.81	140.11	86.58	96.14		145.94	109.31
2004	69.90	171.77	98.51	117.30			114.37
2005	63.35	148.62	94.91	113.02		94.93	102.97
2006	65.57	156.32	93.65	163.02			119.64
2007	62.49	155.74	89.91	134.76			110.73
2008	63.22	160.93	82.17	138.20			111.13
2009	54.60	195.13	98.79	136.29		5.00	97.96
2010	51.81	178.59	77.10	227.15			133.66
2011	68.62	240.13	86.16	157.09			138.00

Table 6. Ocean quahog landings (mt meats) by stock assessment region as reported in logbooks by the ITQ fishery.

YEAR	SVA	DMV	NJ	LI	SNE	GBK	UNK	Total
1980		4,232	7,748	6			147	12,133
1981	56	3,637	8,402	3			23	12,120
1982	6	4,605	8,532				63	13,205
1983		5,396	8,249	21	629		291	14,586
1984	6	7,171	8,851		822		1,125	17,975
1985	160	7,200	10,676	40	693		1,956	20,726
1986		8,237	9,059	396	562		649	18,902
1987		10,540	9,070	1,180	696		27	21,514
1988	42	11,715	7,014	640	841		20	20,273
1989		6,439	14,100	605	1,196		20	22,359
1990	14	3,685	15,590	739	934			20,962
1991		4,839	14,575	1,674	865			21,953
1992		2,378	6,942	11,940	1,143			22,402
1993		1,953	10,205	8,642	1,020			21,820
1994		992	6,938	12,015	954		22	20,921
1995		699	5,356	9,526	5,412			20,994
1996		736	4,864	5,943	8,350		26	19,919
1997		1,072	4,229	5,141	8,968			19,410
1998		1,365	2,684	6,856	6,736		39	17,679
1999		1,090	3,038	6,329	6,618		27	17,102
2000		1,048	3,286	4,745	5,083	49	695	14,905
2001		864	4,544	2,417	4,632	13	4,765	17,235
2002		1,732	2,781	9,113	3,884		635	18,144
2003		896	3,683	11,626	2,177		73	18,456
2004		624	2,761	10,690	3,273			17,348
2005		910	669	9,714	2,021		19	13,334
2006		494	467	11,101	1,847			13,909
2007		100	1,566	11,290	2,311			15,267
2008		270	1,724	11,290	2,255			15,539
2009		211	2,424	8,626	4,594		18	15,873
2010		428	2,315	9,882	3,467		14	16,106
2011		290	1,868	10,235	1,748			14,141

Table 7. Ocean quahog landings by stock assessment region as reported in logbooks for the ITQ fishery. Figures are in thousands of ITQ bushels.

YEAR	SVA	DMV	NJ	LI	SNE	GBK	UNK	Total
1980		933	1,708	1			32	2,675
1981	12	802	1,852	1			5	2,672
1982	1	1,015	1,881				14	2,911
1983		1,190	1,819	5	139		64	3,216
1984	1	1,581	1,951		181		248	3,963
1985	35	1,587	2,354	9	153		431	4,569
1986		1,816	1,997	87	124		143	4,167
1987		2,324	2,000	260	153		6	4,743
1988	9	2,583	1,546	141	185		4	4,469
1989		1,420	3,108	133	264		4	4,929
1990	3	812	3,437	163	206			4,621
1991		1,067	3,213	369	191			4,840
1992		524	1,530	2,632	252			4,939
1993		431	2,250	1,905	225			4,810
1994		219	1,530	2,649	210		5	4,612
1995		154	1,181	2,100	1,193			4,628
1996		162	1,072	1,310	1,841		6	4,391
1997		236	932	1,133	1,977			4,279
1998		301	592	1,511	1,485		9	3,898
1999		240	670	1,395	1,459		6	3,770
2000		231	724	1,046	1,121	11	153	3,286
2001		190	1,002	533	1,021	3	1,050	3,800
2002		382	613	2,009	856		140	4,000
2003		198	812	2,563	480		16	4,069
2004		138	609	2,357	722			3,825
2005		201	148	2,142	446		4	2,940
2006		109	103	2,447	407			3,066
2007		22	345	2,489	510			3,366
2008		59	380	2,489	497			3,426
2009		46	534	1,902	1,013		4	3,499
2010		94	510	2,179	764		3	3,551
2011		64	412	2,256	385			3,117

Table 8. Number of quahogs measured, number of trips sampled, percentage of trips sampled, and number of quahogs measured per bushel landed by year and region, from port samples.

Region	Year	Quahogs measured	Trips sampled	% of trips sampled	Samples per bushel landed
SNE	1996	30	1	0.12	0.00002
	1997	310	10	1.20	0.00016
	1998	796	25	3.88	0.00054
	1999	634	21	2.67	0.00043
	2000	822	27	4.12	0.00073
	2001	761	25	3.84	0.00074
	2002	1353	42	7.18	0.00158
	2003	606	20	6.31	0.00126
	2004	1302	43	10.39	0.00180
	2005	1280	42	14.58	0.00287
	2006	996	32	12.45	0.00245
	2007	1282	42	14.84	0.00252
	2008	2406	80	34.19	0.00507
	2009	2274	75	4.70	0.00225
2010	2797	93	37.20	0.00366	
2011	1320	44	23.04	0.00343	
Region	Year	Quahogs measured	Trips sampled	% of trips sampled	Samples per bushel landed
LI	1996	30	1	0.12	0.00002
	1997	1012	32	5.02	0.00089
	1998	480	16	2.28	0.00032
	1999	1440	48	7.12	0.00103
	2000	390	13	2.63	0.00037
	2001	180	6	1.05	0.00014
	2002	150	5	0.63	0.00007
	2003	990	33	3.26	0.00039
	2004	360	12	1.37	0.00015
	2005	1866	62	9.00	0.00087
	2006	2928	98	12.68	0.00120
	2007	2099	68	8.58	0.00084
	2008	2482	81	11.81	0.00101
	2009	2589	85	16.67	0.00136
2010	3564	119	20.73	0.00164	
2011	5386	180	31.25	0.00239	

Table 8 Cont'd. Number of quahogs measured, number of trips sampled, percentage of trips sampled, and number of quahogs measured per bushel landed by year and region, from port samples.

Region	Year	Quahogs measured	Trips sampled	% of trips sampled	Samples per bushel landed
NJ	1996	30	1	0.14	0.00003
	1997	390	13	2.03	0.00042
	1998	420	14	3.47	0.00071
	1999	420	14	3.13	0.00063
	2000	600	20	4.13	0.00082
	2001	780	26	3.99	0.00078
	2002	510	17	4.59	0.00083
	2003	390	13	2.68	0.00048
	2004	1080	36	9.92	0.00177
	2005	90	3	3.23	0.00061
	2006	243	8	11.59	0.00236
	2007	343	11	6.04	0.00099
	2008	330	11	4.74	0.00086
	2009	630	21	8.43	0.00118
	2010	780	26	8.84	0.00153
2011	690	23	10.95	0.00168	
DMV	1996	180	6	5.08	0.00111
	1997	570	19	10.86	0.00241
	1998	390	13	6.70	0.00130
	1999	960	32	19.39	0.00399
	2000	690	23	14.65	0.00299
	2001	660	22	18.64	0.00335
	2002	120	4	1.78	0.00031
	2003	390	13	10.66	0.00197
	2004	150	5	4.46	0.00109
	2005	511	17	12.32	0.00255
	2006	743	24	29.63	0.00683
	2007	195	6	42.86	0.00887
	2008	120	4	10.00	0.00202
	2009	95	3	10.71	0.00205
	2010	270	9	13.85	0.00286
2011	180	6	16.22	0.00282	

Table 9. Random and nearly random NEFSC survey tows used to estimate trends in ocean quahog abundance. Numbers in each cell are the number of tows used for each combination of stratum and cruise. Figures in plain text are the number of original tows (without borrowing). Outlined cells had zero tows originally but were filled by borrowing tows from the same strata during previous and/or subsequent cruises. Black cells have zero tows that could not be filled by borrowing. Survey/region combinations with relatively poor sampling (large amount of borrowed or missing tows, or borrowed or missing tows from large strata) are shown in grey.

region	stratum	area (nm2)	% total region area	SURVEY YEAR													
				1982	1983	1984	1986	1989	1992	1994	1997	1999	2002	2005	2008	2011	
GBK	54	295	0.04		3	3	3	6	3	3	3	3		2	2	2	
	55	386	0.06	3	3	3	3	1	3	3	3	2	2	4	2	3	
	56	214	0.03									4	4	4			
	57	176	0.02			2	2	1	2	5	2	2	2	4	2	11	
	58	303	0.04									5	5	5			
	59	512	0.07	1	4	5	1	2	6	5	5	4	5	9	4	16	
	60	801	0.11			2	2	2	4	2	5	5	5	9	4	5	
	61	588	0.09	8	1	6	5	12	7	6	6	6	6	11	5	5	
	62	731	0.10			1	1	1	4	4	4	4	4	7	3	4	
	65	184	0.03			2	2	4	2	4	3	4	1	1	3	3	
	67	196	0.03			5	5	5	7	7	7	7		2	2	1	
	68	380	0.06	1	8	7	3	6	6	5	5	5		6	6	6	
	69	902	0.13	2	5	11	6	6	6	7	6	8	8	4	4	1	
	70	544	0.08	1	2	6	4	8	4	4	4	3	2	6	4	19	
	71	168	0.02			2	2	3	1	2	3	3	1	2	3	1	3
	72	472	0.07	2	10	8	1	8	8	8	8	6	6	4	4	5	
73	526	0.08	1	1	4	3	6	6	6	6	5	6	9	3	5		
74	443	0.06	3	4	1	3	7	4	4	4	3	3	6	3	11		

Table 9 Cont'd.

region	stratum	area (nm2)	% total region area	SURVEY YEAR												
				1982	1983	1984	1986	1989	1992	1994	1997	1999	2002	2005	2008	2011
SNE	37	660	0.14	7	4	7	3	6	3	5	4	4	3	3	2	2
	38	268	0.06	3	2	5	3	3	3	5	3	3	3	2	3	7
	39	946	0.20	6	4	6	2	5	5	5	5	5	5	5	5	8
	41	580	0.12	6	5	7	5	6	6	6	6	5	6	6	6	4
	45	407	0.09	3	7	9	4	4	4	4	4	4	3	3	4	7
	46	205	0.04	2	5	5	3	2	3	5	3	3	2	3	3	6
	47	873	0.18	4	3	4	2	2	4	5	4	3	1	7	4	8
	94	215	0.04	1	2	2		1	1	2	2	4	2	2	5	5
	95	278	0.06	4	14	11	4	4	4	4	4	4	4	8	4	5
	96	490	0.10	12	12	13	1	1	3	2	4	4		1	1	1
LI	29	1078	0.24	11	10	20	10	10	10	10	10	11	10	10	16	10
	30	667	0.15	7	8	14	6	6	6	6	6	7	6	7	12	4
	31	932	0.21	9	7	12	5	7	8	8	8	9	8	10	8	8
	33	361	0.08	4	4	8	4	4	4	5	4	4	4	4	10	4
	34	207	0.05	2	2	4	2	2	2	5	2	2	2	2	8	6
	35	614	0.14	4	2	4	2	5	6	6	6	6	6	6	6	5
	91	342	0.08	2	2	4	4	3	3	3	3	3	3	3	5	11
	92	165	0.04	2	2	3	2	2	2	2	2	2	2	2	5	11
	93	97	0.02	1	1	2	1	1	1	1	1	1	2	1	4	6

Table 9 Cont'd.

				SURVEY YEAR													
region	stratum	area (nm2)	% total region area	1982	1983	1984	1986	1989	1992	1994	1997	1999	2002	2005	2008	2011	
NJ	17	703	0.11	11	11	17	12	12	12	12	14	12	12	12	12	5	
	18	240	0.04	3	3	6	3	3	3	3	3	3	3	3	3	5	
	19	266	0.04	3	3	6	3	2	3	3	3	3	3	3	3	4	
	21	1693	0.26	18	18	21	19	20	20	23	26	39	29	20	28	15	
	22	305	0.05	3	3	6	3	3	3	5	3	3	3	3	3	5	
	23	724	0.11	7	6	11	5	4	5	5	5	5	5	5	5	5	
	25	647	0.10	9	9	13	8	9	9	9	12	8	9	9	13	8	
	26	190	0.03	2	2	5	3	3	3	3	3	3	3	3	3	3	
	27	442	0.07	4	4	8	4	4	4	4	4	4	4	4	3	4	4
	87	356	0.05	8	7	10	9	9	9	9	9	9	16	8	9	6	
	88	484	0.07	15	15	24	17	20	20	20	21	23	20	17	19	6	
	89	343	0.05	14	15	21	15	18	17	17	19	18	18	15	18	4	
	90	117	0.02	2	2	3	2	2	2	2	2	2	2	2	2	1	4
DMV	9	1894	0.47	30	26	35	29	37	37	39	39	38	39	36	31	15	
	10	190	0.05	2	2	3	3	3	3	3	3	3	3	3	2	4	
	11	246	0.06	2	2	4	2	2	2	2	2	2	2	2	6	4	
	13	1149	0.28	19	18	25	20	20	20	21	22	19	20	18	15	7	
	14	205	0.05	2	2	3	3	3	3	5	3	3	3	3	26	23	
	15	387	0.10	4	4	8	4	4	4	5	4	5	4	4	9	5	
SVA	5	690	0.97	4	9	13	8	8	8	7	8	16	8	8	17	9	
	6	22	0.03	1	1	1	1	1	1	1	1	3	2	1	1		

Table 10. Trends in abundance and biomass for ocean quahogs ≥ 50 mm SL during 1982-2011, based on NEFSC clam survey data. Survey values are the quahogs caught in the survey dredge. Stock values are the survey values adjusted to account for the selectivity and efficiency of the survey dredge. Fishable values are the stock values adjusted to account for the selectivity of a commercial dredge. Figures include original plus borrowed tows. "N strata surveyed" includes strata sampled by tows borrowed from the previous and subsequent surveys if needed. Survey data for 1994 should be ignored because of gear problems that artificially boosted sampling efficiency. Survey coverage was incomplete on GBK prior to 1986, and in 2005.

region	year	SURVEY				STOCK				FISHABLE				n tows	n positive tows	n strata surveyed
		n/tow	CV	kg/tow	CV	n/tow	CV	kg/tow	CV	n/tow	CV	kg/tow	CV			
GBK	1986	278	0.19	6	0.18	430	0.23	9	0.20	256	0.18	6	0.18	47	21	16
	1989	92	0.26	2	0.25	127	0.24	3	0.25	87	0.26	2	0.25	78	38	16
	1992	346	0.21	9	0.21	486	0.19	11	0.20	328	0.21	9	0.21	74	41	16
	1994	405	0.20	11	0.20	578	0.19	14	0.19	384	0.20	10	0.20	76	40	16
	1997	270	0.19	7	0.19	389	0.19	9	0.18	253	0.19	7	0.19	83	44	18
	1999	273	0.17	8	0.19	366	0.16	9	0.18	260	0.17	8	0.19	77	47	18
	2002	328	0.18	9	0.19	478	0.15	11	0.18	310	0.18	9	0.19	61	38	15
	2008	324	0.30	6	0.28	693	0.31	11	0.29	285	0.29	6	0.27	49	30	15
	2011	317	0.23	7	0.22	606	0.21	10	0.22	284	0.23	6	0.22	94	69	15
SNE	1982	278	0.27	7	0.25	346	0.28	8	0.26	265	0.27	7	0.25	48	30	10
	1983	173	0.29	4	0.30	238	0.31	5	0.29	163	0.29	4	0.30	58	37	10
	1984	188	0.27	5	0.29	234	0.26	6	0.28	180	0.27	5	0.29	69	38	10
	1986	289	0.31	7	0.31	394	0.35	9	0.32	273	0.30	7	0.30	27	23	9
	1989	275	0.19	7	0.18	353	0.21	8	0.19	261	0.19	6	0.18	34	29	10
	1992	333	0.19	9	0.19	400	0.19	10	0.19	320	0.19	8	0.20	36	31	10
	1994	529	0.22	13	0.20	670	0.25	16	0.21	504	0.22	13	0.20	43	32	10
	1997	293	0.54	6	0.45	448	0.61	8	0.52	269	0.53	6	0.43	39	27	10
	1999	252	0.54	6	0.48	313	0.56	7	0.51	241	0.53	6	0.48	39	30	10
	2002	181	0.22	5	0.22	207	0.22	6	0.22	176	0.22	5	0.22	29	28	9
	2005	158	0.26	4	0.23	334	0.42	5	0.27	145	0.25	3	0.22	40	34	10
	2008	201	0.25	4	0.22	524	0.42	7	0.27	180	0.24	4	0.22	37	31	8
	2011	157	0.28	4	0.24	255	0.32	5	0.27	145	0.28	3	0.23	52	42	9

Table 10 Cont'd.

region	year	SURVEY				STOCK				FISHABLE				n tows	n positive tows	n strata surveyed
		n/tow	CV	kg/tow	CV	n/tow	CV	kg/tow	CV	n/tow	CV	kg/tow	CV			
LI	1982	277.91	0.15	6.98	0.16	433.99	0.16	9.29	0.15	256.75	0.15	6.64	0.16	42	36	9
	1983	185.88	0.21	5.23	0.21	253.51	0.22	6.36	0.21	175.68	0.21	5.04	0.21	38	36	9
	1984	239.24	0.17	6.67	0.16	323.92	0.17	8.11	0.16	225.92	0.17	6.42	0.16	71	63	9
	1986	319.60	0.22	8.89	0.20	426.26	0.22	10.78	0.21	302.09	0.21	8.56	0.20	36	31	9
	1989	226.21	0.34	5.06	0.29	367.49	0.38	7.15	0.33	206.65	0.33	4.74	0.28	40	36	9
	1992	323.33	0.18	8.31	0.16	465.23	0.20	10.62	0.17	301.75	0.18	7.93	0.16	42	36	9
	1994	592.57	0.16	15.35	0.16	827.85	0.17	19.30	0.16	555.85	0.16	14.67	0.16	46	44	9
	1997	401.64	0.16	11.16	0.16	518.85	0.17	13.35	0.16	381.48	0.16	10.75	0.16	42	35	9
	1999	232.27	0.17	6.28	0.15	310.52	0.19	7.67	0.16	219.40	0.17	6.03	0.14	45	41	9
	2002	253.06	0.21	6.97	0.20	330.41	0.21	8.39	0.20	240.10	0.21	6.71	0.20	43	40	9
	2005	149.38	0.19	4.07	0.19	215.78	0.19	5.06	0.18	140.78	0.19	3.92	0.19	45	39	9
	2008	155.33	0.16	4.55	0.15	206.67	0.19	5.41	0.16	147.68	0.16	4.40	0.15	74	66	9
	2011	195.56	0.19	5.29	0.20	289.72	0.20	6.61	0.20	184.31	0.19	5.08	0.20	65	54	9
NJ	1982	112.34	0.20	3.56	0.20	129.33	0.20	3.92	0.20	109.15	0.20	3.48	0.20	99	50	13
	1983	86.09	0.21	2.83	0.21	98.42	0.21	3.09	0.21	83.89	0.21	2.78	0.21	98	55	13
	1984	147.61	0.24	4.67	0.24	170.30	0.24	5.15	0.24	143.44	0.24	4.58	0.24	151	79	13
	1986	144.02	0.23	4.91	0.22	159.78	0.24	5.28	0.22	141.03	0.23	4.84	0.22	103	52	13
	1989	72.24	0.22	2.16	0.21	88.60	0.22	2.45	0.21	69.63	0.22	2.11	0.22	109	52	13
	1992	88.04	0.18	3.02	0.17	97.82	0.18	3.25	0.17	86.26	0.18	2.98	0.17	110	52	13
	1994	235.41	0.22	7.61	0.20	269.04	0.22	8.33	0.21	229.38	0.22	7.48	0.20	115	59	13
	1997	122.26	0.15	4.27	0.15	135.78	0.16	4.58	0.15	119.85	0.15	4.21	0.15	124	59	13
	1999	59.48	0.15	2.02	0.14	72.27	0.15	2.22	0.14	57.77	0.15	1.99	0.14	132	61	13
	2002	89.79	0.23	3.23	0.24	101.12	0.22	3.46	0.23	87.98	0.24	3.19	0.24	127	60	13
	2005	47.08	0.16	1.57	0.15	62.36	0.15	1.77	0.15	45.37	0.16	1.54	0.15	103	54	13
	2008	45.15	0.17	1.50	0.16	60.59	0.17	1.70	0.16	43.41	0.17	1.47	0.16	121	65	13
	2011	74.76	0.15	2.27	0.14	99.69	0.16	2.63	0.14	71.55	0.15	2.21	0.14	74	53	13

Table 10 Cont'd.

		SURVEY				STOCK				FISHABLE				n tows	n positive tows	n strata surveyed
region	year	n/tow	CV	kg/tow	CV	n/tow	CV	kg/tow	CV	n/tow	CV	kg/tow	CV			
DMV	1982	79.16	0.32	2.96	0.34	86.64	0.31	3.16	0.33	77.73	0.32	2.92	0.34	59	24	6
	1983	86.23	0.49	2.55	0.42	106.61	0.52	2.99	0.45	82.37	0.49	2.46	0.42	54	28	6
	1984	52.01	0.35	1.67	0.30	63.19	0.36	1.90	0.31	49.99	0.34	1.62	0.30	78	34	6
	1986	75.68	0.23	2.53	0.22	86.74	0.24	2.80	0.22	73.54	0.23	2.48	0.22	61	28	6
	1989	64.35	0.58	1.80	0.46	82.47	0.62	2.18	0.51	60.90	0.56	1.73	0.45	69	31	6
	1992	71.98	0.36	2.29	0.31	85.41	0.40	2.59	0.33	69.42	0.36	2.22	0.30	69	25	6
	1994	39.46	0.25	1.33	0.23	47.97	0.27	1.49	0.24	38.21	0.25	1.30	0.23	75	28	6
	1997	47.74	0.21	1.67	0.21	56.44	0.22	1.85	0.21	46.34	0.21	1.64	0.21	73	28	6
	1999	28.36	0.29	0.95	0.27	33.39	0.29	1.06	0.27	27.51	0.29	0.93	0.26	70	23	6
	2002	31.81	0.25	1.11	0.23	38.77	0.26	1.23	0.23	30.84	0.25	1.08	0.22	71	19	6
	2005	19.41	0.49	0.69	0.53	24.84	0.45	0.78	0.50	18.82	0.50	0.68	0.53	66	21	6
	2008	17.76	0.54	0.62	0.59	22.61	0.49	0.70	0.56	17.22	0.54	0.61	0.59	57	16	6
	2011	16.14	0.31	0.51	0.35	22.55	0.28	0.61	0.32	15.28	0.31	0.50	0.36	58	38	6
SVA	1982	0.04	0.00	0.00	0.00	0.04	0.00	0.00	0.00	0.04	0.00	0.00	0.00	5	1	2
	1983	1.89	0.58	0.10	0.58	1.92	0.58	0.10	0.58	1.89	0.58	0.10	0.58	10	3	2
	1984	0.19	0.85	0.01	0.87	0.19	0.84	0.01	0.87	0.19	0.85	0.01	0.87	14	2	2
	1986	0.29	0.00	0.01	0.00	0.29	0.00	0.01	0.00	0.28	0.00	0.01	0.00	9	1	2
	1989	0.39	0.00	0.02	0.00	0.40	0.00	0.02	0.00	0.39	0.00	0.02	0.00	9	1	2
	1992	0.00		0.00		0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	9	0	2
	1994	4.47	0.79	0.23	0.81	4.56	0.78	0.23	0.80	4.45	0.79	0.22	0.81	8	2	2
	1997	0.15	0.00	0.00	0.00	0.28	0.00	0.01	0.00	0.14	0.00	0.00	0.00	9	1	2
	1999	0.08	0.55	0.00	0.61	0.18	0.50	0.00	0.54	0.07	0.56	0.00	0.61	19	2	2
	2002	0.04	1.00	0.00	1.00	0.13	1.00	0.00	1.00	0.04	1.00	0.00	1.00	10	1	2
	2005	0.00		0.00		0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	9	0	2
	2008	0.00		0.00		0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	9	0	2
	2011	0.00		0.00		0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	9	0	1

Table 11. Efficiency-corrected ocean quahog swept-area biomass estimates by region (in thousands of metric tons fishable stock) during years with NEFSC clam surveys. Data for SVA and GBK for 2005 are basically averages of 2002 and 2008 as SVA and GBK were not sampled during 2005.

Area of region (nm ²) - no unsuitable habitat correction			INPUT: Fraction suitable habitat (u)		
SVA	712	10%	SVA	1	10%
DMV	4071	10%	DMV	1	10%
NJ	6510	10%	NJ	1	10%
LI	4463	10%	LI	1	10%
SNE	4922	10%	SNE	0.958	10%
GBK	7821	10%	GBK	0.9	10%
Total	28499				

Habitat area in assessment region (nm ²)			INPUT: Biomass fraction in unsurveyed deep water		
SVA	712	14%	SVA	0	10%
DMV	4071	14%	DMV	0	10%
NJ	6510	14%	NJ	0	10%
LI	4463	14%	LI	0	10%
SNE	4714.03	14%	SNE	0.02	10%
GBK	7038.9	14%	GBK	0.13	10%

INPUT: Original survey mean catch of fishable stock (kg/tow, tows adjusted to nominal distance using sensors)												
	1997	CV	1999	CV	2002	CV	2005	CV	2008	CV	2011	CV
SVA	0.00	1.00	0.00	0.55	0.00	1.00	0.00	1.00	0.00	1.00	0.00	1.00
DMV	0.69	0.22	0.47	0.27	0.72	0.24	0.44	0.48	0.37	0.33	0.47	0.48
NJ	1.81	0.15	1.02	0.14	1.93	0.23	1.10	0.14	1.25	0.20	1.96	0.22
LI	4.88	0.17	3.22	0.14	3.63	0.17	2.33	0.16	3.61	0.15	3.74	0.23
SNE	2.39	0.38	2.91	0.46	3.47	0.26	2.19	0.22	3.36	0.27	2.39	0.34
GBK	2.87	0.16	3.36	0.18	4.15	0.18	4.64	0.20	4.98	0.27	4.12	0.24

Swept-area biomass without efficiency correction (B', 1000 mt):												
	1997	CV	1999	CV	2002	CV	2005	CV	2008	CV	2011	CV
SVA	0.01	1.02	0.00	0.59	0.00	1.02	0.00	1.02	0.00	1.02	0.00	1.02
DMV	22.62	0.30	15.53	0.33	23.75	0.31	14.58	0.52	12.33	0.39	15.45	0.52
NJ	95.69	0.25	53.62	0.24	101.92	0.30	58.10	0.24	66.12	0.28	103.57	0.29
LI	176.41	0.26	116.51	0.25	131.46	0.27	84.13	0.26	130.45	0.25	135.24	0.31
SNE	93.27	0.43	113.26	0.50	135.17	0.33	85.52	0.30	131.10	0.34	93.26	0.39
GBK	185.23	0.26	216.72	0.27	267.59	0.27	298.84	0.28	321.27	0.33	265.69	0.31
Total w/o GBK	388.00	0.17	298.93	0.22	392.30	0.16	242.34	0.15	340.00	0.17	347.52	0.18
Total	573.22	0.14	515.64	0.17	659.89	0.15	541.18	0.17	661.28	0.18	613.22	0.17

Table 11 Cont'd. Efficiency-corrected ocean quahog swept-area biomass estimates by region (in thousands of metric tons fishable stock) during years with NEFSC clam surveys. Data for SVA and GBK for 2005 are basically averages of 2002 and 2008 as SVA and GBK were not sampled during 2005.

INPUT: Survey dredge efficiency (e)

	1997	CV	1999	CV	2002	CV	2005	CV	2008	CV	2011	CV
	0.169	0.21	0.169	0.21	0.169	0.21	0.169	0.21	0.169	0.21	0.169	0.21

Efficiency adjusted swept area fishable biomass (B, 1000 mt)

	1997	CV	1999	CV	2002	CV	2005	CV	2008	CV	2011	CV
SVA	0.05	1.04	0.02	0.62	0.01	1.04	0.03	1.04	0.03	1.04	0.03	1.04
DMV	133.82	0.37	91.88	0.39	140.56	0.37	86.27	0.56	72.96	0.44	91.40	0.56
NJ	566.19	0.33	317.29	0.32	603.05	0.37	343.81	0.32	391.23	0.35	612.83	0.36
LI	1043.87	0.34	689.40	0.32	777.86	0.34	497.83	0.33	771.91	0.33	800.24	0.37
SNE	551.90	0.47	670.20	0.54	799.83	0.39	506.02	0.37	775.73	0.40	551.85	0.45
GBK	1096.01	0.33	1282.35	0.34	1583.35	0.34	1768.29	0.35	1901.03	0.39	1572.15	0.38
Total w/o GBK	2295.83	0.27	1768.80	0.30	2321.31	0.27	1433.96	0.26	2011.85	0.27	2056.35	0.28
Total	3391.84	0.25	3051.15	0.27	3904.67	0.26	3202.24	0.27	3912.88	0.28	3628.51	0.27

Lower bound for 80% confidence intervals on fishable biomass (1000 mt, for lognormal distribution with no bias correction)

	1997	1999	2002	2005	2008	2011
SVA	0.02	0.01	0.00	0.01	0.01	0.01
DMV	84.99	56.48	88.39	44.14	42.63	46.82
NJ	377.04	212.24	382.13	229.94	252.95	391.38
LI	686.10	460.78	510.22	329.02	512.82	505.03
SNE	309.98	348.96	495.13	321.41	475.13	319.60
GBK	723.40	839.89	1038.19	1144.45	1169.60	985.37
Total w/o GBK	1634.53	1210.57	1659.38	1033.40	1430.58	1448.72
Total	2463.94	2173.60	2829.82	2281.19	2754.97	2581.08

Upper bound for 80% confidence intervals on fishable biomass (1000 mt, for lognormal distribution with no bias correction)

	1997	1999	2002	2005	2008	2011
SVA	0.14	0.05	0.04	0.09	0.09	0.09
DMV	210.70	149.46	223.53	168.61	124.86	178.42
NJ	850.24	474.36	951.68	514.08	605.10	959.58
LI	1588.20	1031.43	1185.92	753.25	1161.89	1268.02
SNE	982.64	1287.18	1292.03	796.66	1266.52	952.90
GBK	1660.56	1957.91	2414.79	2732.17	3089.86	2508.36
Total w/o GBK	3224.69	2584.43	3247.29	1989.78	2829.31	2918.84
Total	4669.19	4282.99	5387.78	4495.19	5557.45	5101.00

Table 12. Fishing mortality estimates based on catch and efficiency-corrected swept-area biomass for fishable ocean quahogs during years with clam surveys. CV's are based on analytical variance calculations assuming log normality, and include uncertainty in catch, survey data, swept-area, amount of suitable habitat, and survey dredge efficiency.

INPUT: Upper bound incidental mortality allowance 0.05

INPUT: Assumed CV for catch 0.1

INPUT: Landings (1000 mt, discard ~ 0)

	1997	1999	2002	2005	2008	2011
SVA	0	0	0	0	0	0
DMV	1.07	1.09	1.74	0.91	0.27	0.29
NJ	4.23	3.04	2.79	0.67	1.73	1.87
LI	5.14	6.34	9.14	9.73	11.12	10.24
SNE	8.97	6.63	3.90	2.02	2.15	1.75
GBK	0	0	0	0	0	0
Total	19.41	17.10	17.56	13.33	15.28	14.14

INPUT: Catch

	1997	1999	2002	2005	2008	2011
SVA	0	0	0	0	0	0
DMV	1.13	1.15	1.82	0.96	0.28	0.30
NJ	4.44	3.20	2.93	0.70	1.82	1.96
LI	5.40	6.66	9.60	10.21	11.68	10.75
SNE	9.42	6.96	4.09	2.12	2.26	1.84
GBK	0	0	0	0	0	0
Total	20.38	17.96	18.44	14.00	16.04	14.85

INPUT: Efficiency Corrected Swept Area Biomass for Fishable Stock (1000 mt)

	1997	CV	1999	CV	2002	CV	2005	CV	2008	CV	2011	CV
SVA	0.05	1.04	0.02	0.62	0.01	1.04	0.03	1.04	0.03	1.04	0.03	1.04
DMV	133.82	0.37	91.88	0.39	140.56	0.37	86.27	0.56	72.96	0.44	91.40	0.56
NJ	566.19	0.33	317.29	0.32	603.05	0.37	343.81	0.32	391.23	0.35	612.83	0.36
LI	1043.87	0.34	689.40	0.32	777.86	0.34	497.83	0.33	771.91	0.33	800.24	0.37
SNE	551.90	0.47	670.20	0.54	799.83	0.39	506.02	0.37	775.73	0.40	551.85	0.45
GBK	1096.01	0.33	1282.35	0.34	1583.35	0.34	1768.29	0.35	1901.03	0.39	1572.15	0.38
Total w/o GBK	2295.83	0.27	1768.80	0.30	2321.31	0.27	1433.96	0.26	2011.85	0.27	2056.35	0.28
Total	3391.84	0.25	3051.15	0.27	3904.67	0.26	3202.24	0.27	3912.88	0.28	3628.51	0.27

Table 12 Cont'd. Fishing mortality estimates based on catch and efficiency-corrected swept-area biomass for fishable ocean quahogs during years with clam surveys. CV's are based on analytical variance calculations assuming log normality, and include uncertainty in catch, survey data, swept-area, amount of suitable habitat, and survey dredge efficiency.

Fishing mortality (y-1)

	1997	CV	1999	CV	2002	CV	2005	CV	2008	CV	2011	CV
SVA	0.000	1.046	0.000	0.630	0.000	1.046	0.000	1.046	0.000	1.046	0.000	1.046
DMV	0.008	0.379	0.012	0.407	0.013	0.388	0.011	0.570	0.004	0.450	0.003	0.569
NJ	0.008	0.341	0.010	0.337	0.005	0.381	0.002	0.337	0.005	0.365	0.003	0.375
LI	0.005	NA	0.010	NA	0.012	0.353	0.021	0.347	0.015	0.343	0.013	0.385
SNE	0.017	0.485	0.010	0.554	0.005	0.401	0.004	0.379	0.003	0.410	0.003	0.458
GBK	0.000	NA	0.000	NA	0.000	NA	0.000	0.000	0.000	0.000	0.000	0.391
Total w/o GBK	0.009	0.288	0.010	0.319	0.008	0.285	0.010	0.279	0.008	0.289	0.007	0.296
Total	0.006	0.273	0.006	0.288	0.005	0.274	0.004	0.288	0.004	0.297	0.004	0.289

Lower bound for 80% confidence intervals for fishing mortality (y-1, for lognormal distribution with no bias correction)

	1997	1999	2002	2005	2008	2011
SVA	NA	NA	NA	NA	NA	NA
DMV	0.005	0.008	0.008	0.006	0.002	0.002
NJ	0.005	0.007	0.003	0.001	0.003	0.002
LI	NA	NA	0.008	0.013	0.010	0.008
SNE	0.009	0.005	0.003	0.003	0.002	0.002
GBK	NA	NA	NA	NA	NA	NA
Total w/o GBK	0.006	0.007	0.006	0.007	0.006	0.005
Total	0.004	0.004	0.003	0.003	0.003	0.003

Upper bound for 80% confidence intervals for fishing mortality (y-1, for lognormal distribution with no bias correction)

	1997	1999	2002	2005	2008	2011
SVA	NA	NA	NA	NA	NA	NA
DMV	0.013	0.021	0.021	0.022	0.007	0.007
NJ	0.012	0.015	0.008	0.003	0.007	0.005
LI	NA	NA	0.019	0.032	0.023	0.022
SNE	0.031	0.020	0.008	0.007	0.005	0.006
GBK	NA	NA	NA	NA	NA	NA
Total w/o GBK	0.013	0.015	0.011	0.014	0.011	0.010
Total	0.008	0.008	0.007	0.006	0.006	0.006

Table 13. KLAMZ estimates of ocean quahog biomass (in thousands of metric tons) from 1978 to 2011 by assessment region, the exploited regions only, and the whole stock. The CVs for the NJ and SNE areas were bootstrapped as the models did not converge.

	DMV		NJ		LI		SNE		GBK		Exploited area		Whole stock	
	B	CV	B	CV	B	CV	B	CV	B	CV	B	CV	B	CV
1978	293	0.11	943	0.14	718	0.28	1,055	0.07	1,161	0.54	2,024	0.27	3,460	0.22
1979	286	0.11	917	0.13	730	0.26	1,035	0.08	1,185	0.50	2,038	0.26	3,500	0.22
1980	273	0.11	892	0.13	741	0.25	1,016	0.10	1,209	0.47	2,046	0.25	3,540	0.21
1981	263	0.12	868	0.13	753	0.24	997	0.11	1,233	0.43	2,057	0.24	3,580	0.21
1982	254	0.12	843	0.12	764	0.23	978	0.12	1,257	0.40	2,068	0.24	3,620	0.20
1983	244	0.12	818	0.12	775	0.22	960	0.12	1,281	0.37	2,077	0.23	3,660	0.20
1984	233	0.12	795	0.12	786	0.21	941	0.13	1,305	0.34	2,087	0.22	3,700	0.19
1985	221	0.13	772	0.11	797	0.20	922	0.13	1,328	0.32	2,093	0.22	3,730	0.19
1986	209	0.13	746	0.11	808	0.19	904	0.14	1,352	0.29	2,096	0.21	3,760	0.19
1987	196	0.14	723	0.11	819	0.19	886	0.14	1,375	0.27	2,101	0.20	3,800	0.18
1988	182	0.15	701	0.10	828	0.18	869	0.15	1,398	0.25	2,103	0.20	3,830	0.18
1989	167	0.16	682	0.10	838	0.18	851	0.15	1,421	0.23	2,106	0.19	3,860	0.18
1990	157	0.16	656	0.10	848	0.17	834	0.15	1,443	0.21	2,107	0.19	3,890	0.18
1991	150	0.17	628	0.10	858	0.17	817	0.15	1,465	0.20	2,110	0.19	3,920	0.17
1992	143	0.17	603	0.10	867	0.16	801	0.16	1,487	0.19	2,112	0.18	3,950	0.17
1993	137	0.18	585	0.10	848	0.16	785	0.16	1,481	0.19	2,073	0.18	3,890	0.17
1994	133	0.18	564	0.10	833	0.16	769	0.16	1,474	0.18	2,034	0.18	3,840	0.17
1995	129	0.18	546	0.10	815	0.17	753	0.16	1,466	0.18	1,996	0.18	3,790	0.17
1996	126	0.18	530	0.10	798	0.17	733	0.16	1,456	0.18	1,957	0.19	3,740	0.17
1997	123	0.18	516	0.10	785	0.17	711	0.16	1,445	0.18	1,919	0.19	3,690	0.17
1998	119	0.18	502	0.10	773	0.17	689	0.16	1,434	0.18	1,882	0.19	3,630	0.17
1999	115	0.19	490	0.10	759	0.17	669	0.17	1,421	0.18	1,846	0.19	3,580	0.17
2000	112	0.19	477	0.10	746	0.17	649	0.17	1,408	0.18	1,810	0.19	3,530	0.18
2001	109	0.19	465	0.10	734	0.17	632	0.17	1,394	0.18	1,777	0.19	3,480	0.18
2002	106	0.19	451	0.11	720	0.17	615	0.17	1,380	0.18	1,742	0.19	3,430	0.18
2003	102	0.19	440	0.11	704	0.17	599	0.17	1,365	0.18	1,705	0.19	3,380	0.18
2004	99	0.20	428	0.11	685	0.18	586	0.17	1,349	0.18	1,668	0.19	3,320	0.18
2005	96	0.20	417	0.11	667	0.18	571	0.17	1,333	0.18	1,632	0.19	3,270	0.18
2006	94	0.20	409	0.11	650	0.18	558	0.17	1,317	0.18	1,600	0.20	3,220	0.18
2007	91	0.20	400	0.11	631	0.18	546	0.17	1,300	0.18	1,567	0.20	3,170	0.18
2008	89	0.20	391	0.12	613	0.19	533	0.17	1,284	0.18	1,534	0.20	3,120	0.18
2009	87	0.20	382	0.12	594	0.19	521	0.17	1,266	0.18	1,500	0.20	3,070	0.18
2010	85	0.20	372	0.12	579	0.19	506	0.17	1,249	0.18	1,466	0.20	3,010	0.18
2011	83	0.20	363	0.13	562	0.20	493	0.17	1,232	0.18	1,431	0.20	2,960	0.18

Table 14. KLAMZ estimates of ocean quahog annual fishing mortality from 1978 to 2011 by assessment region, the exploited regions only, and the whole stock. The CVs for the NJ and SNE areas were bootstrapped as the models did not converge.

	DMV		NJ		LI		SNE		GBK		Exploited area		Whole stock	
	F	CV	F	CV	F	CV	F	CV	F	CV	F	CV	F	CV
1978	0.006	0.11	0.009	0.21					0	0	0.005	0.27	0.003	0.23
1979	0.027	0.11	0.009	0.20					0	0	0.008	0.26	0.005	0.22
1980	0.018	0.12	0.010	0.19	0.000	0.25			0	0	0.007	0.25	0.004	0.21
1981	0.015	0.12	0.011	0.18	0.000	0.24			0	0	0.006	0.25	0.004	0.21
1982	0.020	0.12	0.011	0.17					0	0	0.007	0.24	0.004	0.20
1983	0.023	0.12	0.010	0.16	0.000	0.22	0.001	0.07	0	0	0.007	0.23	0.004	0.20
1984	0.034	0.13	0.012	0.16			0.001	0.07	0	0	0.009	0.22	0.005	0.19
1985	0.037	0.13	0.016	0.15	0.000	0.20	0.001	0.07	0	0	0.010	0.22	0.006	0.19
1986	0.042	0.14	0.013	0.15	0.000	0.19	0.001	0.07	0	0	0.009	0.21	0.005	0.19
1987	0.056	0.14	0.013	0.14	0.001	0.19	0.001	0.07	0	0	0.010	0.21	0.006	0.18
1988	0.067	0.15	0.010	0.13	0.001	0.18	0.001	0.07	0	0	0.010	0.20	0.005	0.18
1989	0.040	0.16	0.021	0.13	0.001	0.18	0.001	0.07	0	0	0.011	0.20	0.006	0.18
1990	0.024	0.17	0.024	0.12	0.001	0.17	0.001	0.07	0	0	0.010	0.19	0.005	0.18
1991	0.033	0.17	0.024	0.12	0.002	0.17	0.001	0.07	0	0	0.011	0.19	0.006	0.17
1992	0.017	0.17	0.012	0.12	0.014	0.16	0.001	0.07	0	0	0.011	0.18	0.006	0.17
1993	0.014	0.18	0.018	0.12	0.010	0.16	0.001	0.07	0	0	0.011	0.18	0.006	0.17
1994	0.008	0.18	0.013	0.12	0.015	0.17	0.001	0.07	0	0	0.010	0.19	0.005	0.17
1995	0.005	0.18	0.010	0.12	0.012	0.17	0.007	0.07	0	0	0.011	0.19	0.006	0.17
1996	0.006	0.18	0.009	0.11	0.008	0.17	0.012	0.07	0	0	0.010	0.19	0.005	0.17
1997	0.009	0.18	0.008	0.11	0.007	0.17	0.013	0.07	0	0	0.010	0.19	0.005	0.17
1998	0.012	0.18	0.005	0.11	0.009	0.17	0.010	0.07	0	0	0.009	0.19	0.005	0.17
1999	0.010	0.19	0.006	0.11	0.008	0.17	0.010	0.07	0	0	0.009	0.19	0.005	0.17
2000	0.010	0.19	0.007	0.11	0.006	0.17	0.008	0.08	0	0	0.008	0.19	0.004	0.18
2001	0.009	0.19	0.011	0.11	0.008	0.17	0.008	0.08	0	0	0.010	0.19	0.005	0.18
2002	0.017	0.19	0.006	0.11	0.013	0.17	0.006	0.08	0	0	0.010	0.19	0.005	0.18
2003	0.009	0.19	0.009	0.11	0.017	0.17	0.004	0.08	0	0	0.011	0.19	0.006	0.18
2004	0.006	0.20	0.007	0.12	0.016	0.18	0.006	0.08	0	0	0.011	0.19	0.005	0.18
2005	0.010	0.20	0.002	0.12	0.015	0.18	0.004	0.08	0	0	0.008	0.20	0.004	0.18
2006	0.005	0.20	0.001	0.12	0.017	0.18	0.003	0.08	0	0	0.009	0.20	0.004	0.18
2007	0.001	0.20	0.004	0.12	0.018	0.19	0.004	0.08	0	0	0.010	0.20	0.005	0.18
2008	0.003	0.20	0.004	0.12	0.018	0.19	0.004	0.08	0	0	0.010	0.20	0.005	0.18
2009	0.002	0.20	0.006	0.12	0.015	0.19	0.009	0.08	0	0	0.011	0.20	0.005	0.18
2010	0.005	0.20	0.006	0.13	0.017	0.19	0.007	0.08	0	0	0.011	0.20	0.005	0.18
2011	0.004	0.20	0.005	0.13	0.019	0.20	0.004	0.08	0	0	0.010	0.20	0.005	0.18

Table 15. Results of 15 year projections for the whole ocean quahog stock from the KLAMZ model under three different harvest scenarios, and the probability of the whole stock becoming overfished in each year ($B < 1.39$ million mt). Natural mortality is 0.02 in all scenarios.

Year	WHOLE STOCK								
	Status quo catch (15,400 mt)			Quota catch (24,190 mt)			Overfishing threshold (0.022 y^{-1})		
	mean B	p overfished	mean F	mean B	p overfished	mean F	mean B	p overfished	mean catch
2011	3192	0	0.005	3192	0	0.005	3192	0	14.3
2012	3145	0	0.005	3145	0	0.008	3145	0	68.0
2013	3098	0	0.005	3089	0	0.008	3046	0	65.9
2014	3051	0	0.005	3033	0	0.008	2949	0	63.8
2015	3004	0	0.005	2978	0	0.008	2855	0	61.7
2016	2957	0	0.005	2923	0	0.009	2764	0	59.8
2017	2910	0	0.005	2868	0	0.009	2676	0	57.9
2018	2864	0	0.006	2813	0	0.009	2591	0	56.0
2019	2818	0	0.006	2759	0	0.009	2508	0	54.2
2020	2772	0	0.006	2706	0	0.009	2428	0	52.5
2021	2727	0	0.006	2652	0	0.009	2350	0	50.8
2022	2682	0	0.006	2600	0	0.010	2275	0	49.2
2023	2637	0	0.006	2547	0	0.010	2203	0	47.6
2024	2593	0	0.006	2496	0	0.010	2132	0	46.1
2025	2549	0	0.006	2444	0	0.010	2065	0	44.6
2026	2506	0	0.006	2394	0	0.011	1999	0	43.2
Cumulative p		0			0			0	

Table 15 cont'd.

	EXPLOITED AREA								
	Status quo catch (15,400 mt)			Quota catch (24,190 mt)			Overfishing threshold (0.022 y ⁻¹)		
Year	mean B	mean F	p overfishing	mean B	mean F	p overfishing	mean B	mean catch	p overfishing
2011	1719	0.009	0	1719	0.009	0	1719	14.3	0
2012	1686	0.009	0	1686	0.015	0.004	1686	36.5	<i>F</i> set to <i>F</i> _{threshold} for projected years
2013	1653	0.010	0	1644	0.015	0.004	1632	35.3	
2014	1620	0.010	0	1603	0.016	0.008	1580	34.1	
2015	1587	0.010	0	1561	0.016	0.014	1529	33.0	
2016	1555	0.010	0	1521	0.017	0.025	1480	32.0	
2017	1523	0.011	0	1480	0.017	0.035	1432	31.0	
2018	1491	0.011	0	1440	0.018	0.068	1386	30.0	
2019	1459	0.011	0	1401	0.018	0.109	1342	29.0	
2020	1428	0.011	0	1362	0.019	0.175	1298	28.1	
2021	1397	0.012	0	1323	0.019	0.261	1257	27.2	
2022	1367	0.012	0	1285	0.020	0.363	1217	26.3	
2023	1337	0.012	0	1247	0.021	0.437	1178	25.4	
2024	1307	0.013	0.002	1210	0.022	0.511	1140	24.6	
2025	1277	0.013	0.002	1173	0.022	0.573	1104	23.8	
2026	1248	0.013	0.004	1137	0.023	0.655	1069	23.1	
Cumulative p			0.004			0.657			




Figure 1. Map of ocean quahog assessment regions, and individual strata used for ocean quahogs.




Figure 2. Commercial size-selectivity and maturity by length (top) and by age (bottom) assuming the von Bertalanffy growth curve for ocean quahogs in the exploited area (from Thorarinsdottir and Jacobson, 1995).




Figure 3. Total ocean quahog landings by region in millions of bushels.





Figure 4. Ocean quahog annual mean ex-vessel price per pound of meats for both the ITQ and Maine fisheries. The gray line shows the actual amount paid (the nominal price) while the black line shows the price adjusted for inflation from the beginning of the time series.




Figure 5. Hours fished by region and year, and sum of hours fished in all four regions.




Figure 6. Number of trips by region and year, and the sum of trips from all regions.




Figure 7. Mean trip length in hours by region and year, and for the four regions combined.




Figure 8. Number of vessels fishing for ocean quahogs by region and year, and the total number of vessels fishing in all four regions combined.




Figure 9. Mean ton class (a number specified by port agents to determine the relative size of a vessel) of the surfclam and ocean quahog fleet from 1994 to 2011. Mean ton class of fleet is the average of all unique vessels that landed greater than 7,500 lbs of meats per trip in a given year. Mean ton class by trip is the mean size of the vessels weighted by the number of trips they made that landed more than 7,500 lbs of meats. The y-axis scale for “ton class” is equivalent to about 140-160 gross tons.




Figure 10. Mean landings per trip in kilobushels by region and year, and for the four regions combined.




Figure 11. Ocean quahog LPUE (bushels per hour fished) by individual region, and the average of the four regions for each year.




Figure 12. Landings of ocean quahogs in millions of bushels by assessment region.




Figure 13. Map showing which statistical areas are used for calculating regional landings.




Figure 14. Assessment regions where vessels reported having fished in a given year. Top: Most vessels fish in two assessment regions. Middle: Vessels that fished in only one assessment region were most likely to fish in LI. Bottom: Vessels that fished in two regions have been moving away from LI and SNE and into NJ and LI in the past several years.




Figure 15. Ocean quahog landings by ten-minute square (TMSQ), the finest scale location for landings reported in logbooks, and time period. TMSQ in light blue had reported landings, but from fewer than three vessels.




Figure 16. Ocean quahog fishing effort by ten-minute square (TMSQ), the most precise location for fishing activity reported in logbooks, and time period. TMSQ in light blue had reported fishing activity, but from fewer than three vessels.




Figure 17. Ocean quahog landings by unit effort by ten-minute square (TMSQ), the most precise location for fishing activity reported in logbooks, and time period. TMSQ in light blue had reported fishing activity, but from fewer than three vessels.

Ocean quahog - DMV region - landings for important 10-minute squares




Figure 18A. Plots of landings from important TMSQ (landings of 100,000+ bushels in one year during the time series) in the DMV region, 1980-2011. Plots are sorted N to S then E to W. The caret marks at zero represent years where less than three vessels fished in the area, so the data point is not shown. The line is a smoother to show trend, fit to all data including those not plotted.

Ocean quahog - DMV region - fishing effort for important 10-minute squares




Figure 18A Cont'd. Plots of landings from important TMSQ (landings of 100,000+ bushels in one year during the time series) in the DMV region, 1980-2011. Plots are sorted N to S then E to W. The caret marks at zero represent years where less than three vessels fished in the area, so the data point is not shown. The line is a smoother to show trend, fit to all data including those not plotted.

Ocean quahog - DMV region - LPUE for important 10-minute squares




Figure 18A Cont'd. Plots of landings from important TMSQ (landings of 100,000+ bushels in one year during the time series) in the DMV region, 1980-2011. Plots are sorted N to S then E to W. The caret marks at zero represent years where less than three vessels fished in the area, so the data point is not shown. The line is a smoother to show trend, fit to all data including those not plotted.

Ocean quahog - NJ region - landings for important 10-minute squares




Figure 18B. Plots of landings from important TMSQ (landings of 100,000+ bushels in one year during the time series) in the NJ region, 1980-2011. Plots are sorted N to S then E to W. The caret marks at zero represent years where less than three vessels fished in the area, so the data point is not shown. The line is a smoother to show trend, fit to all data including those not plotted.

Ocean quahog - NJ region - fishing effort for important 10-minute squares




Figure 18B Cont'd. Plots of landings from important TMSQ (landings of 100,000+ bushels in one year during the time series) in the NJ region, 1980-2011. Plots are sorted N to S then E to W. The caret marks at zero represent years where less than three vessels fished in the area, so the data point is not shown. The line is a smoother to show trend, fit to all data including those not plotted.

Ocean quahog - NJ region - LPUE for important 10-minute squares




Figure 18B Cont'd. Plots of landings from important TMSQ (landings of 100,000+ bushels in one year during the time series) in the NJ region, 1980-2011. Plots are sorted N to S then E to W. The caret marks at zero represent years where less than three vessels fished in the area, so the data point is not shown. The line is a smoother to show trend, fit to all data including those not plotted.

Ocean quahog - LI region - landings for important 10-minute squares




Figure 18C. Plots of landings from important TMSQ (landings of 100,000+ bushels in one year during the time series) in the LI region, 1980-2011. Plots are sorted N to S then E to W. The caret marks at zero represent years where less than three vessels fished in the area, so the data point is not shown. The line is a smoother to show trend, fit to all data including those not plotted.

Ocean quahog - LI region - fishing effort for important 10-minute squares




Figure 18C Cont'd. Plots of landings from important TMSQ (landings of 100,000+ bushels in one year during the time series) in the LI region, 1980-2011. Plots are sorted N to S then E to W. The caret marks at zero represent years where less than three vessels fished in the area, so the data point is not shown. The line is a smoother to show trend, fit to all data including those not plotted.

Ocean quahog - LI region - LPUE for important 10-minute squares




Figure 18C Cont'd. Plots of landings from important TMSQ (landings of 100,000+ bushels in one year during the time series) in the LI region, 1980-2011. Plots are sorted N to S then E to W. The caret marks at zero represent years where less than three vessels fished in the area, so the data point is not shown. The line is a smoother to show trend, fit to all data including those not plotted.

Ocean quahog - SNE region - landings for important 10-minute squares




Figure 18D. Plots of landings from important TMSQ (landings of 100,000+ bushels in one year during the time series) in the SNE region, 1980-2011. Plots are sorted N to S then E to W. The caret marks at zero represent years where less than three vessels fished in the area, so the data point is not shown. The line is a smoother to show trend, fit to all data including those not plotted.

Ocean quahog - SNE region - fishing effort for important 10-minute squares




Figure 18D Cont'd. Plots of landings from important TMSQ (landings of 100,000+ bushels in one year during the time series) in the SNE region, 1980-2011. Plots are sorted N to S then E to W. The caret marks at zero represent years where less than three vessels fished in the area, so the data point is not shown. The line is a smoother to show trend, fit to all data including those not plotted.

Ocean quahog - SNE region - LPUE for important 10-minute squares




Figure 18D Cont'd. Plots of landings from important TMSQ (landings of 100,000+ bushels in one year during the time series) in the SNE region, 1980-2011. Plots are sorted N to S then E to W. The caret marks at zero represent years where less than three vessels fished in the area, so the data point is not shown. The line is a smoother to show trend, fit to all data including those not plotted.




Figure 19A. Length composition of port samples by year from the Delmarva region.




Figure 19B. Length composition of port samples by year from the New Jersey region.




Figure 19C. Length composition of port samples by year from the Long Island region.




Figure 19D. Length composition of port samples by year from the southern New England region.




Figure 20A. Port sample length frequencies from the SNE region plotted with survey length frequencies from the same region and year.




Figure 20B. Port sample length frequencies from the LI region plotted with survey length frequencies from the same region and year.




Figure 21. Fishery and survey selectivity curves for ocean quahog from NEFSC (2007a). The ratio of the fishery and survey selectivity curves, which can be used to convert survey abundance at size directly to fishable abundance at size, is also shown.




Figure 22. Map of an ocean quahog depletion experiment site from 2008. The start and stop locations are correct, but the tow paths were not straight lines. The map was made to look at the relationship of the setup tows (red lines) to the depletion tows (blue lines).




Figure 23. The number of ocean quahogs caught at each 2011 survey station. The small crosses are stations where no ocean quahogs were caught.




Figure 24. The number of ocean quahogs at least 70mm in shell length caught at each 2011 survey station. The small crosses are stations where no ocean quahogs larger than 69mm were caught.




Figure 25. The number of ocean quahogs less than 70mm in shell length caught at each 2011 survey station. The small crosses are stations where no ocean quahogs less than 70mm were caught.




Figure 26. Number per tow of ocean quahogs that would be available to the fishery (number per tow X fishery selectivity curve) from the NEFSC clam survey. The 1994 survey, when increased voltage to the dredge resulted in more than standard fishing power, is indicated by the open diamond.




Figure 27. Kilograms per tow of ocean quahogs that would be available to the fishery (number per tow X fishery selectivity curve) from the NEFSC clam survey. The 1994 survey, when increased voltage to the dredge resulted in more than standard fishing power, is indicated by the open diamond.




Figure 28. Number per tow of ocean quahogs greater than 50 mm SL from the NEFSC clam survey by region. The 1994 survey, when increased voltage to the dredge resulted in more than standard fishing power, is indicated by the open diamond.




Figure 29. Kilograms per tow of ocean quahogs greater than 50 mm SL from the NEFSC clam survey by region. The 1994 survey, when increased voltage to the dredge resulted in more than standard fishing power, is indicated by the open diamond.




Figure 30. Survey catches of ocean quahogs less than 40mm SL from the 2008 survey (top) and the 2011 survey (bottom), illustrating a recruitment event on GBK where the ocean quahogs are growing large enough to be selected by the survey dredge.




Figure 31A. Number per tow of ocean quahogs less than 40mm shell length from all surveys from 1978 to 1983 combined (top) and from all surveys 1984 to 1989 combined (bottom).




Figure 31B. Number per tow of ocean quahogs less than 40mm shell length from the 1992 survey (top) and from the 1997 and 1999 surveys combined (bottom).




Figure 31C. Number per tow of ocean quahogs less than 40mm shell length from the 2002 and 2005 surveys combined (top) and from the 2008 and 2011 surveys combined (bottom).




Figure 32A. Survey length frequencies for the DMV region. Mean proportion per tow (vertical axis) by shell length class (horizontal axis).




Figure 32B. Survey length frequencies for the NJ region. Mean proportion per tow by shell length class.




Figure 32C. Survey length frequencies for the LI region. Mean proportion per tow by shell length class.




Figure 32D. Survey length frequencies for the SNE region. Mean proportion per tow by shell length class.




Figure 32E. Survey length frequencies for the GBK region. Mean proportion per tow by shell length class.




Figure 33. Mean number of ocean quahogs of all sizes caught per survey tow in any given TMSQ during the time period.




Figure 34. Growth curves (size at age) used in the KLAMZ model for ocean quahogs from the MAB (dashed line), and GBK (solid line). The dotted line is for small ocean quahogs from the MAB and is not used in the model, but may be a more realistic depiction of size of the youngest ocean quahogs.







Figure 35. KLAMZ model results for the DMV region.




Figure 36. KLAMZ biomass and fishing mortality estimates and survey biomass and fishing mortality estimates for the DMV region with 80% confidence intervals. Survey biomass is the efficiency-corrected swept-area biomass (ESB). Survey fishing mortality is calculated by dividing catch (reported landings plus 5% incidental mortality) by ESB.







Figure 37. KLAMZ model results for the NJ region.







Figure 38. KLAMZ model results for the LI region.







Figure 39. KLAMZ model results for the SNE region.






Figure 40. KLAMZ model results for the GBK region.




Figure 41. KLAMZ biomass estimates and survey biomass estimates for the GBK region with 80% confidence intervals.




Figure 42. KLAMZ model population dynamics time series variables (above) and total and recruit biomass (below) for the exploited regions DMV, NJ, LI and SNE combined.




Figure 43. KLAMZ model results for the exploited regions DMV, NJ, LI and SNE combined. Survey observations, 95% confidence interval and fitted values (above) and observed catch data used in the model (below).




Figure 44. KLAMZ model diagnostics for the exploited regions DMV, NJ, LI and SNE combined. Standardized residuals vs. predicted values for survey data (above) and standardized residuals vs. year for survey data (below).




Figure 45. KLAMZ model estimates of fishable biomass for the entire stock (top) and the exploited regions (bottom).





Figure 46. KLAMZ model population dynamics time series variables for the whole stock model (above) and total and recruit biomass (below) from the whole stock model.




Figure 47. KLAMZ model results for the whole stock. Survey observations, 95% confidence interval and fitted values (above) and observed catch data used in the model (below).





Figure 48. KLAMZ model diagnostics for the whole stock model. Standardized residuals vs. predicted values for survey data (above) and standardized residuals vs. year for survey data (below).




Figure 49. KLAMZ estimates of fishing mortality for the entire stock (top) and the exploited regions (bottom).





Figure 50. KLAMZ results for total biomass in the DMV (top) and NJ regions with KLAMZ results from past assessments for comparison. SARC 27 was in 1998, SARC 31 was in 2000, SARC 38 was in 2004, SARC 44 was in 2007 and SARC 48 was in 2009.





Figure 51. KLAMZ results for total biomass in the LI (top) and SNE regions with KLAMZ results from past assessments for comparison. SARC 27 was in 1998, SARC 31 was in 2000, SARC 38 was in 2004, SARC 44 was in 2007 and SARC 48 was in 2009.





Figure 52. KLAMZ results for total biomass in the GBK region (top) and the whole stock minus GBK with KLAMZ results from past assessments for comparison. SARC 27 was in 1998, SARC 31 was in 2000, SARC 38 was in 2004, SARC 44 was in 2007 and SARC 48 was in 2009. Exploited area was not modeled for SARC 27.




Figure 53. KLAMZ results for the whole ocean quahog stock with KLAMZ results from past assessments for comparison. SARC 27 was in 1998, SARC 31 was in 2000, SARC 38 was in 2004, SARC 44 was in 2007 and SARC 48 was in 2009.





Figure 54. KLAMZ results for fishing mortality in the DMV (top) and NJ regions with KLAMZ results from past assessments for comparison. SARC 27 was in 1998, SARC 31 was in 2000, SARC 38 was in 2004, SARC 44 was in 2007 and SARC 48 was in 2009.





Figure 55. KLAMZ results for fishing mortality in the LI (top) and SNE regions with KLAMZ results from past assessments for comparison. SARC 27 was in 1998, SARC 31 was in 2000, SARC 38 was in 2004, SARC 44 was in 2007 and SARC 48 was in 2009.




Figure 56. KLAMZ results for fishing mortality for the whole stock minus GBK (top) and the whole ocean quahog stock with KLAMZ results from past assessments for comparison. SARC 27 was in 1998, SARC 31 was in 2000, SARC 38 was in 2004, SARC 44 was in 2007 and SARC 48 was in 2009.




Figure 57A. Biomass and fishing mortality results from 15-year projections for the whole ocean quahog stock (left side) and the exploited area only (right side) under the “status quo” scenario (15,400 mt, the mean annual catch during 2006 to 2011). The center line is the mean of the bootstrap estimates, the dashed lines are 95% confidence intervals, and the red lines are biomass or fishing mortality thresholds.







Figure 57B. Biomass and fishing mortality results from 15-year projections for the whole ocean quahog stock (left side) and the exploited area only (right side) with the catch fixed at the current quota of 24,190 mt. The center line is the mean of the bootstrap estimates, the dashed lines are 95% confidence intervals, and the red lines are the biomass or fishing mortality thresholds.




Figure 57C. Biomass and catch results from 15-year projections for the whole ocean quahog stock (left side) and the exploited area only (right side) with fishing mortality fixed at the overfishing threshold of $F = 0.022y^{-1}$. The center line is the mean of the bootstrap estimates, the dashed lines are 95% confidence intervals and the red line is the biomass threshold. Probabilities of overfishing are not shown as they are 50% in each year when fishing is at the OFL level.

Appendix 1: Updated Report on the Maine Ocean Quahog Resource and Fishery

Introduction

The Maine fishery for Ocean quahogs, although harvesting the same species as the rest of the EEZ fishery (*Artica islandica*), is prosecuted in a different way and fills a different sector of the shellfish market. Maine “mahogany quahogs” are harvested at a smaller size (starting at 38 mm or 1.5 inches in shell length, SL) and marketed as a less expensive alternative for *Mercenaria mercenaria* for home and restaurant consumption (Maine DMR 2003). The offshore beds targeted by the Maine fishery are made up of small ocean quahogs, the maximum size being only about 75mm.

The Maine fishery began to expand into Federal waters in the 1980s due in part to PSP closures within state waters. In 1990 it was determined that this fishing activity conflicted with the Magnuson-Stevens Fishery Management Conservation Act which calls for the stock to be managed as a unit throughout its range. The Maine fishery was granted experimental status from 1990-1997. In 1998, the Maine fishery was fully incorporated under Amendment 10 of the surfclam/ocean quahog FMP and given an initial annual quota of 100,000 bushels based on historical landings data. There was no independent assessment of the resource available at that time. The State of Maine is responsible under Amendment 10 to certify harvest areas free of PSP and to conduct stock assessments.

In 2002 the State of Maine conducted a pilot survey to assess the distribution and abundance of quahogs along the Maine coast. This survey was a critical first step in establishing distribution, size composition and relative abundance information for the Maine fishery and for directing the design of the current survey work. While this initial survey provided valuable information it did not have the resources to estimate dredge efficiency and therefore was not able to estimate total biomass or biological reference points. The survey conducted in 2005 was focused on estimating dredge efficiency and mapping quahog density on the commercial fishing grounds.

Estimates of biomass and mortality presented in this report are only for the two commercial beds south of Addison (west bed) and Jonesport/GreatWass (east bed), Maine. This approach was chosen due to available resources and because it was conservative. Other quahog beds are known to exist along many parts of the Maine coast. If mortality targets can be met using the estimates from the primary fishing grounds then biomass outside the survey area can act as a *de facto* preserve.

Fishery Data

Data throughout this report are presented in metric units. In the case of landings and LPUE, values are reported in units of Maine bushels, which are about two-thirds the size of the “industry” bushels used as a measure of ocean quahog volume for the rest of the EEZ. To determine the meat yield of a Maine bushel of ocean quahogs, all quahogs

caught by the Maine survey (number and size composition estimated by subsampling) are converted to a total meat weight then divided by the total number of bushels caught (known). Shell length is converted to meat weight using the equation

$$W = 4.97 \times 10^{-6} \times SL^{3.5696}$$

where W is meat weight and SL is shell length. The resulting meat yield (10.8 lbs) is for a Maine bushel of quahogs averaging around 57 mm SL.

Historically the bulk of ocean quahog fishing activity in Maine has taken place on two large quahog beds just off the coast north of 43°50' latitude: the east bed off the town of Addison and the west bed off Great Wass Island. The two beds cover an area of approximately 60 square nautical miles (Figure 1).

Harvesting takes place year round with the highest market demand during the summer holidays (Memorial Day through Labor Day). Most vessels in the Maine fleet are between 10.7-13.7 m (35-45 ft) and classified as “undertonnage” or “small” in issuing permits. All of the vessels use a “dry” dredge (having no hydraulic jets to loosen the sediments) with a cutter bar set by regulation at no more than 0.91 m (36 in). There are no restrictions on any other dimension of the dredge.

There are no size limits for this fishery, and there is no discarding. The ocean quahogs fished from the two commercial beds off Maine are much smaller than most of the rest of the EEZ harvest, averaging between 50 and 60 mm SL (Figure 2), which suits the half-shell market. The ex-vessel price for Maine mahogany quahogs in 2011 was over three dollars per pound of meats, about five times the price paid for ocean quahogs from the rest of the EEZ. Even though size of individual ocean quahogs has increased since the beginning of the fishery, all the catch brought up in a dredge is still acceptable for this market and are kept. The fishery has no regulatory closed days, although the beds are occasionally closed to fishing as a paralytic shellfish poisoning (PSP) precaution. Since the summer of 2007, there have been 77 days when the fishing areas were closed and 145 days when the open fishing areas were severely restricted in size due to PSP precautions. These closures usually happen during the summer months.

Maine ocean quahog landings have trended downwards since 2002 (Table 1, Figure 3). The exception to this trend is in 2006 when landings increased to 124,839 bushels after the re-opening of a highly productive portion of the fishing grounds that had been closed in previous years as a precaution against PSP. After the initial boost to landings from additional fishing ground, landings again began to decline. By the end of 2012 only 65,912 bushels out of a 100,000 bushel quota had been landed.

LPUE has been fairly stable since the early 1990s (Figure 3). Changes in LPUE were often the result of regulatory or fishing practice changes, such as the uptick in 2006 which mirrors the peak in landings from the re-opening of productive beds. Despite the intensity of the fishery, the fact that the LPUE has not fallen may be the result of the fishery moving onto the most productive beds (Figure 1), the clams growing larger in

size, and the reduction in total effort since the early 2000s. Since 2008, the number of valid ocean quahog license holders has fallen from 47 to 29, and hours fished per year by the fleet has dropped by more than half in the past 10 years (Figure 4).

Incidental mortality for Maine ocean quahogs is unknown. This is an important topic for future research, especially since Maine has a high level of dredging activity relative to the size of the fleet. For example, in 2008 the ocean quahog fleet fished approximately 10,776 hours, equivalent to 64,656 tows at ten minutes each. Using standard industry dredge dimensions and tow speeds this level of fishing activity represents 31.42 nm² of bottom swept by commercial dredges per year. Five percent is added to ocean quahog landings before calculating exploitation rates to make up for incidental mortality, but it is just an estimate based on the fishery for larger ocean quahogs in the rest of the EEZ.

Research Surveys

Surveys of the Maine ocean quahog resource were conducted in 2002, 2005, 2006, 2008 and 2011. With the limited funds dedicated for survey work on quahogs, it was decided to focus all survey efforts after 2002 on the two large ocean quahog beds off of Addison (west bed) and Great Wass Island (east bed) that are the primary commercial fishing grounds. Therefore survey estimates of biomass pertain only to these two beds and not to the coast of Maine as a whole. Vessel logbooks and the 2002 independent survey abundance indices show that the majority of the ocean quahog resource, and as a result the majority of fishing activity, occurs here.

The first step in designing the 2005 survey was to establish a 1 km² grid overlay (using Arcview 3.2) over the two beds. Based on number of days at sea available, it was estimated 260 stations could be completed during the survey, so the centers of 260 1 km² grids covering the commercial beds were selected as start points for survey tows. These points were transferred to The Cap'n Voyager Software for use on board the survey vessel.

In 2005 the west bed had been the only open fishing grounds for 3 years due to PSP closures. The east bed had been unfished for 3 years but had previously been a productive ocean quahog fishing ground. The 2006 survey took place 9 months after the east bed reopened. All areas were open during the 2008 and 2011 surveys.

Survey gear and procedures

The 2005 and 2006 surveys were conducted from the commercial vessel F/V Promise Land, a 12.8 m (42 ft) Novi Style dragger piloted by Capt. Michael Danforth. All survey tows during these two years were conducted using a dredge with the following dimensions: cutter bar 0.91 m (36 in), 2.44 m (8 ft) long x 1.83 m (6 ft) wide x 1.22 m (4 ft) high, overall weight 1,361 kg (3,000 lbs), bar spacing all grills 19.05 mm (¾ in). The dredge used by the F/V Promise Land during normal fishing activity was used for the survey. After the 2006 survey, The F/V Promise Land was sold and the captain left the fishery. The vessel contracted for the 2008 and 2011 surveys, The F/V Allyson J4, was about the same size as the F/V Promise Land and the captain, Bruce Porter, had been a

quahog fisherman for 24 years. The dredge used in 2008 and 2011 was also built for commercial use with the same specifications as the dredge on the Promise Land. The only difference between the dredges was a custom extension on the Promise Land dredge that could hold more sediment, making it roughly 400 lbs heavier than the one used in 2008 and 2011 (Figure 5). During tow operations it was noted that the teeth on the cutter bar of the new dredge shined to depth of 3 inches just as they had in the original dredge. From this we assumed that the new dredge was cutting to the same depth as the original. It was also felt that since the survey tows were short (two minutes compared to about ten minutes for a commercial tow) in order to avoid any overfilling and subsequent loss of catch the extensions on the catch box of the original dredge would not give it any advantage over the current dredge.

To conduct a survey station, the vessel approaches the center of the selected tow grid, and if suitable bottom is not present at the predetermined start point, the vessel starts crossing runs within the grid. If towable bottom cannot be found, then the grid location is deemed un-towable, a note is made, and the captain continues on to the next grid. When a suitable tow path is found within a grid the dredge is lowered to the bottom by free-spooling until the ratio of cable length to depth is 3:1. Once the desired cable length is reached the drum is locked, a two minute timer is started and a GPS point taken. The dredge is towed into the current at approximately 3.5 knots for two minutes, a second GPS point is taken and the dredge is brought to the surface. Before it is brought onboard, mud is cleaned from the dredge by steaming in tight circles keeping the dredge in the vessel's prop wash (Figure 6).

Once on board, the dredge is emptied and the catch photographed (Figure 7). The catch is placed on a shaker table (Figure 8), bycatch is noted and all live ocean quahogs are sorted out. A 5 liter subsample of ocean quahogs is taken from each tow to count and measure. The entire catch is processed if it is less than 5 L. The remainder of the catch is placed in calibrated buckets to determine total catch volume. The number of quahogs caught in a tow is estimated by counting the number of clams in the 5 L subsample and expanding to the total volume of the catch. All data are analyzed using Excel with variances calculated using a bootstrap program (10,000 iterations) written by Dr. Yong Chen at the University of Maine, Orono.

Tow distances were determined by The Cap'n Software and were checked using ESRI ArcInfo software. All data from each tow are standardized to a 200 m tow prior to further analysis. Due to a number of reasons such as placement of lobster gear, vessel availability and weather, the number of stations completed per survey has varied from 130 to 183.

Estimating dredge efficiency

Maine dry dredges are less efficient (2 to 17 percent, ME DMR 2003) than the hydraulic dredges used in the rest of the EEZ (up to 95 percent, Medcolf and Caddy 1971). A reliable estimate of dredge efficiency is needed to convert survey densities to a biomass estimate (NEFSC 2004). To assess the efficiency of the Maine dredge, boxcore samples were taken to directly estimate quahog density, then tows were made in the same area

with the survey dredge. Considering only ocean quahogs of sizes available to the dredge, the ratio of density estimated from the dredge tows to density estimated from the boxcore samples is an estimate of survey dredge efficiency (Thorarinsdottir and Jacobson 2005). The estimated dredge efficiency was 17.9 percent, with a 95% bootstrap confidence interval of 8.0%-34.4%. More details of the dredge efficiency experiments can be found in NEFSC 2009, Appendix B2.

Dredge survey results

The original 2005 survey visited 259 potential tow grids. 71% of the stations were towable (183) and 29% were untowable, mostly due to inappropriate substrate, but sometimes due to the presence of fixed gear. During future surveys only the 183 towable grids were revisited. In 2006 130 tows were completed, in 2008 181 tows were completed, and in 2011 183 tows were completed.

Tow distance, catch volume and counts were all standardized to a 200m tow. For all surveys the highest concentration of biomass was in the eastern bed. The eastern section has had the most variable open and close status due to PSP. Substrate data (Figure 9) from Kelly et al. (1998) show the complexity of the substrate in the eastern section with highest quahog densities found near the boundary of hard rocky substrate with gravels, sands or mud. Substrate data collected independently using sidescan imaging show that Kelly et al.'s (1998) substrate information was relatively accurate. However, in some cases substrate labeled as "sand" or "gravel-sand mix" near our most productive tows may have been shell hash from old quahog beds that was seen in boxcores from the same area.

Size frequencies for all subsampled quahogs (n = 20,737 in 2005; 2,014 in 2006; 4,055 in 2008 and 4,316 in 2011) show the quahogs in the eastern bed are larger (Figure 10). Cumulative size frequency distributions and a Kolmogorov-Smirnov test were used to test the null hypothesis that the size frequency distributions in the eastern and western areas were the same (Zar 1999). The null hypothesis was rejected (p=0.001).

Because the two beds have different size compositions and densities, abundance and biomass are calculated separately for the two beds before making combined estimates for the entire survey area. Abundance estimates (see Table 2 and Figure 11) are calculated using a dredge efficiency that was estimated by applying 10,000 bootstrapped efficiency estimates from the three boxcore trips to 10,000 average abundance estimates from the surveys, and the swept area of the survey. Biomass estimates are made by dividing the population into 1 mm size bins based on survey size frequencies, then converting shell length (SL) to meat wet weight (W) using $W=4.97 \times 10^{-6} \times SL^{3.5696}$ (Maine DMR 2003).

Growth and per recruit modeling

A sample of 83 ocean quahogs from the east bed was recently (February 2013) aged at the University of Iowa, and although the data are preliminary, there is evidence that ocean quahogs are growing faster and larger there than previously thought. Kraus et al. (1992) estimated a growth curve for Maine ocean quahogs from the east bed which suggested Maine quahogs grow more slowly and to smaller sizes than ocean quahogs

from the mid-Atlantic bight, while the new data places east bed ocean quahogs in the middle of the EEZ and Kraus et al. curves (Figure 12).

Biological and fishery parameters from a variety of sources were used to carry out a per recruit analysis for ocean quahogs in Maine waters in 2005. The length-based per recruit model used was from the NOAA Fisheries Toolbox (Yield Per Recruit program, internet address: <http://nft.nefsc.noaa.gov>). Age at length and growth information was taken from Kraus et al. (1992). Length-weight parameters were from the 2002 Maine Quahog survey. Size at maturity estimates were based on Rowell et al. (1990) who found that ocean quahog females from Nova Scotia became fully mature at an average size of 49.2 mm. Fishery selectivity was modeled as a linear ramp function that was zero at 37 mm SL and one at 47mm, based on the facility of fitting live ocean quahogs of increasing size through the grates of a commercial dredge by hand (19.05 mm, 3/4 in. bar spacing). Clams from 34mm to 38mm SL generally passed through the grate with some getting caught. After 41mm almost all clams were thick enough to be retained. The regression model for shell depth and shell length in Feindel (2003) shows that a 19.05 mm (¾ in) bar spacing is the thickness of an ocean quahog with 38.7 mm SL. The biological reference points estimated in per recruit modeling for ocean quahog were $F_{max} = 0.0561$, $F_{0.1} = 0.0247$ and $F_{50\%} = 0.013 \text{ y}^{-1}$ (Figure 13). These may be reassessed when the new age and growth data have been fully vetted and can be used in the model.

Sensitivity analysis shows biological reference points from the per recruit model for ocean quahog are most sensitive to fishery selectivity parameters and, in particular, the length at which ocean quahog in Maine waters become fully recruited to the fishery. Commercial port sampling conducted in 2009 confirmed the size selectivity estimates used in the modeling (Figure 2).

Fishing mortality rate

Fishing mortality is estimated as catch in meat weight/average biomass. The survey biomass is used as a proxy for average biomass, as annual mortality rates are low. Following NEFSC (2004), the catch for each year used in fishing mortality estimation was landings plus a 5% allowance for incidental mortality to account for clams that are killed during fishing activity but not harvested. Maine ocean quahog catches, biomass estimates and F estimates for 2005, 2006, 2008 and 2011 are given below.

Year	Median Biomass estimate - mt meat weight	landings in mt meat, full year w/ %5 incidental mortality	F - whole year
2005	25,862	528	0.020
2006	19,012	642	0.033
2008	16,574	348	0.021
2011	19,577	446	0.023

Fishing mortality estimates for 2008 and 2011 are roughly equal to the $F_{0.1}$ generated by the per recruit model, but higher than the $F_{50\%}$.

Stock Status

Since the entire population of ocean quahogs in U.S. waters is managed as a single stock and overfishing definitions apply to the whole stock, it is not possible to evaluate the status of ocean quahogs in Maine as the biomass represents less than 1% of the EEZ stock as a whole. It is not possible to compare or evaluate current biomass levels relative to biological reference points associated with maximum productivity, depleted stock or historical levels because no appropriate biological reference points or historical biomass estimates are available.

$F_{0.1}$ might be a reasonable reference point for managers if the goal is to maximize yield per recruit while preserving some spawning stock. Simulation analysis (Clark 2002) indicates that $F_{50\%}$ (1.3% per year) might be a reasonable reference point for managers if the goal was to preserve enough spawning potential to maintain the resource in the long term. However, preservation of spawning potential may not be necessary if recruitment originates mostly outside of Maine waters.

References

- Chenoweth, S. and D. Dennison. 1993. The Ocean Quahog in Maine- Fishery and Biology. Maine Department of Marine Resources Research Ref. Doc. 93/2.
- Clark, W.G. $F_{35\%}$ revisited ten years later. N. Am. J. Fish. Mgmt. 22:251-257.
- Kelley, J.T., W.A. Barhardt, D.F. Belknap, S.M. Dickson, and A.R. Kelley. 1998. The Seafloor Revealed- The Geology of the Northwestern Gulf of Maine Inner Continental Shelf. Maine Geological Survey, Natural Resources Information and Mapping Center, Dept. of Conservation. Open-File 96-6, 55pp.
- Kraus, M-M.G., B.F.Beal, S.R. Chapman, and L. McMartin. 1992. A comparison of growth rates in *Artica islandica* (Linnaeus, 1767) between field and laboratory populations. J. Shellfish Res. 11(2):289-294.
- Maine Department of Marine Resources. 2003. Gulf of Maine ocean quahog (*Artica islandica*) assessment. Completion report. S. Feindel and D. Schick.
- Medcolf, J.C. and J.F.Caddy. 1971. Underwater observations on the performance of clam dredges of three types. ICES CM B10: 5pp.
- NEFSC (Northeast Fisheries Science Center). 2004. 38th Northeast Regional Stock Assessment Workshop (38th SAW) Stock Assessment Review Committee (SARC) Consensus Summary of Assessments. pp.1-173. NEFSC Ref. Doc.04-03; 246 pp.

NEFSC (Northeast Fisheries Science Center). 2009. 48th Northeast Regional Stock Assessment Workshop (48th SAW) Assessment Report. US Dept Commer, Northeast Fish Sci Cent Ref Doc. 09-15; 834 p.

Rowell, T.W., D.R. Chaisson, and J.T. McLane. 1990. Size and age of sexual maturity and annual gametogenic cycle in the ocean quahog, *Artica islandica* (Linnaeus, 1767), from coastal waters in Nova Scotia, Canada. J. Shellfish Res. Vol.9, No.1, pp195-203.

Thorarinsdottir, G.G. and L. Jacobson. 2005. Fishery biology and biological reference points for management of ocean quahogs (*Artica islandica*) off Iceland. Fisheries Res. 75: 97-106

Zar, J.H 1999. Biostatistical Analysis 4th ed. Prentice Hall, New York.

Table 1. Maine landings, effort and landings per unit effort from vessel logbooks for all vessel classes combined. Only records with both effort and catch data were used to calculate LPUE, which in most years are 100% of the logbook entries as can be seen by comparing columns two and three below.

year	Landings (in Maine bushels)	Landings (records with both effort and catch)	Effort (hours fished)	Nominal LPUE (Maine bushels/hr)
1990	1,018	1,018	286	3.56
1991	36,679	34,360	17,163	2.00
1992	24,839	24,519	13,469	1.82
1993	17,144	17,144	5,748	2.98
1994	21,672	21,672	5,106	4.24
1995	37,912	37,912	5,747	6.60
1996	47,025	47,025	8,483	5.54
1997	72,706	72,706	11,829	6.15
1998	72,466	72,152	11,745	6.14
1999	93,015	92,285	11,151	8.28
2000	121,274	119,103	12,739	9.35
2001	110,272	110,272	13,511	8.16
2002	147,191	147,191	19,681	7.48
2003	119,675	119,675	17,853	6.70
2004	102,187	102,187	19,022	5.37
2005	100,115	100,115	17,063	5.87
2006	121,373	121,373	14,902	8.14
2007	102,006	102,006	14,018	7.28
2008	66,926	66,926	10,776	6.21
2009	56,808	56,808	9,928	5.72
2010	56,469	56,469	9,727	5.81
2011	65,307	65,307	9,145	7.14
2012	65,912	65,912	7,132	9.24

Table 2. Maine ocean quahog survey median abundance and biomass estimates.

Year	Bed	Median abundance estimate - billions	Median biomass estimate - mt meat weight	CV
2005	west	1.729	8,653	39%
	east	2.404	17,208	40%
	combined	4.134	25,862	39%
2006	west	1.996	10,166	41%
	east	1.225	8,846	41%
	combined	3.221	19,012	41%
2008	west	0.711	5,471	40%
	east	1.094	11,103	41%
	combined	1.805	16,574	40%
2011	west	0.754	7,053	40%
	east	1.231	13,277	41%
	combined	1.989	19,577	40%




Figure 1. Locations of all reported commercial landings 2003-2008 (top) and 2009-2012 (bottom).




Figure 2. Size frequency for port samples collected in Jan- March 2009 from 6 different vessels.




Figure 3. Commercial LPUE and Landings from clam industry logbooks.




Figure 4. Number of valid and active (fished during the year) ocean quahog licenses for the Maine fishery, with fleet fishing effort in thousands of hours shown by the dotted line.



Figure 5. At left, the dredge used for the 2005 and 2006 surveys, weighing about 3,000 lbs. At right, the dredge used for the 2008 and 2011 surveys, weighing about 2,600 lbs.



Figure 6. After being brought to the surface, the catch is washed and mud rinsed away in the propeller wash of the survey vessel.



Figure 7. Typical catch from a two-minute survey tow. Note very low bycatch and uniform size of clams.



Figure 8. Processing the survey catch on shaker table, used to remove shell fragments and mud. This step is performed in commercial operations as well.





Figure 9. Substrate data from Kelly et al. (1998) showing coincidence of hard bottom edges with high density quahog tows from eastern bed. The tow locations are the dark blue dots, while pink is rock, light blue is mud, yellow is sand and green is gravel substrate.



2011 Shell Sizes Eastern and Western beds





Figure 10. Growth in Maine ocean quahogs between the 2005 and 2011 surveys for the west bed (top left) and the east bed (top right), and 2011 size frequencies for both beds shown together. These quahogs appear to be growing faster than previous Maine growth data (Kraus et al. 1992) would predict. For instance, the mean growth of 5mm between 2005 and 2008 (three years) in the western bed would be expected to take eight years and the 4.45mm increase during the same three years in the eastern bed would be expected to take 14 years.

Median Maine ocean quahog abundance estimates



Median Maine ocean quahog biomass estimates




Figure 11. Estimates of abundance in billions of individual ocean quahogs (top) and biomass in metric tons of meats for 2005 - 2011.




Figure 12. Ocean quahog shell length at age from a major commercial bed off the Maine coast (east bed), and the mid-Atlantic Bight. The blue symbols represent preliminary new age data from the Maine east bed, and the lines represent growth curves from published studies: Kraus et al. 1992 (Maine east bed, bottom) and NEFSC 2004, (Mid-Atlantic Bight, top).







Figure 13. Results of yield per recruit analysis from 2005.


Appendix 2. Maps of Station Locations and Ocean Quahog Catches from the NEFSC Clam Survey, 1980-2011




Maps of station locations and ocean quahog catches from the 1980 (top) and 1982 (bottom) NEFSC clam surveys. The crosses indicate stations where no ocean quahogs were caught, and the black line designate the assessment regions.




Maps of station locations and ocean quahog catches from the 1983 (top) and 1986 (bottom) NEFSC clam surveys. The crosses indicate stations where no ocean quahogs were caught, and the black line designate the assessment regions.




Maps of station locations and ocean quahog catches from the 1989 (top) and 1992 (bottom) NEFSC clam surveys. The crosses indicate stations where no ocean quahogs were caught, and the black line designate the assessment regions.




Maps of station locations and ocean quahog catches from the 1994 (top) and 1997 (bottom) NEFSC clam surveys. The crosses indicate stations where no ocean quahogs were caught, and the black line designate the assessment regions.



Maps of station locations and ocean quahog catches from the 1999 (top) and 2002 (bottom) NEFSC clam surveys. The crosses indicate stations where no ocean quahogs were caught, and the black line designate the assessment regions.



Maps of station locations and ocean quahog catches from the 2005 (top) and 2008 (bottom) NEFSC clam surveys. The crosses indicate stations where no ocean quahogs were caught, and the black line designate the assessment regions.



Maps of station locations and ocean quahog catches from the 2011 NEFSC clam survey. The crosses indicate stations where no ocean quahogs were caught, and the black line designate the assessment regions.

Procedures for Issuing Manuscripts in the *Northeast Fisheries Science Center Reference Document (CRD) Series*

Clearance

All manuscripts submitted for issuance as CRDs must have cleared the NEFSC's manuscript/abstract/webpage review process. If any author is not a federal employee, he/she will be required to sign an "NEFSC Release-of-Copyright Form." If your manuscript includes material from another work which has been copyrighted, then you will need to work with the NEFSC's Editorial Office to arrange for permission to use that material by securing release signatures on the "NEFSC Use-of-Copyrighted-Work Permission Form."

For more information, NEFSC authors should see the NEFSC's online publication policy manual, "Manuscript/abstract/webpage preparation, review, and dissemination: NEFSC author's guide to policy, process, and procedure," located in the Publications/Manuscript Review section of the NEFSC intranet page.

Organization

Manuscripts must have an abstract and table of contents, and (if applicable) lists of figures and tables. As much as possible, use traditional scientific manuscript organization for sections: "Introduction," "Study Area" and/or "Experimental Apparatus," "Methods," "Results," "Discussion," "Conclusions," "Acknowledgments," and "Literature/References Cited."

Style

The CRD series is obligated to conform with the style contained in the current edition of the United States Government Printing Office Style Manual. That style manual is silent on many aspects of scientific manuscripts. The CRD series relies more on the CSE Style Manual. Manuscripts should be prepared to conform with these style manuals.

The CRD series uses the American Fisheries Society's guides to names of fishes, mollusks, and decapod

crustaceans, the Society for Marine Mammalogy's guide to names of marine mammals, the Biosciences Information Service's guide to serial title abbreviations, and the ISO's (International Standardization Organization) guide to statistical terms.

For in-text citation, use the name-date system. A special effort should be made to ensure that all necessary bibliographic information is included in the list of cited works. Personal communications must include date, full name, and full mailing address of the contact.

Preparation

Once your document has cleared the review process, the Editorial Office will contact you with publication needs – for example, revised text (if necessary) and separate digital figures and tables if they are embedded in the document. Materials may be submitted to the Editorial Office as files on zip disks or CDs, email attachments, or intranet downloads. Text files should be in Microsoft Word, tables may be in Word or Excel, and graphics files may be in a variety of formats (JPG, GIF, Excel, PowerPoint, etc.).

Production and Distribution

The Editorial Office will perform a copy-edit of the document and may request further revisions. The Editorial Office will develop the inside and outside front covers, the inside and outside back covers, and the title and bibliographic control pages of the document.

Once both the PDF (print) and Web versions of the CRD are ready, the Editorial Office will contact you to review both versions and submit corrections or changes before the document is posted online.

A number of organizations and individuals in the Northeast Region will be notified by e-mail of the availability of the document online.

Research Communications Branch
Northeast Fisheries Science Center
National Marine Fisheries Service, NOAA
166 Water St.
Woods Hole, MA 02543-1026

**MEDIA
MAIL**

Publications and Reports of the Northeast Fisheries Science Center

The mission of NOAA's National Marine Fisheries Service (NMFS) is "stewardship of living marine resources for the benefit of the nation through their science-based conservation and management and promotion of the health of their environment." As the research arm of the NMFS's Northeast Region, the Northeast Fisheries Science Center (NEFSC) supports the NMFS mission by "conducting ecosystem-based research and assessments of living marine resources, with a focus on the Northeast Shelf, to promote the recovery and long-term sustainability of these resources and to generate social and economic opportunities and benefits from their use." Results of NEFSC research are largely reported in primary scientific media (*e.g.*, anonymously-peer-reviewed scientific journals). However, to assist itself in providing data, information, and advice to its constituents, the NEFSC occasionally releases its results in its own media. Currently, there are three such media:

NOAA Technical Memorandum NMFS-NE -- This series is issued irregularly. The series typically includes: data reports of long-term field or lab studies of important species or habitats; synthesis reports for important species or habitats; annual reports of overall assessment or monitoring programs; manuals describing program-wide surveying or experimental techniques; literature surveys of important species or habitat topics; proceedings and collected papers of scientific meetings; and indexed and/or annotated bibliographies. All issues receive internal scientific review and most issues receive technical and copy editing.

Northeast Fisheries Science Center Reference Document -- This series is issued irregularly. The series typically includes: data reports on field and lab studies; progress reports on experiments, monitoring, and assessments; background papers for, collected abstracts of, and/or summary reports of scientific meetings; and simple bibliographies. Issues receive internal scientific review and most issues receive copy editing.

Resource Survey Report (formerly *Fishermen's Report*) -- This information report is a regularly-issued, quick-turnaround report on the distribution and relative abundance of selected living marine resources as derived from each of the NEFSC's periodic research vessel surveys of the Northeast's continental shelf. This report undergoes internal review, but receives no technical or copy editing.

TO OBTAIN A COPY of a *NOAA Technical Memorandum NMFS-NE* or a *Northeast Fisheries Science Center Reference Document*, either contact the NEFSC Editorial Office (166 Water St., Woods Hole, MA 02543-1026; 508-495-2350) or consult the NEFSC webpage on "Reports and Publications" (<http://www.nefsc.noaa.gov/nefsc/publications/>). To access *Resource Survey Report*, consult the Ecosystem Surveys Branch webpage (<http://www.nefsc.noaa.gov/femad/ecosurvey/mainpage/>).

ANY USE OF TRADE OR BRAND NAMES IN ANY NEFSC PUBLICATION OR REPORT DOES NOT IMPLY ENDORSEMENT.