



July 11, 2016

CRUISE RESULTS

UNOLS R/V *Hugh R. Sharp*
Cruise No. S1 16-01 (Parts I–III)
Sea Scallop Survey

CRUISE PERIOD AND AREA

The cruise period was 19 May – 24 June 2016 and was conducted in three parts: Part I was from 19 – 27 May, Part II was from June 2 – 9 June, Part IIIa was from 8 - 16 June and Part IIIb was from 17-24 June, 2016. The area surveyed was from the Mid-Atlantic Bight to Georges Bank, and average sampling depths ranged from approximately 33 to 134 meters (18 to 65 fathoms). Approximate station locations are shown in Figures 1 and 2.

OBJECTIVES

The objectives of the survey were to: 1) determine the distribution and relative abundance of sea scallops (*Placopecten magellanicus*) and associated fauna utilizing two sampling devices: the 8-foot wide, standardized sea scallop dredge and the stereo-optic towed camera array (HabCam V4); 2) collect biological samples.

METHODS

Operations and gear for cruise S1 16-01 Parts I – III conformed with the Cruise Instructions for the Sea Scallop Survey, dated May 16, 2016. Exceptions to the Cruise Instructions were during Leg 1, where on 20 May, HabCam separated from the tow cable and vessel. HabCam was recovered on 26 May and necessary repairs made. Leg II departed on 2 June instead of the originally planned date of 29 May.

Pre-selected, random stations were sampled using a modified 2.44 m (8-foot) wide, New Bedford-type scallop dredge rigged with 5.1 cm (2 inch) diameter rings and lined with at 3.8 cm (1½ inch) polyethylene stretched mesh liner. Tow duration was 15 minutes, tow speed was 3.8 knots, and the dredge was fished using a 3.5:1 wire out to depth scope. Tow distance was recorded using differential GPS, and a recording inclinometer was mounted on the dredge to collect bottom-contact time data.

All catch and biological data were recorded using the shipboard, automated, data-entry system, Fisheries Scientific Computing System (F S C S). This system uses digital scales, electronic measuring boards, and touch-screen monitors to record data on deck. After each dredge tow of

the scallop survey, the entire catch was sorted into biological and habitat components. Live whole and clapper shells of various scallop species, including sea and Icelandic, were weighed using the motion-compensated, digital scales. Representative length frequencies for all scallop species were collected to the nearest millimeter using electronic measuring boards (Icthystick); selected fish species incidentally caught in the dredge were also measured to the nearest millimeter. Weights and total numbers were recorded for all other fish species at each station. Furthermore, the weights and total numbers of cancer crabs and starfish were recorded at selected stations. Habitat portions were estimated by basket volume, converted into number of liters (where one basket = 46 liters), recorded into the FSCS system, and then finally discarded. Surface temperatures were measured using the R/V *Hugh R. Sharp*'s hull-mounted temperature sensor and logged by the Scientific Computer System (SCS) at all stations. Temperature and conductivity profiles were made at approximately every fourth dredge station using a conductivity, temperature, and depth instrument (CTD). A bottom salinity sample was obtained twice a day, when applicable, to calibrate the CTD instrument.

Additionally, cooperative work was conducted throughout all legs of the sea scallop survey to determine the most economical and valid methodology for consecutive sampling of the dredge and the Habitat Camera system (HabCam) as a surveying tool, or part of the suite of survey tools, for future NOAA scallop surveys. Developed by a group of researchers associated with Woods Hole Oceanographic Institute (WHOI), as well as in conjunction with particular members of the commercial fishing industry, HabCam is a towed, seafloor-imaging camera system with the following capabilities: 1) acquisition of stereo-optical and acoustic imagery, which can be viewed in "real time"; 2) the ability to count and measure scallops and groundfish; 3) measurement of biodiversity and community structure; 4) characterization of substrate; and 5) measurement of oceanic properties (salinity, temperature, nutrients).

The system is designed to operate over the range of the continental shelf and, while at sea, is able to image a track of over 100 nautical miles each 24-hour day. The current Northeast Fisheries Science Center's 8-foot wide scallop dredge can make approximately 24, 15-minute tows at 3.8 knots per day, covering about 4,500 square meters (m²) per tow and 106,704 m² per day. Continuous operation with HabCam towing at about six knots covers over 260,000 m² per day. Thus, the spatial coverage of HabCam is over 2.5 times the area covered by the survey dredge.

The HabCam system was mounted in a ten-foot long by three-foot wide steel frame and towed one to three meters off the ocean floor at a speed of approximately six knots. An operator controlled the system by means of the R/V *Hugh R. Sharp*'s winch-driven, fiber optic cable, which allowed for real-time data collection and provided power to the unit.

RESULTS

The survey sampled at 152 dredge stations, with 2, 49 and 101 dredge hauls made on Parts I, II and III, respectively; 136 of those stations were representative. The dredge flipped 4 times and stations were re-towed in most flip cases. Bottom temperatures were collected at 16 stations using the CTD system, while bottom water samples for CTD calibration were taken at 4 stations. A total of 2,906 samples were collected to support seven internal and external investigations (Table 1).

During the three legs of the survey, NOAA HabCam V4 was deployed concurrently throughout the scallop strata. HabCam V4 captured images along a cruise track of approximately 483 nm in the Mid-Atlantic Bight (MAB) and 1,148 nm on Georges Bank, including the Great South Channel. The HabCam track is shown in Figure 3. A total of 5.8 million images were collected; 1.6 million images in the MAB, while 4.2 million images were collected on Georges Bank.

DISPOSITION OF DATA

Catch data and hydrographic data will be analyzed at the NEFSC Laboratory in Woods Hole, Massachusetts. The various collections were forwarded to researchers listed in Table 1. Resulting data will be audited, edited, and archived in an Oracle database.

HabCam images will be further analyzed for biological data both at the Woods Hole Oceanographic Institute and at NOAA's NMFS Woods Hole Lab.

SCIENTIFIC PERSONNEL

National Marine Fisheries Service, NEFSC, Woods Hole, MA

Larry Brady³
Jonathan Duquette²
Joseph Godlewski²
Dvora Hart^{1,3b}
Robert Johnston¹
Victor Nordahl³, Chief Scientist¹
Eric Robillard²
Burton Shank²

National Marine Fisheries Service, Cape May, NJ

Joshua O'Connor¹

National Marine Fisheries Service, Hampton, VA

Steven Ellis¹

Contractors, Integrated Statistics, Woods Hole, MA

Michael Bergman^{2,3}
Glenn Chamberlain³
Jui-Han Chang³
Nicole Charriere^{1,3}, Chief Scientist²
Tasha O'Hara², Chief Scientist³
Jill Price^{3b}
Katie Rogers²
Chris Tholke³

WHOI, HabCam Group, Woods Hole, MA

Philip Alatalo²
Scott Gallager^{1,2,3a}
Steve Lerner¹

NOAA Sea Grant

Katherine Kaplan²

Cornell University, Ithaca, NY

NOAA Teacher At Sea

Donna Knutson³

Century High School, Bismarck, ND

Endicott College, Beverly, MA

Lloyd Hayes¹

Old Dominion University, Norfolk, VA

Aris-Aja Horsey¹

University of ME, Fort Kent, ME

Thomas Kaupelis¹

University of ME, Orono, ME

Chen-Tien Hsu^{3b}

University of NH, Durham, NH

Aliza Ray^{3a}

University of NC, Chapel Hill, NC

Justin McNabb¹

Volunteers

Suzanne Rab Green¹

Weehawken, NJ

Jean-Julien Hugo-Webb¹

Acton, MA

Kateryn Thompson-Delgado^{3a}

Bronx, NY

¹ 19 – 27 May, 2016

² 29 May – 7 June, 2016

³ 8 – 24 June, 2016

^{3a} 8 – 16 June, 2016

^{3b} 17 – 24 June, 2016

For further information, contact Robert Johnston, National Marine Fisheries Service, Northeast Fisheries Science Center, Woods Hole, MA 02543. Phone (508) 495-2061; FAX (508) 495-2115; Robert.Johnston@noaa.gov. The Resource Survey Report for this survey and the Cruise Results can be viewed at: [NEFSC Ecosystems Survey Branch website](#).

Table 1. Special samples obtained for various investigators on UNOLS R/V *Hugh R. Sharp* Sea Scallop Survey, during 16 May- 21 June 2017.

Investigator and Affiliation	Samples Saved	Approximate Number
Gallager, Scott WHOI, Woods Hole, MA	sea scallop meats	404 frozen
Hart, Dvora NMFS, NEFSC, Woods Hole, MA	sea scallop shells	530 shells frozen
	sea scallop meat weights	465 weights
	sea scallop gonad weights	468 weights
	sea scallop shell widths	468 examined
	<i>Asterias</i> sp. sea stars	551 examined
Richards, Anne NMFS, NEFSC, Woods Hole, MA	Goosefish	20 frozen whole




Figure 1. Dredge tows made from UNOLS R/V *Hugh R. Sharp*, during NOAA Fisheries Service, Northeast Fisheries Science Center's Sea Scallop Survey, 19 May - 24 June 2016




Figure 2. Dredge tows made from UNOLS R/V *Hugh R. Sharp*, during NOAA Fisheries Service, Northeast Fisheries Science Center's Sea Scallop Survey, 19 May - 24 June 2016




Figure 3. NOAA HabCam track from Mid-Atlantic Bight to Georges Bank, conducted by UNOLS R/V *Hugh R. Sharp* during NOAA Fisheries Service, Northeast Fisheries Science Center's Sea Scallop Survey, 19 May – 24 June 2016.