

perimental evidence, the whole question of the factors influencing the behavior of oyster larvae at the time of setting needs to be examined.

TO THE ANGLE OF SURFACE

French oyster growers take advantage of the preference of oyster larvae for the under surfaces of submerged objects and use special spat collectors made of tiers of tiles set one upon the other with their concave surfaces underneath. New spat is always found in larger numbers on the lower surfaces. According to Cole and Knight-Jones (1939) the larvae of *O. edulis* reared in large tanks in Conway, Wales, set more intensely on the under surfaces of test shells. A study of the effect of the angle of a flat surface on the attachment of larvae was made by Hopkins (1935) in his work on *O. lurida* of the Pacific Coast. He used glass plates, each 2,400 sq. in., placed at different angles over the oyster grounds. The under horizontal surface was designated as 0° and the upper horizontal as 180°. Other plates were set at 45° intervals between the two extremes. The average number of larvae attached to each surface were:

0° (under horizontal)	1,195
45°	181
90°	11
135°	3
180° (upper horizontal)	1

Similar observations were made by Schaefer (1937) with the larvae of *C. gigas*. He set 150 glass plates in positions varying by 45°; some of the plates were parallel to the direction of the tidal current while others were transverse to it. The plates were left for 5 days before the spat were counted. There is a functional relationship between the intensity of setting and the angle of the surface on which the larvae set, the number of larvae being greatest on the under horizontal surface (0° angle) and lowest as the angle approaches 180°. The curve shown in figure 354 is drawn between the points taken as the average numbers of larvae attaching during 5 days on a glass surface held at different angles. The curve is hyperbolic. Schaefer attributes the setting behavior of *C. gigas* to the upward position of the foot of the swimming larvae and possibly to negative geotaxis. No experimental evidence is given to substantiate either point.

The behavior of oyster larvae does not differ from that of many other fouling organisms which




FIGURE 354.—Effect of angle of the surface of plate glass on the number of spat of *C. virginica* which attach to it within 5 days. Glass area 2,400 sq. in.; 0° is under horizontal surface. (Figure 1 from Schaefer, 1937).

were found by Pomerat and Reiner (1942) to attach in greatest abundance to the under surfaces of plates held in a horizontal position.

Contradictory results were obtained, however, in experiments with cement covered boards held at several angles either suspended in water or placed near the bottom. These tests made by Butler (1955) on oyster bottoms near Pensacola, Fla., showed a preponderance of spat on the upper surfaces of the boards. Setting on upper surfaces (135° and 180°) comprised 78 percent of the total number set, and only 18 percent were counted on the lower surfaces. The remaining 4 percent were found on vertical boards. Butler was not able to confirm the results of Pomerat's and Riener's tests made earlier at Pensacola in which frosted glass plates suspended in water attracted the greater percentage of oysters (and barnacles) to the lower surfaces. Bonnot (1937, 1940) built a spat collector from plywood strips dipped in a mixture of cement and sand. He fastened five or six strips of plywood one above the other so that the horizontal surfaces were separated by three-fourths of an inch. Setting of *O. lurida* averaged 98 larva per square inch on the upper surfaces and 67 per square inch on the lower ones. The turbulent current caused by narrow spaces between the strips possibly was

responsible for the greater number of larvae setting on the upper surfaces.

A lack of consistency in observations of various investigators in different environments indicates that it is impossible to ascertain the effects of a single factor of the environment while testing under complex and variable natural conditions. Real progress in the study of the reaction of oyster larvae may be achieved if further observations are made under controlled conditions.

Larvae of *C. virginica* that are grown artificially in culture jars and not disturbed by stirring or aeration are more or less uniformly distributed. Eyed larvae frequently congregate on the surface, swimming with their vela uppermost and touching one another with the tips of the cilia. They form groups or "rafts" visible to the naked eye. Some of them close their valves, fall rapidly to the bottom, and after a short time resume swimming. Falling to the bottom should not be confused with negative geotaxis, which has not been demonstrated for oyster larva. In the laboratory, larvae often attach themselves to the sides of plastic or glass containers and apparently do not discriminate between light and dark surfaces.

TO THE PROPERTIES OF SURFACE

Oyster larvae attach themselves to many kinds of hard and semihard surfaces. They are found on rocks, gravel, cement, wood, shells of other mollusks, on stems and leaves of marsh grass, and on a great variety of miscellaneous objects such as tin cans, rubber boots and tires, glass, tar paper, and pieces of plastic that may be accidentally thrown on the bottom or deliberately used as spat collectors. There is no evidence that the larvae are selective in finding a suitable place to set, provided the surface is not covered with a slimy film, detritus, or soft mud. Under natural conditions they are never found on shifting sand or on a bottom covered with loose sediment. Success of setting always depends primarily on the availability of clean surfaces rather than on other factors. Shells covered with oil and greasy substances in polluted areas are not suitable for the attachment of larvae. Cole and Knight-Jones (1939) found that it is difficult to induce *O. edulis* to set on smooth glass, but the larvae of *C. virginica* raised in the laboratory readily attach to polished glass. In fact live preparations of spat may be obtained for microscopic examination

of small oysters by suspending glass slides in a tank with fully grown larvae.

GREGARIOUSNESS

An interesting gregarious tendency has been observed by Cole and Knight-Jones (1949) among the larvae of *O. edulis*. During experiments in large rearing tanks they found that larvae set more readily on shells already bearing 50 to 100 spat than on shells bearing fewer spat. They suggest that a substance secreted into the surrounding water by the spat, and possibly by the fully developed larvae, encourages the setting. No attempts were made to isolate the substance and test its effect. The authors make another observation which may throw some doubt on the validity of their interpretation. They state (p. 36) that "Larvae set more readily on shells which had remained uncleaned in the tanks for 2 or more weeks, and which bore a visible film of bacteria or diatoms, than on similar shells which were cleaned daily." In their study of gregariousness they placed shells of uniform size and shape in pairs in a tank containing fully developed larvae, and the number of spat attached to them was counted daily. One shell of the pair was considered a control and was cleaned every day, and the other (experimental) remained uncleaned. By the end of the setting period the total spat settled on the experimental shells significantly exceeded the total spat settled on the controls by a ratio of 2.5 to 1. The figures suggest that the observed differences may be due to the attraction of larvae by those which had already settled on the shell, but the conclusion cannot be accepted without further verification. The possibility is not excluded that some other unknown factor, such as the position of the controls in relation to the experimentals, affected the results or that handling and removal of spat from the control shells caused changes to the surface which made them less attractive to the larvae. It would be profitable to conduct a series of tests designed to eliminate bias by placing experimental and control shells at random and making a statistical analysis of the significance of the differences.

Yonge (1960) expresses no doubt "that larvae (of *O. edulis*) settle more readily on surfaces to which others are already attached," and points out that this tendency aids in reproductive efficiency and is, therefore, a major benefit to attached animals. In view of the fact that a single oyster

shell has sufficient space for only a few spat to grow to maturity, heavy concentrations of spat are of doubtful value for reproduction and may even be harmful by creating overcrowded conditions.

ARTIFICIAL REARING OF OYSTER LARVAE

Early attempts to rear oyster larvae under artificial conditions produced uncertain results. Sometimes a small number of spat were obtained, but the experiments could not be repeated under similar conditions. At that time oyster larvae were placed in 5-gallon carboys, and the water was aerated and circulated. At 2-day intervals the larvae were concentrated by centrifuging and transferred into fresh sea water (Wells, 1920). In another method, tried with only partial success, the larvae were reared in slowly running sea water which was filtered through a 2-inch layer of white sand or porous stone (filtrose) placed on the bottom of a container. The rates of filtration and of addition of new water were regulated by a valve placed below the filtering layer (Prytherch, 1924). In both types of experiments no food was added to the containers under the assumption that enough was present in the water. Then Gaarder and Spärck (1933) and Gaarder (1933) studied the food of the larva of *O. edulis* in Norwegian oyster ponds and made what may be considered the first significant step toward solving the problem of rearing larvae under artificial conditions. Spärck observed that the water of the ponds contained considerable numbers of a small green unicellular alga which later on was isolated and cultured in the laboratory. It appeared to be a species of *Chlorella* which was consumed by the larvae. Studies by these investigators revealed also that nannoplankton of the ponds consisted principally of small green algae and flagellates measuring from 2μ to 3μ . Fertilization of experimental tanks by the addition of liquid manure greatly increased the production not only of *Chlorella* but also of various diatoms, chiefly *Nitzschia*, flagellates, various large unicellular green algae, and bacteria. In this enriched water a few larvae grew to a size of 300μ but failed to attach (Spärck, 1927). After it was found that *Chlorella* is present in the Norwegian oyster ponds, in experimental tanks in Conway, Wales, and in certain experimental basins in Denmark, Kändler (1933) attempted to grow oyster larvae on a diet

of this alga alone but had little success. This led him to conclude that oyster larvae are unable to digest *Chlorella*, which left the intestine apparently unchanged. Feeding experiments with *Carteria* and *Chlamydomonas* were also unsuccessful. More critical experiments conducted at Conway, Wales, showed that the larvae are unable to utilize nonmotile green algae such as *Chlorella* and *Collomyxa* but that yellow-brown chrysoomonads (not identified but designated as flagellate C) gave satisfactory results (Cole, 1937).

The Conway experiments demonstrated that organic enrichment of the water of the large tanks was consistently successful in giving rise to a good crop of flagellates with the resulting good growth and setting of larvae (Cole, 1939). The most satisfactory fertilizer was the meat of the shore crab *Carcinus* ground with sand and heated to the boiling point. The suspension of meat was added to a 90,000-gallon tank at the average rate corresponding to 12.5 medium-sized crabs per day for a period of 3 to 4 weeks. Production of nannoplankton was judged by pH readings, and as soon as the readings reached 8.3 to 8.4 and the tank had a distinct slight cloudiness, no more crab meat was added.

Evidence presented by Cole showed that growth and attachment of *O. edulis* larvae in tanks were significantly increased by organic enrichment which stimulated the development of the nannoplankton. Under laboratory conditions the oyster larvae grew and set satisfactorily in the water containing cultures of *Platymonas tetrahele*. The larvae of oysters and other bivalves apparently are not able to swallow microorganisms which exceed 8μ , but according to Thorson (1950) the size of nannoplankton normally devoured by larval forms is smaller (2μ to 3μ).

Difficulties in obtaining reproducible results from using organic enrichment for rearing larvae suggested that variations in the composition and quantity of nannoplankton may be responsible. To determine the food requirements of *O. edulis* larvae, Bruce, Knight, and Parke (1940) isolated from sea water six flagellate organisms ranging in size from 1.5μ to 7μ in diameter. A known number of oyster larvae were introduced into glass vessels filled with 16 l. of uncontaminated, sterile sea water which was stirred and aerated. The water was changed continuously by a drop feed; the loss of larvae was prevented by covering the outflow tubes with bolting silk. The larvae were

fed pure cultures of flagellates grown in the so-called "Erdschreiber" medium of the following composition (Gross, 1937):

Sodium nitrate (NaNO ₃)	0.1 g.
Sodium orthophosphate (Na ₂ HPO ₄)	0.02 g.
Soil extract	50 ml.
Sea water	1,000 ml.

Soil extract is made by boiling 1 kg. of good potting or garden soil with 1 l. of distilled water in an autoclave for 1 hour. The flask is set aside for 2 or 3 days, and the muddy dark fluid is decanted and sterilized by heating to the boiling point. After standing 3 to 4 weeks the suspended particles settle on the bottom, and the transparent brown or red fluid is poured into another container and boiled for a short time. Boiling of the medium should be avoided once the required quantities of nitrates and phosphate have been added.

Since the six flagellates used in these experiments (Bruce, Knight, and Parke, 1940) were not identified and were labeled only by letters, inconsistencies in the results reported may be attributed to the appearance in the culture of other species, or, as the authors state, "to the supervision of factors outside experimental control." The authors suggest that one of these conditions may be the fact that larvae from different oysters are not equally viable.

The feeding of oyster larvae (*O. edulis*) with pure cultures of nannoplankton was repeated by Walne (1956). In this case the larvae were kept in vessels of 1 l. capacity without change of sea water, and the species of flagellates grown in cultures were identified. Among the Chlorophyceae, only *Pyramimonas grossii* Parke gave consistently good results. Tests made with *Chlorella stigmatophora* Butcher seemed to indicate that those chlorococcales which have a thick cell wall are poor food for oyster larvae. The best results were obtained with *Isochrysis galbana* Parke, a chrysophycean of about 5 μ to 6 μ in length. *Prymnesium parvum* Carter was found to be toxic to larvae. So far there is no proof that the species of flagellates used in these experiments form a significant component of the natural population of nannoplankton and that their presence in estuaries is necessary for larvae living under natural conditions.

Imai and Hatanaka (1949, 1950) reported that the larvae of *C. gigas* can be reared on a culture of colorless flagellate, *Monas* sp., which abounds in brackish waters of Japan. The authors believe

that the flagellate of the *Monas* type plays an important role in the production of oysters in Japan. The possibility remains, however, that in their experiments other flagellates were present in the culture of *Monas* enriched with glucose, cane sugar, nitrates, and phosphates.

The pelagic life of *C. virginica* and *C. gigas*, and probably of all oviparous oysters, is longer than that of larviparous *O. edulis* and *O. lurida*. Consequently, the rearing of these oviparous larvae under artificial conditions presents additional difficulties. Considerable advances in the rearing of larvae of various bivalve species were made by Loosanoff, Davis, and their collaborators at the Bureau of Commercial Fisheries Biological Laboratory, Milford, Conn. Phases of the work are summarized by Loosanoff (1954) and Loosanoff and Davis (1963a, 1963b). Oysters were induced to spawn by increasing the temperature and by adding sperm suspension (see p. 308, Chapter XIV). The fertilized eggs were freed from debris by passing the water through a series of fine screens and placed in 5-gallon earthenware jars until free-swimming larvae emerged. Then the water was changed every 24 to 48 hours by straining it through fine sieves which retained the larvae. The sea water in which the larvae lived was filtered through cotton to remove detritus and zooplankton. Aeration and mechanical agitation were considered unnecessary if the water was changed every other day. The larvae were given measured amounts of cultures of various micro-organisms. In general the results obtained in Milford corroborate the findings of British investigators. Davis (1953) established that oyster larvae can utilize as food the following species of flagellates: *Dicrateria inornata*, *Chromulina pleiades*, *Isochrysis galbana*, *Hemiselmis rufescens*, and *Pyramimonas grossii*. *Chlorella* sp. was used only by advanced larval stages and not by young veligers.

The utilizable flagellates were added to the rearing tanks at the rates of 15,000 and 25,000 cells per ml. per day but no toxic effects were noticed in these heavy concentrations, and the larval oyster population of approximately 5,000 per l. showed satisfactory growth. The actual number of flagellates ingested by the larvae was not determined, but the inference was made that "the rate of growth of oyster larvae had an inverse relation to the number of larvae per unit volume" (Davis, 1953). Cole (1939) states that a population of 20,000 to 30,000 small flagellates per 1 ml.

is adequate to promote growth of larvae of *O. edulis* but is insufficient for the spat.

With the exception of the toxic *Prymnesium parvum*, the naked flagellates provided better food for young oyster larvae than the organisms with heavy cell walls, which can be utilized only by older larvae. The best single foods were found to be *Isochrysis galbana*, *Monochrysis lutheri*, *Chromulina pleiades*, *Dicrateria inornata*, and some other unidentified species of *Dicrateria*. Since the cultures used in these experiments were not free of bacteria, the question naturally arises whether the marine bacteria are utilized as food. Davis (1953) states that none of the 13 species of marine bacteria tested by him were used by the larvae. The species of bacteria have not been identified. However, the probability that larvae may derive a certain amount of food from some bacteria is strengthened by the observation reported by Davis (1953) and Loosanoff (1954) that larvae kept in cotton-filtered sea water without algal food continued to grow for as long as 14 to 18 days. The role of marine bacteria in the feeding of oyster larvae needs further experimental study.

Apparently the best results in rearing larvae under artificial conditions are obtained with a mixed food of *Isochrysis galbana*, *Monochrysis lutheri*, *Chromulina pleiades*, and *Dicrateria* sp. With such a diet and at 30° C., the larvae of *C. virginica* begin setting between the 10th and 12th days after fertilization; at 24° C., the sibling larvae are ready to set on the 24th to 26th day; at 20° C., only a few of the larvae set by the 38th day. Setting of larvae of *O. lurida* at a temperature of 22° C. takes place on the 7th day after release of larvae from the brood chamber (Loosanoff and Davis, 1963a).

Under laboratory conditions in Woods Hole the young larvae of *C. virginica* are often found on the bottom of vessels entangled in lumps of several individuals. These larvae never recover and usually die within the next 24 hours. Sometimes the larvae of oysters and clams are attacked and killed by a fungus which has been tentatively identified by the workers at the Bureau of Commercial Fisheries Biological Laboratory at Milford, Conn., as belonging to the genus *Sirolopidium zoophthorum* Vishniac (Davis, Loosanoff, Weston, and Martin, 1954; Johnson and Sparrow, 1961). There are undoubtedly other bacteria and possibly viruses which inflict epizootic mortality on larval

populations in the laboratory and in natural waters.

The technique of rearing oyster larvae has progressed sufficiently to be applicable to practical purposes of oyster culture. Details of techniques, organization, and operation of a mollusk hatchery are summarized by Loosanoff and Davis (1963b).

BIBLIOGRAPHY

- ANDREWS, JAY D.
1951. Seasonal patterns of oyster setting in the James River and Chesapeake Bay. *Ecology*, vol. 32, No. 4, pp. 752-758.
- BERNARD, FÉLIX.
1898. Recherches ontogéniques et morphologiques sur la coquille des lamellibranches. Partie I. Taxodontes et anisomyaires. *Annales des Sciences Naturelles, Zoologie, série 8, tome 8*, pp. 1-208.
- BONNOT, PAUL.
1937. Setting and survival of spat of the Olympia oyster, *Ostrea lurida*, on upper and lower horizontal surfaces. *California Fish and Game*, vol. 23, No. 3, pp. 224-228.
1940. Methods of collecting oyster spat. *Transactions of the American Fisheries Society*, vol. 69, for the year 1939, pp. 263-267.
- BORISIAK, A.
1909. Pelecypoda du plankton de la Mer Noire. *Bulletin Scientifique de la France et de la Belgique*, tome 42, pp. 149-184.
- BRUCE, J. R., MARGERY KNIGHT, and MARY W. PARKE.
1940. The rearing of oyster larvae on an algal diet. *Journal of the Marine Biological Association of the United Kingdom*, vol. 24, No. 1, pp. 337-374.
- BUTLER, PHILIP A.
1955. Selective setting of oyster larvae on artificial cultch. *Proceedings of the National Shellfisheries Association*, vol. 45, August 1954, pp. 95-105.
- CARRIKER, MELBOURNE ROMAINE.
1951. Ecological observations on the distribution of oyster larvae in New Jersey estuaries. *Ecological Monographs*, vol. 21, No. 1, pp. 19-38.
1959. The role of physical and biological factors in the culture of *Crassostrea* and *Mercenaria* in a salt-water pond. *Ecological Monographs*, vol. 29, No. 3, pp. 219-266.
1961. Interrelation of functional morphology, behavior, and autoecology in early stages of the bivalve *Mercenaria mercenaria*. *Journal of the Elisha Mitchell Scientific Society*, vol. 77, No. 2, pp. 168-241.
- CHESTNUT, A. F., and WILLIAM E. FAHY.
1953. Studies on the vertical distribution of setting of oysters in North Carolina. *Proceedings of the Gulf and Caribbean Fisheries Institute, Fifth Annual Session, November 1952*, pp. 106-112.
- COLE H. A.
1937. Experiments in the breeding of oysters (*Ostrea edulis*) in tanks, with special reference to the food of the larva and spat. *Fishery Investigations, series II*, vol. 15, No. 4, 1936, 25 pp.

1938. A system of oyster culture. *Journal du Conseil*, vol. 13, No. 2, pp. 221-235.
- 1938b. The fate of the larval organs in the metamorphosis of *Ostrea edulis*. *Journal of the Marine Biological Association of the United Kingdom*, vol. 22, No. 2, pp. 469-484.
1939. Further experiments in the breeding of oysters, (*Ostrea edulis*) in tanks. *Fishery Investigations*, series II, vol. 16, No. 4, 47 pp.
- COLE, H. A., and E. W. KNIGHT-JONES.
1939. Some observations and experiments on the setting behaviour of larvae of *Ostrea edulis*. *Journal du Conseil*, vol. 14, No. 1, pp. 86-105.
1949. The setting behaviour of larvae of the European flat oyster *Ostrea edulis* L., and its influence on methods of cultivation and spat collection. *Fishery Investigations*, series II, vol. 17, No. 3, 39 pp.
- DANTAN, J. L.
1917. La larve de l'*Ostrea edulis* (L.). *Annales de l'Institut Océanographique*, tome 7, fascicule 6, pp. 1-20.
- DAVAINE, C.
1853. Recherches sur la génération des huîtres. *Comptes Rendus des Séances et Mémoires de la Société de Biologie*, tome 4, série 1, année 1852, pp. 297-339.
- DAVIS, HARRY C.
1953. On food and feeding of larvae of the American oyster, *C. virginica*. *Biological Bulletin*, vol. 104, No. 3, pp. 334-350.
1958. Survival and growth of clam and oyster larvae at different salinities. *Biological Bulletin*, vol. 114, No. 3, pp. 296-307.
- DAVIS, HARRY C., and ALAN D. ANSELL.
1962. Survival and growth of larvae of the European oyster, *O. edulis*, at lowered salinities. *Biological Bulletin*, vol. 122, No. 1, pp. 33-39.
- DAVIS, H. C., V. L. LOOSANOFF, W. H. WESTON, and C. MARTIN.
1954. A fungus disease in clam and oyster larvae. *Science*, vol. 120, No. 3105, pp. 36-38.
- DAWSON, C. E.
1955. A study of the oyster biology and hydrography at Crystal River, Florida. *Publications of the Institute of Marine Science, University of Texas*, vol. 4, No. 1, pp. 279-302.
- DIMICK, R. E., GEORGE EGLAND, and J. B. LONG.
1941. Native oyster investigations of Yaquina Bay, Oregon. Progress Report II covering the period July 4, 1939 to September 30, 1941. Oregon Agricultural Experiment Station, Corvallis, Oreg. Co-operating with the Fish Commission of the State of Oregon and the Lincoln County Court, 153 pp.
- ERDMANN, WILHELM.
1935. Untersuchungen über die Lebensgeschichte der Auster. Nr. 5. Über die Entwicklung und die Anatomie der "ansatzreifen" Larve von *Ostrea edulis* mit Bemerkungen über die Lebensgeschichte der Auster. *Wissenschaftliche Meeresuntersuchungen* herausgegeben von der Kommission zur wissenschaftlichen Untersuchung der deutschen Meere in Kiel und der Biologischen Anstalt auf Helgoland, Neue Folge, Band 19, Abteilung Helgoland, Heft 3, Abhandlung Nr. 6, pp. 1-25.
- FASTEN, NATHAN.
1931. The Yaquina oyster beds of Oregon. *American Naturalist*, vol. 65, No. 700, pp. 434-468.
- FOX, DENIS L., JOHN D. ISAACS, and EUGENE F. CORCORAN.
1952. Marine leptocephala, its recovery, measurement and distribution. *Journal of Marine Research*, vol. 11, No. 1, pp. 29-46.
- FUJITA, TSUNENOBU.
1934. Note on the Japanese oyster larva. *Proceedings of the Fifth Pacific Science Congress*, vol. 5, pp. 4111-4117.
- GAARDER, TORBJØRN.
1933. Untersuchungen über Produktions- und Lebensbedingungen in norwegischen Austern-Pollen. *Bergens Museums Årbok*, 1932, *Naturvidenskapelig Rekke*, nr. 3, pp. 1-64.
- GAARDER, TORBJØRN, and R. SPÆRCK.
1933. Hydrographisch-biochemische Untersuchungen in norwegischen Austern-Pollen. *Bergens Museums Årbok*, 1932, *Naturvidenskapelig Rekke*, nr. 1, pp. 1-144.
- GALTSOFF, PAUL S., and R. H. LUCE.
1930. Oyster investigations in Georgia. [U.S.] Bureau of Fisheries, Report of the Commissioner of Fisheries for the fiscal year 1930, appendix 5 (Document 1077), pp. 61-100.
- GALTSOFF, P. S., H. F. PRYTHERCH, and H. C. McMILLIN.
1930. An experimental study in production and collection of seed oysters. *Bulletin of the U.S. Bureau of Fisheries*, vol. 46, for 1930, pp. 197-263. (Document 1088.)
- GARSTANG, WALTER.
1929. The origin and evolution of larval forms. Sectional Presidents' Addresses, Section D, Zoology. *British Association for the Advancement of Science*, Report of the ninety-sixth meeting (ninety-eighth year), Glasgow, 1928, September 5 to 12. Office of the British Association, Burlington House, London, pp. 77-98.
1951. Larval forms and other zoological verses. *Basil Blackwell*, Oxford, England, 85 pp.
- GROSS, F.
1937. Notes on the culture of some marine plankton organisms. *Journal of the Marine Biological Association of the United Kingdom*, vol. 21, No. 2, pp. 753-768.
- HAGMEIER, A., and A. SCHUBERT.
1930. Untersuchungen über die Biologie der Auster. Nr. 4. Die Austernbrut im Wattenmeer. *Wissenschaftliche Meeresuntersuchungen* herausgegeben von der Kommission zur wissenschaftlichen Untersuchung der deutschen Meere in Kiel und der Biologischen Anstalt auf Helgoland, Neue Folge, Band 18, Abteilung Helgoland, Heft 1, Abhandlung Nr. 1, pp. 1-26.
- HAVINGA, B.
1939. Prediction of the time of setting of oyster spat and a method of control. *Journal du Conseil*, vol. 14, No. 3, pp. 394-400.

- HESS, OSWALD.
1962. Entwicklungsphysiologie der Mollusken. Fortschritte der Zoologie, Band 14, pp. 130-163.
- HOPKINS, A. E.
1935. Attachment of larvae of the Olympia oyster, *Ostrea lurida*, to plane surfaces. Ecology, vol. 16, No. 1, pp. 82-87.
1936. Ecological observations on spawning and early larval development in the Olympia oyster (*Ostrea lurida*). Ecology, vol. 17, No. 4, pp. 551-566.
1937. Experimental observations on spawning, larval development, and setting in the Olympia oyster, (*Ostrea lurida*). [U.S.] Bureau of Fisheries, Bulletin No. 23, vol. 48, pp. 439-503.
- HORI, J.
1926. Notes on the full-grown larva and the Japanese common oyster, *Ostrea gigas* Thunberg. Journal of the Imperial Fisheries Institute, vol. 22, No. 1, pp. 1-7.
- HORST, R.
1882. On the development of the European oyster (*Ostrea edulis* L.). Quarterly Journal of Microscopical Science, vol. 22, pp. 341-346.
1883. A contribution to our knowledge of the development of the oyster (*Ostrea edulis* L.). Bulletin of the U.S. Fish Commission, vol. 2, for 1882, pp. 159-167.
- HUXLEY, T. H.
1883. Oysters and the oyster question. The English Illustrated Magazine, vol. 1, No. 1, pp. 47-55; vol. 1, No. 2, pp. 112-121. Macmillan and Company, London.
- IMAI, TAKEO, and MASAYOSHI HATANAKA.
1949. On the artificial propagation of Japanese common oyster, *Ostrea gigas* Thunb., by non-colored naked flagellates. Bulletin of the Institute of Agricultural Research, Tôhoku University, Sendai, Japan, vol. 1, No. 1, pp. 39-46. [In Japanese with English summary.]
1950. Studies on marine non-colored flagellates, *Monas* sp., favorite food of larvae of various marine animals. I. Preliminary research on cultural requirements. Science Reports of the Tôhoku University, series 4, Biology, vol. 18, No. 3, pp. 304-315.
- INGLE, ROBERT M.
1952. Spawning and setting of oysters in relation to seasonal environmental changes. Bulletin of Marine Science of the Gulf and Caribbean, vol. 1, No. 2, pp. 111-135.
- JACKSON, ROBERT TRACY.
1888. The development of the oyster with remarks on allied genera. Proceedings of the Boston Society of Natural History, vol. 23, pp. 531-556.
1890. Phylogeny of the Pelecypoda. The Aviculidae and their allies. Memoirs of the Boston Society of Natural History, vol. 4, No. 8, pp. 277-400.
- JOHNSON, T. W., JR., and F. K. SPARROW, JR.
1961. Fungi in oceans and estuaries. J. Cramer, Weinheim, Germany; Hafner Publishing Company, New York, 668 pp.
- JØRGENSEN, C. BARKER.
1946. Lamellibranchia. In Gunnar Thorson's Reproduction and larval development of Danish marine bottom invertebrates, with special reference to the planktonic larvae in the Sound (Øresund), ch. 9, pp. 277-311. Meddelelser Fra Kommissionen for Danmarks Fiskeri-og Havundersøgelser, serie: Plankton, bind 4, nr. 1.
- KÄNDLER, RUDOLPH.
1927. Muschellarven aus dem Helgoländer Plankton. Bestimmung ihrer Artzugehörigkeit durch Aufzucht. Wissenschaftliche Meeresuntersuchungen herausgegeben von der Kommission zur wissenschaftlichen Untersuchung der deutschen Meere in Kiel und der Biologischen Anstalt auf Helgoland, Neue Folge, Band 16, Abteilung Helgoland, Heft 2, Abhandlung Nr. 5, pp. 1-8.
1928. Untersuchungen über die Biologie der Auster. Nr. 3, Verbreitung und Wachstum der Austernbrut im Wattenmeer. Planktonuntersuchungen im Sommer 1926 und 1927 nach der Aussetzung holländischer Saataustern. Wissenschaftliche Meeresuntersuchungen herausgegeben von der Kommission zur wissenschaftlichen Untersuchung der deutschen Meere in Kiel und der Biologischen Anstalt auf Helgoland, Neue Folge, Band 17, Abteilung Helgoland, Heft 1, Abhandlung Nr. 3, pp. 1-35.
1933. Die Kultur der Auster. Emil Abderhalden's Handbuch der biologischen Arbeitsmethoden, Abteilung 9, Methoden der Erforschung der Leistungen des tierischen Organismus, Teil 5, Band 1, Methoden der Meerwasserbiologie, pp. 599-716. Urban and Schwarzenberg, Berlin.
- KNIGHT-JONES, E. W.
1952. Gregariousness and some other aspects of the setting behaviour of *Spirobrhis*. Journal of the Marine Biological Association of the United Kingdom, vol. 30, No. 2, pp. 201-222.
- KORRINGA, P.
1941. Experiments and observations on swarming, pelagic life and setting in the European flat oyster, *Ostrea edulis* L. Archives Néerlandaises de Zoologie, tome 5, pp. 1-249.
- LEES, H.
1930. Paper and glass collectors for oyster spat. Journal du Conseil, vol. 5, No. 1, pp. 383-384.
- LOOSANOFF, VICTOR L.
1932. Observations on propagation of oysters in James and Corrotoman Rivers and the seaside of Virginia. The Virginia Commission of Fisheries, Newport News, Va., 46 pp.
1949. Vertical distribution of oyster larvae of different ages during the tidal cycle. [Abstract.] Anatomical Record, vol. 105, No. 3, pp. 591-592.
1954. New advances in the study of bivalve larvae. American Scientist, vol. 42, No. 4, pp. 607-624.
- LOOSANOFF, VICTOR L., and HARRY C. DAVIS.
1963a. Rearing of bivalve mollusks. In F. S. Russell (editor), Advances in marine biology, vol. 1, pp. 1-136. Academic Press, Inc., London.
1963b. Shellfish hatcheries and their future. U.S.

- Fish and Wildlife Service, Commercial Fisheries Review, vol. 25, No. 1, pp. 1-11.
- LOOSANOFF, VICTOR L., and JAMES B. ENGLE.
1940. Spawning and setting of oysters in Long Island Sound in 1937, and discussion of the method for predicting the intensity and time of oyster setting. [U.S.] Bureau of Fisheries, Bulletin No. 33, vol. 49, pp. 217-255.
- MACBRIDE, ERNEST WILLIAM.
1914. Invertebrata. In Walter Heape (editor), Text-book of embryology, vol. 1, 692 pp. Macmillan and Company, Ltd., London.
- MACKIN, JOHN G.
1946. A study of oyster strike on the seaside of Virginia. Virginia Fisheries Laboratory of the College of William and Mary and Commission of Fisheries of Virginia, Contribution No. 25, 18 pp. Commonwealth of Virginia, Division of Purchase and Printing, Richmond, Va.
- MARSHALL, S. M., and A. P. ORR.
1955. The biology of a marine copepod, *Calanus finmarchicus* (Gunnerus). Oliver and Boyd, London, 188 pp.
- MAZZERELLI, GIUSEPPE.
1923. Note sulla biologia dell' *Ostrica (Ostrea edulis* L.). I. Nascita delle larve e durata del periodo larvale. Bollettino della Societa dei Naturalisti in Napoli, vol. 34 (serie 2, vol. 14), pp. 151-159.
- MEDCOF, J. C.
1961. Oyster farming in the Maritimes. Fisheries Research Board of Canada, Bulletin No. 131, 158 pp.
- MEISENHEIMER, JOHANNES.
1901. Entwicklungsgeschichte von *Dreissensia polymorpha* Pall. Zeitschrift für wissenschaftliche Zoologie, Band 69, pp. 1-137.
- NELSON, THURLOW C.
1917. Report of the Department of Biology of the New Jersey Agricultural College Experiment Station, New Brunswick, N.J., for the year ending October 31, 1916, pp. 399-430. MacCrellish and Quigley Company, State Printers, Trenton, N.J.
1921. Aids to successful oyster culture. I. Procuring the seed. New Jersey Agricultural Experiment Stations, Bulletin No. 351, New Brunswick, N.J., 59 pp.
1924. The attachment of oyster larvae. Biological Bulletin, vol. 46, No. 3, pp. 143-151.
1926. Report of the Department of Biology of the New Jersey State Agricultural Experiment Station for the year ending June 30, 1925, pp. 281-288. Published by the State, Trenton, N.J.
1928. Relation of spawning of the oyster to temperature. Ecology, vol. 9, No. 2, pp. 145-154.
- NELSON, THURLOW C., and EARLE B. PERKINS.
1931. Annual report of the Department of Biology, July 1, 1929-June 30, 1930. New Jersey Agricultural Experiment Station, New Brunswick, N.J., Bulletin 522, 47 pp.
- POMERAT, C. M., and E. R. REINER.
1942. The influence of surface angle and of light on the attachment of barnacles and other sedentary organisms. Biological Bulletin, vol. 82, No. 1, pp. 14-25.
- PRITCHARD, D. W.
1952. A review of our present knowledge of the dynamics and flushing of estuaries. Chesapeake Bay Institute of the Johns Hopkins University, Technical Report 4, Reference 52-7, 45 pp.
1953. Distribution of oyster larvae in relation to hydrographic conditions. Proceedings of the Gulf and Caribbean Fisheries Institute, Fifth Annual Session, November 1952, pp. 123-132.
1954. A study of flushing in the Delaware model. Chesapeake Bay Institute of the Johns Hopkins University, Technical Report 7, Reference 54-4, 143 pp.
- PRITCHARD, D. W., and RICHARD E. KENT.
1953. The reduction and analysis of data from the James River operation oyster spat. Chesapeake Bay Institute of the Johns Hopkins University, Technical Report 6, Reference 53-12, 92 pp.
- PRYTHERCH, HERBERT F.
1924. Experiments in the artificial propagation of oysters. [U.S.] Bureau of Fisheries, Report of the Commissioner of Fisheries for the fiscal year 1923, appendix 11 (Document 961), pp. 1-14.
1929. Investigation of the physical conditions controlling spawning of oysters and the occurrence, distribution, and setting of oyster larvae in Milford Harbor, Connecticut. Bulletin of the U.S. Bureau of Fisheries, vol. 44, for 1928, pp. 429-503. (Document 1054).
1930. Improved methods for the collection of seed oysters. [U.S.] Bureau of Fisheries. Report of the Commissioner of Fisheries for the fiscal year 1930, appendix 4 (Document 1076), pp. 47-59.
1931. The role of copper in the setting and metamorphosis of the oyster. Science, vol. 73, No. 1894, pp. 429-431.
1934. The role of copper in the setting, metamorphosis, and distribution of the American oyster, *Ostrea virginica*. Ecological Monographs, vol. 4, No. 1, pp. 47-107.
- RANSON GILBERT
1943. Les prodissoconques des Ostréidés actuels et fossiles. Classification et évolution des Ostréidés. Titres et Travaux Scientifiques de M. Gilbert Ranson, pp. 52-58. Masson et Cie, Paris.
1960. Les prodissoconques (coquilles larvaires) des Ostréidés vivants. Bulletin de l'Institut Océanographique No. 1183, 41 pp.
- REES, C. B.
1950. The identification and classification of lamelli-branch larvae. Hull Bulletins of Marine Ecology, vol. 3, No. 19, pp. 73-104.
- RYDER, JOHN A.
1883. On the mode of fixation of the fry of the oyster. Bulletin of the U.S. Fish Commission, vol. 2, for 1882, pp. 383-387.
- SCHAEFER, MILNER B.
1937. Attachment of the larvae of *Ostrea gigas*, the Japanese oyster, to plane surfaces. Ecology, vol. 18, No. 4, pp. 523-527.
1938. The rate of attachment of the larvae of the

- Japanese oyster, *Ostrea gigas*, as related to the tidal periodicity. *Ecology*, vol. 19, No. 4, pp. 543-547.
- SPÄRCK, R.
1925. Studies on the biology of the oyster (*Ostrea edulis*) in the Limfjord, with special reference to the influence of temperature on the sex change. Report of the Danish Biological Station to the Board of Agriculture, 30, 1924, 84 pp.
1927. Studies on the biology of the oyster (*Ostrea edulis*). Report of the Danish Biological Station to the Board of Agriculture, 33, 1927, pp. 43-65.
- STAFFORD, J.
1912. On the recognition of bivalve larvae in plankton collections. Contributions to Canadian Biology being studies from the marine biological stations of Canada, 1906-1910, No. 14, pp. 221-242.
1913. The Canadian oyster, its development, environment and culture. Committee on Fisheries, Game and Fur-bearing Animals, Commission of Conservation, Canada. The Mortimer Company, Ltd., Ottawa, Canada, 159 pp.
- SULLIVAN, CHARLOTTE M.
1948. Bivalve larvae of Malpeque Bay, P. E. I. Fisheries Research Board of Canada, Bulletin No. 77, 36 pp.
- THORSON, GUNNAR.
1946. Reproduction and larval development of Danish marine bottom invertebrates, with special reference to the planktonic larvae in the Sound (Øresund). Meddelelser Fra Kommissionen for Danmarks Fiskeri-Og Havundersøgelser, Serie: Plankton, bind 4, nr. 1, 523 pp. C. A. Reitzels Forlag, Copenhagen.
1950. Reproductive and larval ecology of marine bottom invertebrates. Biological Reviews of the Cambridge Philosophical Society, vol. 25, No. 1, pp. 1-45.
- WALNE, P. R.
1956. Experimental rearing of the larvae of *Ostrea edulis* L. in the laboratory. Fishery Investigations, series 2, vol. 20, No. 9, 23 pp.
- WELLS, WILLIAM FIRTH.
1920. Artificial propagation of oysters. Transactions of the American Fisheries Society, vol. 50, pp. 301-306.
- WERNER, BERNHARD.
1940. Über die Entwicklung und Artunterscheidung von Muschellarven des Nordseeplanktons, unter besonderer Berücksichtigung der Schalenentwicklung. Mit einem Beitrag zur Kenntnis der Laichzeiten der Nordseemuschelein. Zoologische Jahrbücher, Abteilung für Anatomie und Ontogenie der Tiere, Band 66, Heft 1, pp. 1-54.
- WOODS HOLE OCEANOGRAPHIC INSTITUTION.
1952. Marine fouling and its prevention. Prepared for the Bureau of Ships, Navy Department, United States Naval Institute, Annapolis, Maryland, 1952, 388 pp. (Chapter 4, pp. 42-47, Temporal sequences and biotic successions.) Woods Hole Oceanographic Institution, Contribution No. 580.
- YONGE, C. M.
1926. Structure and physiology of the organs of feeding and digestion in *Ostrea edulis*. Journal of the Marine Biological Association of the United Kingdom, vol. 14, No. 2, pp. 295-386.
1959. Evolution within the bivalve mollusca. Proceedings of the XVth International Congress of Zoology, sec. 4, Invertebrate zoology, pp. 367-370.
1960. Oysters. Collins Clear-Type Press, London, 209 pp.
- ZOBELL, CLAUDE E.
1938. The sequence of events in the fouling of submerged surfaces. Official Digest, Federation of Paint and Varnish Production Clubs, September 1938, Scripps Institution of Oceanography, University of California, Contributions, 1938, new series, No. 35, 8 pp. La Jolla, Calif.
- ZOBELL, CLAUDE E., and ESTHER C. ALLEN.
1935. The significance of marine bacteria in the fouling of submerged surfaces. Journal of Bacteriology, vol. 29, No. 3, pp. 239-251.