

PB89141303



NOAA Technical Memorandum NMFS-F/NEC-60

Antarctic Marine Living Resources Program

**Survey of Antarctic Fur Seals
in the South Shetland Islands,
Antarctica, during the 1986-1987
Austral Summer**

**U.S. DEPARTMENT OF COMMERCE
National Oceanic and Atmospheric Administration
National Marine Fisheries Service
Northeast Fisheries Center
Woods Hole, Massachusetts**

September 1988

REPRODUCED BY:
U.S. Department of Commerce
National Technical Information Service
Springfield, Virginia 22161

NTIS

Recent issues in this series

32. **Secondary Production of Benthic Macrofauna at Three Stations of Delaware Bay and Coastal Delaware.** By Stavros Howe and Wayne Leathem. November 1984. ix + 62 p., 6 figs., 19 tables. NTIS Access. No. PB85-145753/AS.
33. **MARMAP Surveys of the Continental Shelf from Cape Hatteras, North Carolina, to Cape Sable, Nova Scotia (1977-1983). Atlas No. 1. Summary of Operations.** By John D. Sibunka and Myron J. Silverman. November 1984. vii + 306 p., 52 figs., 2 tables. NTIS Access. No. PB85-150985/AS.
34. **Oceanology: Biology of the Ocean. Volume 2. Biological Productivity of the Ocean.** By M. E. Vinogradov, editor in chief. First printed by Nauka Press, Moscow, 1977. Translated from the Russian by Albert L. Peabody. January 1985. x + 518 p., 81 figs., 59 tables. NTIS Access. No. PB85-204683/AS.
35. **Annual NEMP Report on the Health of the Northeast Coastal Waters, 1982.** By John B. Pearce, Carl R. Berman, and Marlene R. Rosen, eds., and Robert N. Reid (benthos), Catherine E. Warsh (water quality), and Edith Gould (biological effects), topic coords. January 1985. xi + 68 p., 29 figs., 5 tables. NTIS Access. No. PB85-219129/AS.
36. **Growth and Survival of Larval Fishes in Relation to the Trophodynamics of Georges Bank Cod and Haddock.** By Geoffrey C. Laurence and R. Gregory Lough. January 1985. xvi + 150 p., 67 figs., 15 tables, 1 app. NTIS Access. No. PB85-220093/AS.
37. **Regional Action Plan.** By Bruce E. Higgins, Ruth Rehfus, John B. Pearce, Robert J. Pawlowski, Robert L. Lippson, Timothy Goodger, Susan Mello Roe, and Douglas W. Beach. April 1985. ix + 84 p., 4 figs., 6 tables, 9 app. NTIS Access. No. PB85-219962/AS.
38. **The Shelf/Slope Front South of Nantucket Shoals and Georges Bank as Delineated by Satellite Infrared Imagery and Shipboard Hydrographic and Plankton Observations.** By J. B. Colton, Jr., J. L. Anderson, J. E. O'Reilly, C. A. Evans-Zetlin, and H. G. Marshall. May 1985. vi + 22 p., 14 figs. NTIS Access. No. PB85-221083/AS.
39. **USA Historical Catch Data, 1904-82, for Major Georges Bank Fisheries.** By Anne M. T. Lange and Joan E. Palmer. May 1985. iii + 21 p., 12 figs., 2 tables. NTIS Access. No. PB85-233948/AS.
40. **Indexing the Economic Health of the U.S. Fishing Industry's Harvesting Sector.** By Virgil J. Norton, Morton M. Miller, and Elizabeth Kenney. May 1985. v + 42 p., 44 figs., 25 tables, 1 app. NTIS Access. No. PB85-217958/AS.
41. **Calculation of Standing Stocks and Energetic Requirements of the Cetaceans of the Northeast United States Outer Continental Shelf.** By Robert D. Kenney, Martin A. M. Hyman, and Howard E. Winn. May 1985. iv + 99 p., 1 fig., 5 tables, 1 app. NTIS Access. No. PB85-239937/AS.
42. **Status of the Fishery Resources Off the Northeastern United States for 1985.** By Conservation & Utilization Division, Northeast Fisheries Center. August 1985. iii + 137 p., 46 figs., 49 tables. NTIS Access. No. PB86-125473/AS.
43. **Status of the Fishery Resources Off the Northeastern United States for 1986.** By Conservation & Utilization Division, Northeast Fisheries Center, September 1986. iii + 130 p., 45 figs., 48 tables. NTIS Access. No. PB87-122115/AS.

(continued on inside back cover)

GENERAL DISCLAIMER

This document may be affected by one or more of the following statements

- **This document has been reproduced from the best copy furnished by the sponsoring agency. It is being released in the interest of making available as much information as possible.**
- **This document may contain data which exceeds the sheet parameters. It was furnished in this condition by the sponsoring agency and is the best copy available.**
- **This document may contain tone-on-tone or color graphs, charts and/or pictures which have been reproduced in black and white.**
- **This document is paginated as submitted by the original source.**
- **Portions of this document are not fully legible due to the historical nature of some of the material. However, it is the best reproduction available from the original submission.**



BIBLIOGRAPHIC INFORMATION

PB89-141303

Report Nos: NOAA-TM-NMFS-F/NEC-60

Title: Survey of Antarctic Fur Seals in the South Shetland Islands, Antarctica, during the 1986-1987 Austral Summer.

Date: Sep 88

Authors: J. L. Bengtson, L. M. Ferm, T. J. Harkonen, E. G. Schaner, and B. S. Stewart.

Performing Organization: National Marine Fisheries Service, Woods Hole, MA. Northeast Fisheries Center. **Goeteborg Univ. (Sweden). Tjarno Marine Biological Lab. **Hubbs Marine Research Inst., San Diego, CA.

Type of Report and Period Covered: Technical memo.,

Supplementary Notes: Prepared in cooperation with Goeteborg Univ. (Sweden). Tjarno Marine Biological Lab., and Hubbs Marine Research Inst., San Diego, CA.

NTIS Field/Group Codes: 47D, 57Z

Price: PC A03/MF A01

Availability: Available from the National Technical Information Service, Springfield, VA. 22161

Number of Pages: 19p

Keywords: *Seals (Mammals), *Aquatic animals, Tables (Data), Surveys, Antarctica, South Shetland Islands, Marine biology, Population growth, *Antarctic fur seals, Antarctic Marine Living Resources Program, Arctocephalus gazella.

Abstract: A survey of Antarctic fur seals in the South Shetland Islands, Antarctica, during the 1986/87 austral summer indicated that this species is continuing its population recovery and recolonization of rookery sites following 19th Century commercial exploitation. Twelve fur seal pupping sites were identified, some of which had not previously been reported. The largest pupping sites are at Telmo Island and Cape Shirreff on the north coast of Livingston Island. Total fur seal pup production in the South Shetland Islands in 1986/87 is estimated to be approximately 4000 individuals. Notes on other pinniped species observed during the survey are presented. The optimal sites for combined fur seal and penguin monitoring activities, as part of the Commission for the Conservation of Antarctic Marine Living Resources (CCAMLR) Ecosystem Monitoring Program, are recommended as Seal Island, Elephant Island; Stigant Point, King George Island; and Cape Shirreff/Telmo Island, Livingston Island.



NOAA Technical Memorandum NMFS-F/NEC-60

This TM series is used for documentation and timely communication of preliminary results, interim reports, or special purpose information, and has not received complete formal review, editorial control, or detailed editing.

Antarctic Marine Living Resources Program

Survey of Antarctic Fur Seals in the South Shetland Islands, Antarctica, during the 1986-1987 Austral Summer

**John L. Bengtson¹, Lisa M. Ferm¹, Tero J. Harkonen²,
Everett G. Schaner³, and Brent S. Stewart⁴**

¹ National Marine Mammal Lab., National Marine Fisheries Serv., Seattle, WA 98815

² Tjarno Marine Biological Lab., Univ. of Goteborg, Stromstad, Sweden

³ 8224 Windsor View Terr., Potomac, MD 20854

⁴ Hubbs Marine Research Ctr., San Diego, CA 92109

U. S. DEPARTMENT OF COMMERCE

C. William Verity, Secretary

National Oceanic and Atmospheric Administration

William E. Evans, Administrator


National Marine Fisheries Service

James W. Brennan, Assistant Administrator for Fisheries

Northeast Fisheries Center

Woods Hole, Massachusetts

September 1988



PROGRAM STATEMENT

The U.S. Antarctic Marine Living Resources (AMLR) Program supports U.S. policy regarding the Convention for the Conservation of Antarctic Marine Living Resources (CCAMLR). The CCAMLR is an international agreement that supports an ecosystem approach to the conservation and management of living resources found in ocean areas surrounding the continent of Antarctica. The Convention mandates a management regime committed to applying measures to ensure that harvesting of Antarctic marine living resources, such as finfish and krill, is conducted in a manner that considers ecological relationships among dependent and related species. Member countries of CCAMLR are: Argentina, Australia, Belgium, Brazil, Chile, European Economic Community, France, German Democratic Republic, Federal Republic of Germany, India, Japan, Korea, New Zealand, Norway, Poland, South Africa, Spain, USSR, United Kingdom, and United States.

U.S. objectives for the CCAMLR were established with the signing into law of the Antarctic Marine Living Resources Convention Act of 1984 (P.L. 98-623). The legislation charges the Secretary of Commerce with the design, conduct and implementation of directed scientific research in support of U.S. objectives in the CCAMLR. Responsibility for these activities has been delegated to the National Fisheries Service (NMFS) of NOAA.

The U.S. AMLR Program supports the CCAMLR need for information through analysis of commercial fisheries data and directed research on key species groups in the Antarctic marine ecosystem. This information, along with research done by other member countries, is used by the CCAMLR to detect and record significant changes in critical components of the Antarctic ecosystem. The Scientific Committee of the CCAMLR then makes conservation recommendations to the Commission, which establishes required conservation measures.

The U.S. AMLR Program focuses its field research activities in the southwest Atlantic Ocean, Scotia Arc, and Antarctic Peninsula. Special attention is directed to the vicinity of Bransfield Strait, South Shetland Islands, and the Palmer Archipelago. In addition, the AMLR Program conducts field work in other areas, as needed, to provide comparative data.

ABSTRACT

← A survey of Antarctic fur seals in the South Shetland Islands, Antarctica, during the 1986/87 austral summer indicated that this species is continuing its population recovery and recolonization of rookery sites following 19th Century commercial exploitation. Twelve fur seal pupping sites were identified, some of which had not previously been reported. The largest pupping sites are at Telmo Island and Cape Shirreff on the north coast of Livingston Island. Total fur seal pup production in the South Shetland Islands in 1986/87 is estimated to be approximately 4000 individuals. Notes on other pinniped species observed during the survey are presented. The optimal sites for combined fur seal and penguin monitoring activities, as part of the Commission for the Conservation of Antarctic Marine Living Resources (CCAMLR) Ecosystem Monitoring Program, are recommended as Seal Island, Elephant Island; Stigant Point, King George Island; and Cape Shirreff/Telmo Island, Livingston Island.

↖

TABLE OF CONTENTS

	Page
PROGRAM STATEMENT.....	iii
ABSTRACT	iv
LIST OF TABLES.....	vi
LIST OF FIGURES.....	vi
ACKNOWLEDGEMENTS.....	vii
INTRODUCTION.....	1
METHODS	1
RESULTS	2
DISCUSSION.....	2
LITERATURE CITED.....	4

LIST OF TABLES


	Page
Table 1. Pinniped census in the South Shetland Islands, Antarctica, during the 1986/87 austral summer.....	5
Table 2. Summary of Antarctic fur seal pup production in the South Shetland Islands, Antarctica, in 1986/87.....	6
Table 3. Changes in Antarctic fur seal pup production at important rookery sites in the South Shetland Islands, Antarctica. Sites listed are for those areas for which there are past census data comparable with the 1986/87 survey.....	7

LIST OF FIGURES

	Page
Figure 1. South Shetland Islands, Antarctica study area.....	8

ACKNOWLEDGEMENTS

The authors are grateful to the officers and crew of the USCGC Glacier and the R/V Prof. Siedlecki for excellent logistic support in the field. We also thank M. E. Goebel, R. L. Merrick, S. D. Osmeck, W. D. Shuford, and L. B. Spear for assistance in the field and laboratory. This work was supported by the National Marine Fisheries Service's Antarctic Marine Living Resources Program and the National Science Foundation's Division of Polar Programs.



INTRODUCTION

Antarctic fur seals, Arctocephalus gazella, were commercially harvested and nearly exterminated in the 19th century. In the 1930's small numbers of fur seals (tens of individuals) were once again seen at Bird Island, South Georgia (Bonner, 1968). Since 1956, when censuses were begun in the South Georgia vicinity, the number of pups born at Bird Island and South Georgia increased to a population estimated in the early 1980's at just under one million (Payne, 1977; Bonner, 1981; Laws, 1984). At present, the majority of Antarctic fur seal pups are born at Bird Island, South Georgia, and the Willis Island, although daughter rookeries have been re-established during the past several decades throughout the Scotia Arc and elsewhere (Bonner, 1968; Laws, 1984).

Surveys have been made during the breeding season at several of these sites, but census coverage has been incomplete. Fur seals in the South Shetland Islands have been censused sporadically over the past 80 years, with the first post-exploitation record of successful pup production being reported by O'Gorman (1961) at Cape Shirreff, Livingston Island, in January of 1958. However, no attempts to census the entire archipelago have been made since 1965/66 when Aguayo and Torres (1967) surveyed fur seal abundance.

The survey reported here focused on the South Shetland Islands and had three principal objectives: 1) to document the recolonization and current distribution of Antarctic fur seal rookeries, 2) to estimate the current annual pup production in the archipelago, and 3) to identify fur seal rookeries suitable as potential monitoring sites in the land-based network of the CCAMLR Ecosystem Monitoring Program.

METHODS

Surveys, which were conducted between 23 December 1986 and 12 February 1987 during research cruises aboard the USCGC Glacier, were used to survey all coastlines of the Elephant Island group, including the Seal Islands, Cornwallis, Clarence, Gibbs, Aspland, and O'Brien Islands (Figure 1). Surveys were flown at approximately 100 m altitude, with photographs taken of large groups to assist in estimating group size (e.g., southern elephant seals, Mirounga leonina). Although the survey focused on Antarctic fur seals, all pinnipeds observed hauled out on land were counted. Species which routinely haul out on ice (e.g., crabeater seals, Lobodon carcinophagus, and leopard seals, Hydrurga leptonyx), were sighted on land only infrequently whereas southern elephant seals (hauled out on beaches for their annual molting period) were sighted often. Counts of all pinnipeds were tallied by species, with no differentiation made between age and sex classes except for fur seals. At sites where fur seal pups were present, observers counted living and dead pups by censusing colonies on foot.

Surveys from the R/V Prof. Siedlecki were conducted between 25 January to 12 February 1987 and were performed by navigating inflatable boats near shore. All ice-free coasts of the South Shetland Islands between Low and Smith Islands and King George Island were examined in this fashion (Figure 1). Where fur seals were observed in abundance, observers landed to search for pups by walking through haulout areas.

RESULTS

A total of 12 Antarctic fur seal pupping sites were identified, some of which were not known prior to this survey (Tables 1 and 2). Numbers of Antarctic fur seal pups at various sites ranged between a single pup each at Fildes Peninsula, Desolation Island, and Smith Island, to 1,875 pups (including 215 dead pups) at the Telmo Island north of Livingston Island. Most pupping sites were located on or near King George, Livingston, or Elephant Islands. Only bachelor male fur seals were observed at other sites. No pupping sites were located on the southern coasts of islands along the Bransfield Strait.

Even though the fur seal population is increasing in the South Shetland Islands, not all of the islands known to have had fur seal rookeries prior to exploitation are currently occupied. Our census revealed that although fur seal colonies are being re-established successfully along the northern coastlines of the South Shetland Islands, recolonization of the southern coasts has not yet begun. Southern coasts are known to have been the sites of large fur seal rookeries which were subjected to heavy commercial exploitation in the 1820/21 and 1821/22 seasons (Stackpole, 1955; Bertrand, 1971). In addition, no fur seals were observed at Cornwallis, Clarence, Gibbs, Aspland, or O'Brien Islands. The absence of animals at these islands conforms to previous survey data which indicated few Antarctic fur seals in these areas (Aguayo and Torres, 1967; Hunt 1973; Aguayo, 1978).

DISCUSSION

Our survey results indicate that both the pup production and the distribution of Antarctic fur seals are continuing to increase in the South Shetland Islands (Table 3). Even for those sites in years when data on the number of pups are not available, the total number of individuals observed appears to be increasing in most areas.

The reason for the differences in growth rates of rookery size between the three islands listed in Table 3 is unknown. Whereas rookeries at Elephant and Livingston Islands have expanded significantly over the past two decades, the number of pups born at Stigant Point, King George Island, has remained essentially unchanged. To the human observer, there appears to be sufficient space on the Stigant Point beaches for rookery growth. These areas are currently utilized as haulout areas by bachelor males. Determining the extent to which potential differences in habitat and local prey resources between these sites may influence pup survival and rookery expansion requires further study.

Three fur seal pupping sites were identified as potentially good locations for incorporation into CCAMLR Ecosystem Monitoring Program network: 1) Seal Island, Elephant Island, 2) Stigant Point, King George Island, and 3) Cape Shirreff, Livingston Island. Each of these sites meets the following criteria which are considered important for establishing a field camp for monitoring purposes: 1) at least 100 fur seal pups born annually, 2) at least 10,000 nesting Chinstrap Penguins, *Pygoscelis antarctica*, available for similar monitoring studies, and 3) a suitable camp site available for a field team. Cape Shirreff and Seal Island are particularly well suited as monitoring sites because of their abundance of fur seals and geographic

position in relation to krill fishing areas. The relative advantages of establishing a monitoring program at these sites will be considered further national programs and within the Working Group for the CCAMLR Ecosystem Monitoring Program.

LITERATURE CITED

- Aguayo, L. A. 1978. Present status of the Antarctic fur seal Arctocephalus gazella at South Shetland Islands. *Polar Rec.* 19:167-173.
- Aguayo, L. A., and D. Torres. 1967. Observaciones sobre mamíferos marinos durante la Vigésima Comisión Antártica Chilena. *Rev. Biol. Mar.* 13:1-57.
- Bertrand, K. J. 1971. *Americans in Antarctica, 1775-1948.* Am. Geogr. Soc., New York.
- Bonner, W. N. 1968. The fur seal of South Georgia. *Sci. Rep. Brit. Antarc. Surv.* 56:1-81.
- Bonner, W. N. 1968. Southern fur seals, Arctocephalus (Geoffroy Saint-Hilaire and Cuvier, 1826). Pp. 161-208 in S. H. Ridgway and R. J. Harrison, (Eds.) *Handbook of Marine Mammals*, Vol. 1. Academic Press, London and New York.
- Cattan, P. E., J. V. Yáñez, D. N. Gajardo, and J. C. Cárdenas. 1982. Censo, marcaje y estructura poblacional del lobo fino antártico Arctocephalus gazella (Peters, 1875) en las islas Shetland del Sur, Chile. *Publ. INACH Ser. Cie.* 29:31-38.
- Hunt, J. F. 1973. Observations on the seals of Elephant Island, South Shetland Islands, 1970-71. *Brit. Antarc. Surv. Bull.* 36:99-104.
- Laws, R. M. 1984. Seals. Pp. 621-715 in R. M. Laws, (Ed.) *Antarctic Ecology*, Vol. 2, Academic Press, London and New York.
- Llano, G. A. 1971. Frontispiece. In W. H. Burt, (Ed.) *Antarctic Pinnipedia Antarctic Research Series, Volume 18.* Am. Geophys. Union, Acad. Sci., Nat. Res. Council.
- O'Gorman, F. A. 1961. Fur seals breeding in the Falkland Islands Dependencies. *Nature* 192:914-916.
- Oliva, D. L., L. R. Duran, D. Torres, and M. Gajardo. 1986. Changes in the population structure and growth of the fur seal Arctocephalus gazella (Peters, 1875) in South Shetland Islands. Unpubl. ms.
- Payne, M. R. 1977. Growth of a fur seal population. *Phil. Trans. Roy. Soc. Lond. (B: Biol. Sci.)* 279:67-79.
- Stackpole, E. A. 1955. *The voyage of the Huron and the Huntress.* The Marine Historical Association, No. 29, Mystic, Conn.

Table 1. Pinniped census in the South Shetland Islands, Antarctica, during the 1986/87 austral summer (E=elephant seal, W=Weddell seal, C=crabeater seal, L=leopard seal).

Location	Male	Female	Pups		E	W	C	L
			Ative	Dead				
King George Island	3,339	147	146	12	3,599	456	2	1
Nelson Island	173	0	0	0	403	362	1	1
Robert Island	194	0	0	0	353	141	0	2
Greenwich Island	235	0	0	0	423	40	0	0
Livingston Island	1,974	129	298	0	4,898	264	1	1
Cape Shirreff	1,586	844	673	45	885	134	1	0
Telmo Island	1,607	2,299	1,660	215	3	3	0	0
Snow Island	652	0	0	0	1,494	157	5	0
Smith Island	105	2	1	0	0	8	0	0
Low Island	418	0	0	0	251	34	0	0
Deception Island	520	0	0	0	3	48	1	0
Elephant Island	158	191	235	15	1,315	0	21	4
Seal Island	66	178	241	8	232	4	1	3
Large Leap Island	73	167	254	21	0	0	0	0
Total	11,100	3,957	3,508	316	13,859	1,651	33	12

Table 2. Summary of Antarctic fur seal pup production in the South Shetland Islands, Antarctica, in 1986/87.

Location	Live	Dead	Total
Seal Island (Elephant Island)	241	8	249
Large Leap Island (Elephant Island)	254	21	275
Cape Valentine (Elephant Island)	42	3	45
Cape Lindsey (Elephant Island)	191	12	203
Stinker Point (King George Island)	2	0	2
Stigant Point (King George Island)	145	12	157
Fildes Peninsula (King George Island)	1	0	1
Desolation Island (Livingston Island)	1	0	1
Cape Shirreff (Livingston Island)	673	45	718
Telmo Island (Livingston Island)	1,660	215	1,875
Window Island (Livingston Island)	297	0	297
Smith Island	1	0	1
Total	3,508	316	3,824

Table 3. Changes in Antarctic fur seal pup production at important rookery sites in the South Shetland Islands, Antarctica. Sites listed are for those areas for which there are past census data comparable with the 1986/87 survey.

Location	Date	Pups ¹	Total ²	Source
<u>ELEPHANT ISLAND</u>				
Seal Island ³	7 Jan 66	0	20	Aguayo, 1978
	13 Dec 70	16	62	Hunt, 1973
	24-31 Dec 86	249	493	1986/87 survey
Cape Valentine	16 Feb 66	2	30	Aguayo, 1978
	9 Feb 71	3	100+	Hunt, 1973
	23 Dec 86	45	127	1986/87 survey
Cape Lindsey	16 Feb 66	3	70	Aguayo, 1978
	Feb 71	0	30	Hunt, 1973
	30 Jan 87	203	468	1986/87 survey
<u>KING GEORGE ISLAND</u>				
Stigant Point	Jan 66	0	0	Aguayo, 1978
	16 Jan 70	123	213	Llano, 1971
	Feb 73	80	250	Aguayo, 1978
	Jan 82	168	293	Cattan et al., 1982
	1982/83	123	367	Oliva et al., 1986
	31 Jan 87	157	507	1986/87 survey
<u>LIVINGSTON ISLAND</u>				
Cape Shirreff	14 Jan 58	0	27	O'Gorman, 1961
	3 Feb 59	2	11	O'Gorman, 1961
	1 Jan 66	12	50	Aguayo, 1978
	Feb 71	27	201	Aguayo, 1978
	25 Jan 73	300	1,741	Aguayo, 1978
	Jan 82	180	532	Cattan et al., 1982
	1982/83	--	564	Oliva et al., 1986
	1983/84	248	969	Oliva et al., 1986
	1984/85	384	1,590	Oliva et al., 1986
	2 Feb 87	718	3,148	1986/87 survey
	Window Island	21 Jan 66	50	150
25 Jan 73		70	320	Aguayo, 1978
3 Feb 87		297	646	1986/87 survey

¹Includes both living and dead pups.

²Total Antarctic fur seals, including pups.

³Compares censuses only at the largest of the Seal Islands.




Figure 1. South Shetland Islands, Antarctica study area.

(continued from inside front cover)

44. **NOAA's Northeast Monitoring Program (NEMP): A Report on Progress of the First Five Years (1979-84) and a Plan for the Future.** By Robert N. Reid, Merton C. Ingham, and John B. Pearce, eds., and Catherine E. Warsh (water quality), Robert N. Reid (sediments & bottom organisms), Adriana Y. Cantillo (trace contaminants in tissues), and Edith Gould (biological effects), topic coords. May 1987. xi + 138 p., 13 figs., 1 table, 9 app. NTIS Access. No. PB87-210100.
45. **Food and Distribution of Juveniles of Seventeen Northwest Atlantic Fish Species, 1973-1976.** By Ray E. Bowman, Thomas R. Azarovitz, Esther S. Howard, and Brian P. Hayden. May 1987. xi + 57 p., 10 figs., 19 tables. NTIS Access. No. PB87-215851/AS.
46. **Influence of Freshwater Inflows on Estuarine Productivity.** By James G. Turek, Timothy E. Goodger, Thomas E. Bigford, and John S. Nichols. May 1987. iii + 26 p. NTIS Access. No. PB87-213666/AS.
47. **MARMAP Surveys of the Continental Shelf from Cape Hatteras, North Carolina, to Cape Sable, Nova Scotia (1977-1984). Atlas No. 2. Annual Distribution Patterns of Fish Larvae.** By Wallace W. Morse, Michael P. Fahay, and Wallace G. Smith. May 1987. viii + 215 p., 27 figs., 2 tables. NTIS Access. No. PB87-232831/AS.
48. **Indexed Bibliography of the Bay Scallop (Argopecten irradians).** By Barbara D. Sabo (Gibson) and Edwin W. Rhodes. May 1987. iii + 85 p. NTIS Access. No. PB87-231411/AS.
49. **Northeast Fisheries Center Framework for Inshore Research.** By Research Planning & Coordination Staff, Northeast Fisheries Center. July 1987. vi + 44 p., 2 figs., 2 tables. NTIS Access. No. PB87-232286/AS.
50. **Status of the Fishery Resources Off the Northeastern United States for 1987.** By Conservation & Utilization Division, Northeast Fisheries Center. October 1987. iii + 132 p., 48 figs., 50 tables. NTIS Access. No. PB88-148549.
51. **An Annotated List of the Fishes of Massachusetts Bay.** By Bruce B. Collette and Karsten E. Hartel. February 1988. x + 70 p., 1 fig., 1 table. NTIS Access. No. PB88-179247/AS.
52. **An Evaluation of the Bottom Trawl Survey Program of the Northeast Fisheries Center.** By Survey Working Group, Northeast Fisheries Center. March 1988. ix + 83 p., 33 figs., 13 tables. NTIS Access. No. PB88-201983/AS.
53. **Contaminants in Hudson-Raritan Estuary Water and Influence of Cold Storage upon Its Chemical Composition.** By Anthony Calabrese, Lawrence J. Buckley, and J. Christopher Powell. May 1988. vii + 37 p., 10 figs., 11 tables. NTIS Access. No. PB88-225628/AS.
54. **Epizootic Ulcerative Syndromes in Coastal/Estuarine Fish.** By Carl J. Sindermann. June 1988. v + 37 p., 8 figs., 1 table.
55. **A Plan for Study: Response of the Habitat and Biota of the Inner New York Bight to Abatement of Sewage Sludge Dumping.** By Environmental Processes Division, Northeast Fisheries Center. June 1988. iii + 34 p., 5 figs., 3 tables, 4 app.
56. **Characterization of the Middle Atlantic Water Management Unit of the Northeast Regional Action Plan.** By Anthony L. Pacheco, ed. July 1988. v + 322 p., 136 figs., 21 tables.
57. **An Analysis and Evaluation of Ichthyoplankton Survey Data from the Northeast Continental Shelf Ecosystem.** By Wallace G. Smith, ed. August 1988. xiii + 132 p., 53 figs., 12 tables, 1 app.
58. **An Indexed Bibliography of Northeast Fisheries Center Publications and Reports for 1987.** By Jon A. Gibson. August 1988. iii + 20 p.
59. **Surveys of Breeding Penguins and Other Seabirds in the South Shetland Islands, Antarctica, January-February 1987.** By W. David Shuford and Larry B. Spear. September 1988. vii + 27 p., 14 figs., 1 table.

PUBLICATIONS AND REPORTS OF THE NORTHEAST FISHERIES CENTER

NOAA's National Marine Fisheries Service (NMFS) seeks to "achieve a continued optimum utilization of living resources for the benefit of the Nation." As the research arm of the NMFS's Northeast Region, the Northeast Fisheries Center (NEFC) supports the NMFS mission by "planning, developing, and managing multidisciplinary programs of basic and applied research to: (1) better understand the living marine resources (including marine mammals) of the Northwest Atlantic, and the environmental quality essential for their existence and continued productivity; and (2) describe and provide to management, industry, and the public, options for the utilization and conservation of living marine resources and maintenance of environmental quality which are consistent with national and regional goals and needs, and with international commitments." To provide its data, information, and advice to constituents, the NEFC issues publications and reports in three categories:

Technical Memorandums—Issued irregularly as NOAA Technical Memorandum NMFS-F/NEC series. Series includes data reports of long-term or large area studies; synthesis reports for major resources or habitats; annual reports of assessment or monitoring programs; documentary reports of oceanographic conditions or phenomena; manuals describing field and lab techniques; literature surveys of major resource or habitat topics; findings of task forces or working groups; and summary reports of scientific or technical workshops. Issues do not undergo exhaustive technical review and editing, but are reliable sources of information. Limited free copies are available from authors or the Center. Issues are also available from the National Technical Information Service, 5285 Port Royal Rd., Springfield, VA, 22161.

Research Documents—Issued irregularly as the Woods Hole Laboratory Reference Document, Narragansett Laboratory Reference Document, and Sandy Hook Laboratory Report series. Series include: data reports on field and lab observations or experiments; progress reports on continuing experiments, monitoring, and assessments; and background papers for scientific or technical workshops. Issues receive minimal internal scientific review and no technical editing. No subscriptions. Free distribution of single copies.

Information Reports—Issued in several series, including: Monthly Highlights (monthly); End-of-Year Report (annual); News Release (irregular); Fishermen's Report (up to four times per year); and The Shark Tagger (two times per year). Content is timely, special-purpose data and/or information. Level of scientific review and technical editing varies by series. All series available through free subscription except for The Shark Tagger which is available only to participants in the NMFS Cooperative Shark Tagging Program.

To obtain a copy of a Technical Memorandum or a Research Document, or to subscribe to an Information Report, write: Information Services Section, Northeast Fisheries Center, Water St., Woods Hole, MA 02543. An annual list of NEFC publications and reports is available upon request at the above address.