

## FIN WHALE (*Balaenoptera physalus*): Western North Atlantic Stock


### STOCK DEFINITION AND GEOGRAPHIC RANGE

The Scientific Committee of the International Whaling Commission (IWC) has proposed stock boundaries for North Atlantic fin whales. Fin whales off the eastern United States, Nova Scotia and the southeastern coast of Newfoundland are believed to constitute a single stock under the present IWC scheme (Donovan 1991). However, the stock identity of North Atlantic fin whales has received relatively little attention, and whether the current stock boundaries define biologically isolated units has long been uncertain. The existence of a subpopulation structure was suggested by local depletions that resulted from commercial overharvesting (Mizroch *et al.* 1984).

A genetic study conducted by Bérubé *et al.* (1998) using both mitochondrial and nuclear DNA provided strong support for an earlier population model proposed by Kellogg (1929) and others. This postulates the existence of several subpopulations of fin whales in the North Atlantic and Mediterranean, with limited gene flow among them. Bérubé *et al.* (1998) also proposed that the North Atlantic population showed recent divergence due to climatic changes (i.e., postglacial expansion), as well as substructuring over even relatively short distances. The genetic data are consistent with the idea that different subpopulations use the same feeding ground, a hypothesis that was also originally proposed by Kellogg (1929).

Fin whales are common in waters of the U. S. Atlantic Exclusive Economic Zone (EEZ), principally from Cape Hatteras northward (Figure 1). Fin whales accounted for 46% of the large whales and 24% of all cetaceans sighted over the continental shelf during aerial surveys (CETAP 1982) between Cape Hatteras and Nova Scotia during 1978-82. While much remains unknown, the magnitude of the ecological role of the fin whale is impressive. In this region fin whales are probably the dominant large cetacean species during all seasons, having the largest standing stock, the largest food requirements, and therefore the largest impact on the ecosystem of any cetacean species (Hain *et al.* 1992; Kenney *et al.* 1997).

There is little doubt that New England waters represent a major feeding ground for fin whales. There is evidence of site fidelity by females, and perhaps some segregation by sexual, maturational or reproductive class in the feeding area (Agler *et al.* 1993). Seipt *et al.* (1990) reported that 49% of fin whales sighted on the Massachusetts Bay area feeding grounds were resighted within the same year, and 45% were resighted in multiple years. The authors suggested that fin whales on these grounds exhibited patterns of seasonal occurrence and annual return that in some respects were similar to those shown for humpback whales. This was reinforced by Clapham and Seipt (1991), who showed maternally directed site fidelity for fin whales in the Gulf of Maine. Information on life history and vital rates is also available in data from the Canadian fishery, 1965-1971 (Mitchell 1974). In seven years, 3,528



**Figure 1.** Distribution of fin whale sightings from NEFSC and SEFSC shipboard and aerial surveys during the summers of 1998, 1999, 2002, 2004 and 2006. Isobaths are the 100-m, 1000-m and 4000-m depth contours.

fin whales were taken at three whaling stations. The station at Blandford, Nova Scotia, took 1,402 fin whales.

Hain *et al.* (1992), based on an analysis of neonate stranding data, suggested that calving takes place during October to January in latitudes of the U.S. mid-Atlantic region; however, it is unknown where calving, mating, and wintering occurs for most of the population. Results from the Navy's SOSUS program (Clark 1995) indicate a substantial deep-ocean distribution of fin whales. It is likely that fin whales occurring in the U. S. Atlantic EEZ undergo migrations into Canadian waters, open-ocean areas, and perhaps even subtropical or tropical regions. However, the popular notion that entire fin whale populations make distinct annual migrations like some other mysticetes has questionable support in the data; in the North Pacific, year-round monitoring of fin whale calls found no evidence for large-scale migratory movements (Watkins *et al.* 2000).

## **POPULATION SIZE**

The best abundance estimate available for the western North Atlantic fin whale stock is 2,269 (CV= 0.37). This August 2006 estimate is recent and provides an estimate when the largest portion of the population was within the study area. However, this estimate must be considered extremely conservative in view of the incomplete coverage of the known habitat of the stock and the uncertainties regarding population structure and whale movements between surveyed and unsurveyed areas. Estimates for animals identified as fin whales were calculated separately from animals identified as either fin or sei whales. The final estimate of fin whales was the sum of the estimate of animals identified as fin whales plus a proportion of the estimate of animals identified as fin or sei whales, where the proportion was defined as the percent of fin whales out of the total number of positively identified fin whales and sei whales.

### **Earlier abundance estimates**

An abundance of 2,200 (CV=0.24) fin whales was estimated from a July to September 1995 sighting survey conducted by two ships and an airplane. The survey covered waters from Virginia to the mouth of the Gulf of St. Lawrence (Palka 1995).

An estimate of abundance of 2,814 (CV=0.21) fin whales was derived from a 28 July to 31 August 1999 line-transect sighting survey conducted by a ship and airplane covering waters from Georges Bank to the mouth of the Gulf of St. Lawrence. Shipboard data were analyzed using the modified direct duplicate method (Palka 1995) that accounts for school size bias and for  $g(0)$ , the probability of detecting a group on the track line. Aerial data were not corrected for  $g(0)$  (Palka 2000). As recommended in the GAMMS Workshop Report (Wade and Angliss 1997), estimates older than eight years are deemed unreliable, and should not be used for PBR determinations. Further, due to changes in survey methodology these data should not be used to make comparisons to more current estimates.

### **Recent surveys and abundance estimates**

An abundance estimate of 2,933 (CV=0.49) fin whales was obtained from an aerial survey conducted in August 2002 which covered 7,465 km of trackline over waters from the 1000 m depth contour on the southern edge of Georges Bank to Maine (Table 1; Palka 2006). The value of  $g(0)$  used for this estimation was derived from the pooled data of 2002, 2004 and 2006 aerial survey data.

An abundance estimate of 1,925 (CV=0.55) fin whales was derived from a line-transect sighting survey conducted during 12 June to 4 August 2004 by a ship and plane that surveyed 10,761 km of trackline in waters north of Maryland (38°N) (Table 1; Palka 2006). Shipboard data were collected using the two independent team line transect method and analyzed using the modified direct duplicate method (Palka 1995) accounting for biases due to school size and other potential covariates, reactive movements (Palka and Hammond 2001), and  $g(0)$ , the probability of detecting a group on the track line. Aerial data were collected using the Hiby circle-back line transect method (Hiby 1999) and analyzed accounting for  $g(0)$  and biases due to school size and other potential covariates (Palka 2005). The value of  $g(0)$  used for this estimation was derived from the pooled data of 2002, 2004 and 2006 aerial survey data.

An abundance of 2,269 (CV=0.37) fin whales was estimated from an aerial survey conducted in August 2006 which covered 10,676 km of trackline in the region from the 2000-m depth contour on the southern edge of Georges Bank to the upper Bay of Fundy and to the entrance of the Gulf of St. Lawrence (Table 1; Palka pers. comm.). The value of  $g(0)$  used for this estimation was derived from the pooled data of 2002, 2004 and 2006 aerial survey data.

Table 1. Summary of recent abundance estimates for western North Atlantic fin whales. Month, year, and area covered during each abundance survey, and resulting abundance estimate ( $N_{best}$ ) and coefficient of variation (CV).			
Month/Year	Area	$N_{best}$	CV
Aug 2002	S. Gulf of Maine to Maine	2,933	0.49
Jun-Jul 2004	Gulf of Maine to lower Bay of Fundy	1,925	0.55
Aug 2006	S. Gulf of Maine to upper Bay of Fundy to Gulf of St. Lawrence	2,269	0.37

### Minimum Population Estimate

The minimum population estimate is the lower limit of the two-tailed 60% confidence interval of the log-normally distributed best abundance estimate. This is equivalent to the 20th percentile of the log-normal distribution as specified by Wade and Angliss (1997). The best estimate of abundance for fin whales is 2,269 (CV=0.37). The minimum population estimate for the western North Atlantic fin whale is 1,678.

### Current Population Trend

There are insufficient data to determine population trends for this species.

### CURRENT AND MAXIMUM NET PRODUCTIVITY RATES

Current and maximum net productivity rates are unknown for this stock. Based on photographically identified fin whales, Agler *et al.* (1993) estimated that the gross annual reproduction rate was at 8%, with a mean calving interval of 2.7 years.

For purposes of this assessment, the maximum net productivity rate was assumed to be 0.04. This value is based on theoretical modeling showing that cetacean populations may not grow at rates much greater than 4% given the constraints of their reproductive life history (Barlow *et al.* 1995).

### POTENTIAL BIOLOGICAL REMOVAL

Potential Biological Removal (PBR) is the product of minimum population size, one-half the maximum productivity rate, and a "recovery" factor (MMPA Sec. 3. 16 U.S.C. 1362; Wade and Angliss 1997). The minimum population size is 1,678. The maximum productivity rate is 0.04, the default value for cetaceans. The "recovery" factor, which accounts for endangered, depleted, or threatened stocks, or stocks of unknown status relative to optimum sustainable population (OSP), is assumed to be 0.10 because the fin whale is listed as endangered under the Endangered Species Act (ESA). PBR for the western North Atlantic fin whale is 3.4.

### ANNUAL HUMAN-CAUSED MORTALITY AND SERIOUS INJURY

For the period 2002 through 2006, the minimum annual rate of human-caused mortality and serious injury to fin whales was 2.0 per year (U.S. waters, 1.6; Canadian waters, 0.4). This value includes incidental fishery interaction records, 0.8 (U.S. waters, 0.8; Canadian waters, 0); and records of vessel collisions, 1.2 (U.S. waters, 0.8; Canadian waters, 0.4) (Glass *et al.* 2008). No fishery-related mortality or serious injury to fin whales was observed by NMFS fishery observers during 2002 through 2006.

### Fishery-Related Serious Injury and Mortality

No confirmed fishery-related mortalities or serious injuries of fin whales have been reported in the NMFS Sea Sampling bycatch database. A review of the records of stranded, floating or injured fin whales for the period 2002 through 2006 on file at NMFS found two records with substantial evidence of fishery interactions causing mortality, and two records resulting in serious injury (Table 2), which results in an annual rate of serious injury and mortality of 0.8 fin whales from fishery interactions. While these records are not statistically quantifiable in the same way as the observer fishery records, they give a minimum count of entanglements for the species. In addition to the records above, there were four additional records of entanglement within the period that either lacked substantial

evidence for a serious injury determination, or did not provide the detail necessary to determine if an entanglement had been a contributing factor in the mortality.

Date <sup>a</sup>	Report Type <sup>b</sup>	Age, Sex, Length	Location <sup>a</sup>	Assigned Cause: P=primary, S=secondary		Notes/Observations
				Ship strike	Entang./ Fsh.inter	
7/28/02	mortality	age & sex unknown	Georges Bank		P	Heavy line seen on tail stock; appeared embedded; no gear recovered
2/12/04	serious injury	age & sex unknown	Pea Island, NC		P	Entangled whale noticeably emaciated; no gear recovered
2/25/04	mortality	Adult Female 16.3m	Port Elizabeth, NJ	P		Displaced vertebrae; ruptured aorta
6/30/04	mortality	age & sex unknown 12m (est)	Georges Bank		P	Freshly dead; heavy line constricting mid-section; no gear recovered
9/26/04	mortality	age & sex unknown 15m (est)	St. Johns, NB	P		Fresh carcass on bow of ship
3/26/05	mortality	Adult <sup>c</sup> Female 16.3m	off Virginia Beach, VA	P		Extensive hemorrhaging and vertebral fractures
4/3/05	mortality	Adult <sup>c</sup> Female 18.8m	Southampton, NY	P		Subdermal hemorrhaging
8/23/05	mortality	Juvenile <sup>c</sup> Male 13.7m	Port Elizabeth, NJ	p		Brought in on bow of ship
9/11/05	mortality	Juvenile <sup>c</sup> Male 11 m	Bonne Esperance, QC	P		Bottom jaw completely severed/broken
9/17/06	serious injury	age & sex unknown 18m (est)	off Mt. Desert Rock, ME		P	Pale skin overall; cyamid load at point of attachment; emaciated

- a. The date sighted and location provided in the table are not necessarily when or where the serious injury or mortality occurred; rather, this information indicates when and where the whale was first reported beached, entangled, or injured.
- b. National guidelines for determining what constitutes a serious injury have not been finalized. Interim criteria as established by NERO/NMFS (Nelson *et al.* 2007) have been used here. Some assignments may change as new information becomes available and/or when national standards are established.
- c. The gender and length were misreported in the 2006 Stock Assessment Report. This table shows the correct values.

### Other Mortality

After reviewing NMFS records for 2002 through 2006, six were found that had sufficient information to confirm the cause of death as collisions with vessels (Table 2) (Glass *et al.* 2008). These records constitute an annual rate of serious injury or mortality of 1.2 fin whales from vessel collisions. NMFS data include three additional records of fin whale collisions with vessels, but the available supporting documentation is insufficient to determine if the whales sustained mortal injuries from the encounters. The number of fin whales taken at 3 whaling stations in Canada from 1965 to 1971 totaled 3,528 whales (Mitchell 1974). Reports of non-directed takes of fin whales are fewer over the last two decades than for other endangered large whales such as right and humpback whales.

### STATUS OF STOCK

The status of this stock relative to OSP in the U.S. Atlantic EEZ is unknown, but the species is listed as endangered under the ESA. There are insufficient data to determine the population trend for fin whales. The total level of human-caused mortality and serious injury is unknown. NMFS records represent coverage of only a portion of the area surveyed for the population estimate for the stock. The total U.S. fishery-related mortality and serious injury for this stock derived from the available records is not less than 10% of the calculated PBR, and therefore cannot be considered insignificant and approaching the ZMRG. This is a strategic stock because the fin whale is listed as an endangered species under the ESA. A Draft Recovery Plan for fin whales has been prepared and is available for review (NMFS 2006).

### REFERENCES CITED

- Agler, B. A., R. L. Schooley, S. E. Frohock, S. K. Katona and I. E. Seipt 1993. Reproduction of photographically identified fin whales, *Balaenoptera physalus*, from the Gulf of Maine. *J. Mamm.* 73(3): 577-587.
- Barlow, J., S. L. Swartz, T. C. Eagle and P. R. Wade 1995. U.S. Marine Mammal Stock Assessments: Guidelines for preparation, background, and a summary of the 1995 assessments. NOAA Tech. Memo. NMFS-OPR-6. 73 pp.
- Bérubé, M., A. Aguilar, D. Dendanto, F. Larsen, G. N. d. Sciara, R. Sears, J. Sigurjónsson, J. Urban-R. and P. J. Palsbøll 1998. Population genetic structure of North Atlantic, Mediterranean and Sea of Cortez fin whales, *Balaenoptera physalus* (Linnaeus 1758): analysis of mitochondrial and nuclear loci. *Mol. Ecol.* 15: 585-599.
- CETAP 1982. A characterization of marine mammals and turtles in the mid- and north Atlantic areas of the U.S. outer continental shelf, final report, Cetacean and Turtle Assessment Program, University of Rhode Island. Washington, DC, Bureau of Land Management. #AA551-CT8-48: 576.
- Clapham, P. J. and I. E. Seipt 1991. Resightings of independent fin whales, *Balaenoptera physalus*, on maternal summer ranges. *J. Mamm.* 72: 788-790.
- Clark, C. W. 1995. Application of U.S. Navy underwater hydrophone arrays for scientific research on whales. *Rep. Int. Whal. Comm.* 45: 210-212.
- Donovan, G. P. 1991. A review of IWC stock boundaries. *Rep. Int. Whal. Comm. (Special Issue)* 13: 39-68.
- Glass, A. H., T. V. N. Cole, M. Garron, R. L. Merrick and R. M. Pace, III 2008. Mortality and serious injury determinations for baleen whale stocks along the United States Eastern Seaboard and adjacent Canadian Maritimes, 2002-2006. *Northeast Fish. Sci. Cent. Ref. Doc.* 08-04. 18 pp.
- Hain, J. H. W., M. J. Ratnaswamy, R. D. Kenney and H. E. Winn 1992. The fin whale, *Balaenoptera physalus*, in waters of the northeastern United States continental shelf. *Rep. Int. Whal. Comm.* 42: 653-669.

- Hiby, L. 1999. The objective identification of duplicate sightings in aerial survey for porpoise. Pages 179-189 *in*: G. W. Garner, S. C. Amstrup, J. L. Laake *et al.*, (eds.) Marine Mammal Survey and Assessment Methods. Balkema, Rotterdam.
- Kellogg, R. 1929. What is known of the migration of some of the whalebone whales. *Ann. Rep. Smithsonian Inst.* 1928: 467-494.
- Kenney, R. D., G. P. Scott, T. J. Thompson and H. E. Winn 1997. Estimates of prey consumption and trophic impacts of cetaceans in the USA northeast continental shelf ecosystem. *J. Northwest Fish. Sci.* 22: 155-171.
- Mitchell, E. 1974. Present status of northwest Atlantic fin and other whale stocks. Pages 108-169 *in*: W. E. Schevill, (ed.) *The whale problem: A status report.* Harvard University Press, Cambridge, MA.
- Mizroch, A. A., D. W. Rice and J. M. Breiwick 1984. The fin whale, *Balaenoptera physalus*. *Mar. Fish. Rev.* 46: 20-24.
- Nelson, M., M. Garron, R. L. Merrick, R. M. Pace, III and T. V. N. Cole 2007. Mortality and Serious Injury Determinations for Baleen Whale Stocks Along the United States Eastern Seaboard and Adjacent Canadian Maritimes, 2001-2005. *Northeast Fish. Sci. Cent. Ref. Doc.* 07-05. 18 pp.
- NMFS 2006. Draft recovery plan for the fin whale (*Balaenoptera physalus*). National Marine Fisheries Service, Silver Spring, MD. [http://www.nmfs.noaa.gov/pr/pdfs/recovery/draft\\_finwhale.pdf](http://www.nmfs.noaa.gov/pr/pdfs/recovery/draft_finwhale.pdf)
- Palka, D. 1995. Abundance estimate of the Gulf of Maine harbor porpoise. *Rep. Int. Whal. Comm.* (special issue) 16: 27-50.
- Palka, D. 2000. Abundance of the Gulf of Maine/Bay of Fundy harbor porpoise based on shipboard and aerial surveys during 1999. *Northeast Fish. Sci. Cent. Ref. Doc.* 00-07. 29 pp. <http://www.nefsc.noaa.gov/psb/pubs/palkalabref00-07.pdf>
- Palka, D. L. 2005. Aerial surveys in the northwest Atlantic: estimation of  $g(0)$ . *Proceedings of a Workshop on Estimation of  $g(0)$  in Line-Transect Surveys of Cetaceans*, European Cetacean Society's 18th Annual Conference; Kolmården, Sweden; Mar. 28, 2004.
- Palka, D. L. 2006. Summer abundance estimates of cetaceans in US North Atlantic Navy Operating Areas. *Northeast Fish. Sci. Cent. Ref. Doc.* 06-03. 41 pp. <http://www.nefsc.noaa.gov/nefsc/publications/crd/crd0603/crd0603.pdf>
- Palka, D. L. and P. S. Hammond 2001. Accounting for responsive movement in line transect estimates of abundance. *Can. J. Fish. Aquat. Sci.* 58: 777-787.
- Seipt, I. E., P. J. Clapham, C. A. Mayo and M. P. Hawvermale 1990. Population Characteristics of Individually Identified Fin Whales *Balaenoptera physalus* in Massachusetts Bay. *Fish. Bull.* 88(2): 271-278.
- Wade, P. R. and R. P. Angliss 1997. Guidelines for assessing marine mammal stocks: Report of the GAMMS Workshop April 3-5, 1996, Seattle, Washington. NOAA Tech. Memo. NMFS-OPR-12. 93 pp.
- Watkins, W. A., M. A. Daher, G. M. Reppucci, J. E. George, D. L. Martin, N. A. DiMarzio and D. P. Gannon 2000. Seasonality and distribution of whale calls in the North Pacific. *Oceanography* 13: 62-67.