

HARBOR PORPOISE (*Phocoena phocoena phocoena*): Gulf of Maine/Bay of Fundy Stock

STOCK DEFINITION AND GEOGRAPHIC RANGE

This stock is found in U.S. and Canadian Atlantic waters. The distribution of harbor porpoises has been documented by sighting surveys, strandings and takes reported by NMFS observers in the Sea Sampling Programs. During summer (July to September), harbor porpoises are concentrated in the northern Gulf of Maine and southern Bay of Fundy region, generally in waters less than 150 m deep (Gaskin 1977; Kraus *et al.* 1983; Palka 1995a; Palka 1995b), with a few sightings in the upper Bay of Fundy and on Georges Bank (Palka 2000). During fall (October-December) and spring (April-June), harbor porpoises are widely dispersed from New Jersey to Maine, with lower densities farther north and south. They are seen from the coastline to deep waters (>1800 m; Westgate *et al.* 1998), although the majority of the population is found over the continental shelf. During winter (January to March), intermediate densities of harbor porpoises can be found in waters off New Jersey to North Carolina, and lower densities are found in waters off New York to New Brunswick, Canada. There does not appear to be a temporally coordinated migration or a specific migratory route to and from the Bay of Fundy region. However, during the fall, several satellite tagged harbor porpoises did favor the waters around the 92-m isobath, which is consistent with observations of high rates of incidental catches in this depth range (Read and Westgate 1997). There were two stranding records from Florida during the 1980s (Smithsonian strandings database) and one in 2003 (NE Regional Office/NMFS strandings and entanglement database).

Gaskin (1984, 1992) proposed that there were four separate populations in the western North Atlantic: the Gulf of Maine/Bay of Fundy, Gulf of St. Lawrence, Newfoundland, and Greenland populations. Analyses involving mtDNA (Wang *et al.* 1996; Rosel *et al.* 1999a; 1999b), organochlorine contaminants (Westgate *et al.* 1997; Westgate and Tolley 1999), heavy metals (Johnston 1995), and life history parameters (Read and Hohn 1995) support Gaskin's proposal. Genetic studies using mitochondrial DNA (Rosel *et al.* 1999a) and contaminant studies using total PCBs (Westgate and Tolley 1999) indicate that the Gulf of Maine/Bay of Fundy females were distinct from females from the other populations in the Northwest Atlantic. Gulf of Maine/Bay of Fundy males were distinct from Newfoundland and Greenland males, but not from Gulf of St. Lawrence males according to studies comparing mtDNA (Palka *et al.* 1996; Rosel *et al.* 1999a) and CHLORs, DDTs, PCBs and CHBs (Westgate and Tolley 1999). Nuclear microsatellite markers have also been applied to samples from these four populations, but this analysis failed to detect significant population sub-division in either sex (Rosel *et al.* 1999a). These patterns may be indicative of female philopatry coupled with dispersal of males. Both mitochondrial DNA and microsatellite




Figure 1. Distribution of harbor porpoises from NEFSC and SEFSC shipboard and aerial surveys during the summers of 1995, 1998, 1999, 2002, 2004, 2006, 2007, 2008, 2010, and 2011 and DFO's 2007 TNASS survey. Isobaths are the 100-m, 1000-m, and 4000-m depth contours.

analyses indicate that the Gulf of Maine/Bay of Fundy stock is not the sole contributor to the aggregation of porpoises found off the mid-Atlantic states during winter (Rosel *et al.* 1999a; Hiltunen 2006). Mixed-stock analyses using twelve microsatellite loci in both Bayesian and likelihood frameworks indicate that the Gulf of Maine/Bay of Fundy is the largest contributor (~60%), followed by Newfoundland (~25%) and then the Gulf of St. Lawrence (~12%), with Greenland making a small contribution (<3%). For Greenland, the lower confidence interval of the likelihood analysis includes zero. For the Bayesian analysis, the lower 2.5% posterior quantiles include zero for both Greenland and the Gulf of St. Lawrence. Intervals that reach zero provide the possibility that these populations contribute no animals to the mid-Atlantic aggregation. This report follows Gaskin's hypothesis on harbor porpoise stock structure in the western North Atlantic, where the Gulf of Maine and Bay of Fundy harbor porpoises are recognized as a single management stock separate from harbor porpoise populations in the Gulf of St. Lawrence, Newfoundland, and Greenland.

POPULATION SIZE

To estimate the population size of harbor porpoises in the Gulf of Maine/Bay of Fundy region, line-transect sighting surveys were conducted during the summers of 1991, 1992, 1995, 1999, 2002, 2004, 2006, 2007, and 2011. The best current abundance estimate of the Gulf of Maine/Bay of Fundy harbor porpoise stock is the result of the 2011 survey: 79,883 (CV=0.32).

Earlier abundance estimates

Please see Appendix IV for a summary of abundance estimates, including earlier estimates and survey descriptions.

Recent surveys and abundance estimates

An abundance estimate of 89,054 (CV=0.47) harbor porpoises was generated from an aerial survey conducted in August 2006 using the Hiby circle-back line-transect method (Hiby 1999) and analyzed accounting for $g(0)$ and biases due to school size and other potential covariates (Palka 2005).. This survey covered 10,676 km of trackline in the region from the 2000-m depth contour on the southern edge of Georges Bank to the upper Bay of Fundy and to the entrance of the Gulf of St. Lawrence. (Table 1; NMFS 2006).

An abundance estimate of 12,732 (CV=0.61) harbor porpoises on the Scotian Shelf and in the Gulf of St. Lawrence was generated from the Canadian Trans North Atlantic Sighting Survey in July–August 2007 (and see Lawson and Gosselin 2009). The total estimate of harbor porpoises from the Gulf of Maine/Bay of Fundy, Gulf of St. Lawrence, and Newfoundland stocks was 16,058 (CV=0.50). This aerial survey covered waters from northern Labrador to the Scotian Shelf, providing full coverage of the Atlantic Canadian coast. The abundance estimates from this survey have been corrected for perception and availability bias, when possible. In general, this involved correcting for perception bias using mark-recapture distance sampling (MCDS), and correcting for availability bias using dive/surface times, as reported in the literature, and the Laake *et al.* (1997) analysis method (Lawson and Gosselin 2011).

An abundance estimate of 79,883 (CV=0.32) harbor porpoises was generated from a shipboard and aerial survey conducted during June–August 2011 (Palka 2012). The aerial portion that contributed to the abundance estimate covered 5,313 km of tracklines that were over waters north of New Jersey from the coastline to the 100-m depth contour through the U.S. and Canadian Gulf of Maine and up to and including the lower Bay of Fundy. The shipboard portion covered 3,107 km of tracklines that were in waters offshore of central Virginia to Massachusetts (waters that were deeper than the 100-m depth contour out to beyond the U.S. EEZ). Both sighting platforms used a double-platform team data collection procedure, which allows estimation of abundance corrected for perception bias of the detected species (Laake and Borchers 2004). Estimation of the abundance was based on the independent observer approach assuming point independence (Laake and Borchers 2004) and calculated using the mark-recapture distance sampling option in the computer program Distance (version 6.0, release 2, Thomas *et al.* 2009).

No harbor porpoises were detected in an abundance survey that was conducted concurrently (June–August 2011) in waters between central Virginia and central Florida. This shipboard survey included shelf-break and inner continental slope waters deeper than the 50-m depth contour within the U.S. EEZ. The survey employed the double-platform methodology searching with 25x bigeye binoculars. A total of 4,445 km of tracklines were surveyed, yielding 290 cetacean sightings.

Table 1. Summary of recent abundance estimates for the Gulf of Maine/Bay of Fundy harbor porpoise (<i>Phocoena phocoena phocoena</i>). Month, year, and area covered during each abundance survey and the resulting abundance estimate (N_{best}) and coefficient of variation (CV).			
Month/Year	Area	N_{best}	CV
Aug 2006	S. Gulf of Maine to upper Bay of Fundy to Gulf of St. Lawrence	89,054	0.47
Jul-Aug 2007 ^a	Scotian Shelf and Gulf of St. Lawrence	12,732	0.61
Jul-Aug 2011	Central Virginia to lower Bay of Fundy	79,883	0.32
a. A portion of this survey covered habitat of the Gulf of Maine/Bay of Fundy stock. The estimate also includes animals from the Gulf of St. Lawrence and Newfoundland stocks.			

Minimum Population Estimate

The minimum population estimate is the lower limit of the two-tailed 60% confidence interval of the log-normal distributed best abundance estimate. This is equivalent to the 20th percentile of the log-normal distribution as specified by Wade and Angliss (1997). The best estimate of abundance for harbor porpoises is 79,883 (CV=0.32). The minimum population estimate for the Gulf of Maine/Bay of Fundy harbor porpoise is 61,415.

Current Population Trend

A trend analysis has not been conducted for this stock. The statistical power to detect a trend in abundance for this stock is poor due to the relatively imprecise abundance estimates and long survey interval. For example, the power to detect a precipitous decline in abundance (i.e., 50% decrease in 15 years) with estimates of low precision (e.g., CV > 0.30) remains below 80% (alpha = 0.30) unless surveys are conducted on an annual basis (Taylor *et al.* 2007).

CURRENT AND MAXIMUM NET PRODUCTIVITY RATES

Several attempts have been made to estimate potential population growth rates. Barlow and Boveng (1991), who used a re-scaled human life table, estimated the upper bound of the annual potential growth rate to be 9.4%. Woodley and Read (1991) used a re-scaled Himalayan tahr life table to estimate a likely annual growth rate of 4%. In an attempt to estimate a potential population growth rate that incorporates many of the uncertainties in survivorship and reproduction, Caswell *et al.* (1998) used a Monte Carlo method to calculate a probability distribution of growth rates. The median potential annual rate of increase was approximately 10%, with a 90% confidence interval of 3-15%. This analysis underscored the considerable uncertainty that exists regarding the potential rate of increase in this population. Moore and Read (2008) conducted a Bayesian population modeling analysis to estimate the potential population growth of harbor porpoise in the absence of bycatch mortality. Their method used fertility data, in combination with age-at-death data from stranded animals and animals taken in gillnets, and was applied under two scenarios to correct for possible data bias associated with observed bycatch of calves. Demographic parameter estimates were 'model averaged' across these scenarios. The Bayesian posterior median estimate for potential natural growth rate was 0.046. This last, most recent, value will be the one used for the purpose of this assessment.

POTENTIAL BIOLOGICAL REMOVAL

Potential Biological Removal (PBR) is the product of minimum population size, one-half the maximum productivity rate, and a recovery factor (MMPA Sec. 3. 16 U.S.C. 1362; Wade and Angliss 1997). The minimum population size is 61,415. The maximum productivity rate is 0.046. The recovery factor, which accounts for endangered, depleted, threatened stocks, or stocks of unknown status relative to optimum sustainable population (OSP) is assumed to be 0.5 because the CV of the average mortality estimate is less than 0.3 (Wade and Angliss 1997). PBR for the Gulf of Maine/Bay of Fundy harbor porpoise is 706.

ANNUAL HUMAN-CAUSED MORTALITY AND SERIOUS INJURY

Data to estimate the mortality and serious injury of harbor porpoise come from U.S. and Canadian Sea Sampling Programs, from records of strandings in U.S. and Canadian waters, and from records in the Marine Mammal Authorization Program (MMAP). See Appendix III for details on U.S. fisheries and data sources.

Estimates using Sea Sampling Program and MMAP data are discussed by fishery under the Fishery Information section (Table 2). Strandings records are discussed under the Other Mortality section (Table 3).

The total annual estimated average human-caused mortality is 683 harbor porpoises per year. This is derived from two components: 640 harbor porpoise per year (CV=0.17) from U.S. fisheries using observer and MMAP data, and 43 per year (unknown CV) from Canadian fisheries using observer data.

New Serious Injury Guidelines

NMFS updated its serious injury designation and reporting process, which uses guidance from previous serious injury workshops, expert opinion, and analysis of historic injury cases to develop new criteria for distinguishing serious from non-serious injury (Angliss and DeMaster 1998; Andersen *et al.* 2008; NOAA 2012). NMFS defines serious injury as an “*injury that is more likely than not to result in mortality*”. Injury determinations for stock assessments revised in 2013 or later incorporate the new serious injury guidelines, based on the most recent 5-year period for which data are available.

Fishery Information

Recently, Gulf of Maine/Bay of Fundy harbor porpoise takes have been documented in the U.S. Northeast sink gillnet, mid-Atlantic gillnet, and Northeast bottom trawl fisheries and in the Canadian herring weir fisheries (Table 2). Detailed U.S. fishery information is reported in Appendix III.

Earlier Interactions

One harbor porpoise was observed taken in the Atlantic pelagic drift gillnet fishery during 1991-1998; the fishery ended in 1998. This observed bycatch was notable because it occurred in continental shelf edge waters adjacent to Cape Hatteras (Read *et al.* 1996). Estimated annual fishery-related mortality (CV in parentheses) attributable to this fishery was 0.7 in 1989 (7.00), 1.7 in 1990 (2.65), 0.7 in 1991 (1.00), 0.4 in 1992 (1.00), 1.5 in 1993 (0.34), 0 during 1994-1996 and 0 in 1998. The fishery was closed during 1997. Information on Canadian fisheries that interact with stocks other than the Gulf of Maine/Bay of Fundy stock can be found in Hooker (1997), Lesage *et al.* (2006) and Benjanims *et al.* (2007).

U.S.

Northeast Sink Gillnet

In 1990, an observer program was started by NMFS to investigate marine mammal takes in the Northeast sink gillnet fishery (Appendix III). Bycatch in the northern Gulf of Maine occurs primarily from June to September, while in the southern Gulf of Maine, bycatch occurs from January to May and September to December. During 2008-2012 no serious injuries were observed (Table 2). Estimated annual bycatch (CV in parentheses) from this fishery was 2,900 in 1990 (0.32), 2,000 in 1991 (0.35), 1,200 in 1992 (0.21), 1,400 in 1993 (0.18) (CUD 1994; Bravington and Bisack 1996), 2,100 in 1994 (0.18), 1,400 in 1995 (0.27) (Bisack 1997), 1,200 in 1996 (0.25), 782 in 1997 (0.22), 332 in 1998 (0.46), 270 in 1999 (0.28) (Rossman and Merrick 1999), 507 in 2000 (0.37), 53 (0.97) in 2001, 444 (0.37) in 2002, 592 (0.33) in 2003, 654 (0.36) in 2004, 630 (0.23) in 2005, 514 (0.31) in 2006, 395 (0.37) in 2007, 666 (0.48) in 2008, 591 (0.23) in 2009, 387 (0.27) in 2010, 273 (0.20) in 2011, and 277 (0.59) in 2012 (Table 2; Orphanides 2013, Hatch and Orphanides 2014). There appeared to be no evidence of differential mortality in U.S. or Canadian gillnet fisheries by age or sex in animals collected before 1994, although there was substantial inter-annual variation in the age and sex composition of the bycatch (Read and Hohn 1995). Using observer data collected during 1990-1998 and a logit regression model, females were 11 times more likely to be caught in the offshore southern Gulf of Maine region, males were more likely to be caught in the south Cape Cod region, and the overall proportion of males and females caught in a gillnet and brought back to land were not significantly different from 1:1 (Lamb 2000).

Scientific experiments that demonstrated the effectiveness of pingers in the Gulf of Maine were conducted during 1992 and 1993 (Kraus *et al.* 1997). After the scientific experiments, experimental fisheries were allowed in the general fishery during 1994 to 1997 in various parts of the Gulf of Maine and south of Cape Cod areas. During these experimental fisheries, bycatch rates of harbor porpoises in pingered nets were less than in non-pingered nets.

A study on the effects of two different hanging ratios in the bottom-set monkfish gillnet fishery on the bycatch of cetaceans and pinnipeds was conducted by NEFSC in 2009 and 2010 with 100% observer coverage which took place in both the Northeast and mid-Atlantic gillnet fisheries. Commercial fishing vessels from Massachusetts and New Jersey were used for the study, which took place south of the Harbor Porpoise Take Reduction Cape Cod South Management Area (south of 40° 40' N) in February-April. Researchers purposely picked an area of historically high bycatch rates in order to have a chance of finding a significant difference. Eight research strings of fourteen nets

each were fished and 159 hauls were completed during the course of the 2009–2010 study. Results showed that while a 0.33 mesh performed better at catching commercially important finfish than a 0.50 mesh, there was no statistical difference in cetacean or pinniped bycatch rates between the two hanging ratios. Twelve harbor porpoises were caught in this project in 79 hauls during 2009 and one animal was caught in 72 hauls during the 2010 experiment in the Northeast (A.I.S., Inc. 2010). These animals were included in the observed interactions and added into the total estimates (Table 2), though these animals and the fishing effort from this experiment were not included in the estimation of the bycatch rate that was expanded to the rest of the fishing effort.

Average estimated harbor porpoise mortality and serious injury in the Northeast sink gillnet fishery during 1994-1998, before the Take Reduction Plan, was 1,163 (0.11). The average annual harbor porpoise mortality and serious injury in the Northeast sink gillnet fishery from 2008 - 2012 was 439 (0.18; Table 2).

Mid-Atlantic Gillnet

Before an observer program was in place for this fishery, Polacheck *et al.* (1995) reported one harbor porpoise incidentally taken in shad nets in the York River, Virginia. In July 1993 an observer program was initiated in the mid-Atlantic gillnet fishery by the NEFSC Sea Sampling program (Appendix III). Documented bycatch after 1995 was from December to May. Bycatch estimates were calculated using methods similar to that used for bycatch estimates in the Northeast sink gillnet fishery (Bravington and Bisack 1996; Bisack 1997). During 2008-2012 no serious injuries were observed (Table 2). The estimated annual mortality (CV in parentheses) attributed to this fishery was 103 (0.57) for 1995, 311 (0.31) for 1996, 572 (0.35) for 1997, 446 (0.36) for 1998, 53 (0.49) for 1999, 21 (0.76) for 2000, 26 (0.95) for 2001, unknown in 2002, 76 (1.13) in 2003, 137 (0.91) in 2004, 470 (0.51) in 2005, 511 (0.32) in 2006, 58 (1.03) in 2007, 350 (0.75) in 2008, 201 (0.55) in 2009, 259 (0.88) in 2010, 123 (0.41) in 2011 and 63(0.83;Orphanides 2013; Hatch and Orphanides 2014).

In the Northeast gillnet fishery section above, see the description of the study on the effects of two different hanging rations in the bottom-set gillnet fishery which took place in both the Northeast and mid-Atlantic gillnet fisheries. Ten harbor porpoises were caught in 8 hauls in the mid-Atlantic as part of this experiment (A.I.S., Inc. 2010). Harbor porpoises that were caught in this study were included in the observed interactions and added into the total estimates (Table 2), though these animals and the fishing effort from this experiment were not included in the estimation of the bycatch rate that was expanded to the rest of the fishing effort.

Annual average estimated harbor porpoise mortality and serious injury from the mid-Atlantic gillnet fishery during 1995 to 1998, before the Take Reduction Plan, was 358 (CV=0.20). The average annual harbor porpoise mortality and serious injury in the mid-Atlantic gillnet fishery from 2008–2012 was 199 (0.37) (Table 2).

Northeast Bottom Trawl

This fishery is active in New England waters in all seasons. Twenty harbor porpoise mortalities were observed in the Northeast bottom trawl fishery between 1989 and 2008, but many of these are not attributable to this fishery. Decomposed animals are presumed to have been dead prior to being taken by the trawl. One fresh dead take was observed in the Northeast bottom trawl fishery in 2003, 4 in 2005, 1 in 2006, 1 in 2008, and 1 in 2011. Revised serious injury guidelines were applied for this period (Waring *et al.* 2014, 2015). One serious injury was observed in 2011. Fishery related bycatch rates for years 2008-2012 were estimated using an annual stratified ratio-estimator. These estimates replace the 2008-2010 annual estimates reported in the 2013 stock assessment report that were generated using a different method. The estimated annual mortality (CV in parentheses) attributed to this fishery was 7.2 (0.48) for 2005, 6.5 (0.49) for 2006, 5.6 (0.46) for 2007, 5.6 (0.97) for 2008, 0 for 2009 and 2010, 5.9 (0.71) for 2011, and 0 for 2012. Annual average estimated harbor porpoise mortality and serious injury from the Northeast bottom trawl fishery from 2008-2012 is 2.3 (0.60) (Table 2).

CANADA

Bay of Fundy Sink Gillnet

During the early 1980s, harbor porpoise bycatch in the Bay of Fundy sink gillnet fishery, based on casual observations and discussions with fishermen, was thought to be low. The estimated harbor porpoise bycatch in 1986 was 94-116 and in 1989 it was 130 (Trippel *et al.* 1996). The Canadian gillnet fishery occurs mostly in the western portion of the Bay of Fundy during the summer and early autumn months, when the density of harbor porpoises is highest. Polacheck (1989) reported there were 19 gillnetters active in 1986, 28 active in 1987, and 21 in 1988.

An observer program implemented in the summer of 1993 provided a total bycatch estimate of 424 harbor porpoises (± 1 SE: 200-648) from 62 observed trips, (approximately 11.3% coverage of the Bay of Fundy trips) (Trippel *et al.* 1996). During 1994, the observer program was expanded to cover 49% of the gillnet trips (171

observed trips). The bycatch was estimated to be 101 harbor porpoises (95% confidence limit: 80-122), and the fishing fleet consisted of 28 vessels (Trippel *et al.* 1996). During 1995, due to groundfish quotas being exceeded, the gillnet fishery was closed from July 21 to August 31. During the open fishing period of 1995, 89% of the trips were observed, all in the Swallowtail region. Approximately 30% of these observed trips used pingered nets. The estimated bycatch was 87 harbor porpoises (Trippel *et al.* 1996). No confidence interval was computed due to lack of coverage in the Wolves fishing grounds. During 1996, the Canadian gillnet fishery was closed during 20-31 July and 16-31 August due to groundfish quotas. From the 107 monitored trips, the bycatch in 1996 was estimated to be 20 harbor porpoises (DFO 1998; Trippel *et al.* 1999). Trippel *et al.* (1999) estimated that during 1996, gillnets equipped with acoustic alarms reduced harbor porpoise bycatch rates by 68% over nets without alarms in the Swallowtail area of the lower Bay of Fundy. During 1997, the fishery was closed to the majority of the gillnet fleet during 18-31 July and 16-31 August, due to groundfish quotas. In addition a time-area closure to reduce porpoise bycatch in the Swallowtail area occurred during 1-7 September. From the 75 monitored trips, 19 harbor porpoises were observed taken. After accounting for total fishing effort, the estimated bycatch in 1997 was 43 animals (DFO 1998). Trippel *et al.* (1999) estimated that during 1997, gillnets equipped with acoustic alarms reduced harbor porpoise bycatch rates by 85% over nets without alarms in the Swallowtail area of the lower Bay of Fundy. The number of monitored trips (and observed harbor porpoise mortalities were 111 (5) for 1998, 93 (3) for 1999, 194 (5) for 2000, and 285 (39) for 2001. The estimated annual mortality estimates were 38 for 1998, 32 for 1999, 28 for 2000, and 73 for 2001 (Trippel and Shepherd 2004). Estimates of variance are not available.

Since 2002 there has been no observer program in the Bay of Fundy region, but the fishery is still active. Bycatch for these years is unknown. The annual average of most recent five years with available data (1997-2001) was 43 animals, so this value is used to estimate the annual average for more recent years. However, in 2011 there was little gillnet effort in New Brunswick waters in the summer; thus the Canadian porpoise by-catch estimates could have been near zero. The fishermen that sought groundfish went into the mid-Bay of Fundy where traditionally by-catch levels were extremely low. Trippel (pers. comm.) estimated that less than 10 porpoise were bycaught in the Canadian fisheries in the Bay of Fundy in 2011. Analysis of port catch records might allow estimation of bycatch for more recent times, however, it would be difficult to also accurately account for the changes in the spatial distribution of the harbor porpoises and fisheries. .

Herring Weirs

Harbor porpoises are taken in Canadian herring weirs, but there have been no recent efforts to observe takes in the U.S. component of this fishery. Smith *et al.* (1983) estimated that in the 1980s approximately 70 harbor porpoises became trapped annually and, on average, 27 died annually. In 1990, at least 43 harbor porpoises were trapped in Bay of Fundy weirs (Read *et al.* 1994). In 1993, after a cooperative program between fishermen and Canadian biologists was initiated, over 100 harbor porpoises were released alive (Read *et al.* 1994). Between 1992 and 1994, this cooperative program resulted in the live release of 206 of 263 harbor porpoises caught in herring weirs. Mortalities (and releases) were 11 (50) in 1992, 33 (113) in 1993, and 13 (43) in 1994 (Neimanis *et al.* 1995). Since that time, additional harbor porpoises have been documented in Canadian herring weirs: mortalities (releases and unknowns) were 5 (60, 0) in 1995; 2 (4, 0) in 1996; 2 (24, 0) in 1997; 2 (26, 0) in 1998; 3 (89, 0) in 1999; 0 (13, 0) in 2000 (A. Read, pers. comm), 14 (296, 0) in 2001, 3 (46, 4) in 2002, 1 (26, 3) in 2003, 4 (53, 2) in 2004; 0 (19, 5) in 2005; 2 (14, 0) in 2006; 3 (9, 3) in 2007, 0 (8, 6) in 2008, 0 (3,4) in 2009, 1 in 2010 (7, 0), 0 (2, 3) in 2011, and 0 (2, 3) in 2012. (Neimanis *et al.* 2004; H. Koopman and A. Westgate, pers. comm.).

Average estimated harbor porpoise mortality in the Canadian herring weir fishery during 2008–2012 was 0.2 (Table 2). An estimate of variance is not possible.

Table 2. From observer program data, summary of the incidental mortality of Gulf of Maine/Bay of Fundy harbor porpoise (<i>Phocoena phocoena phocoena</i>) by commercial fishery including the years sampled, the type of data used, the annual observer coverage, the mortalities and serious injuries recorded by on-board observers, the estimated annual serious injury and mortality, the estimated CV of the annual mortality, and the mean annual combined mortality (CV in parentheses).										
Fishery	Years	Data Type ^a	Observer Coverage ^b	Observed Serious Injury ⁱ	Observed Mortality	Estimated Serious Injury	Estimated Mortality	Combined Serious Injury	Estimated CVs	Mean Annual Combined Mortality
U.S.										
Northeast	08-12	Obs. Data,	.05, .04, .17, .19,	0, 0, 0, 0, 0	30, 45, 50, 66, 34	0, 0, 0, 0, 0	666, 591, 387, 273, 277	666, 591, 387, 273,	.48, .23, .27, .20, .59	439 (0.18)

Sink Gillnet ^{c, h}		Weighout, Trip Logbook	.15					277		
Mid-Atlantic Gillnet	08-12	Obs. Data Weighout	.03, .03, .04, .02, .02	0, 0, 0, 0, 0	9, 7, 18, 11, 2	0, 0, 0, 0	350, 201, 259, 123, 63	350, 201, 259, 123, 63	.75, .55, .88, .41, .83	199 (0.37)
Northeast bottom trawl ^g	08-12	Obs. Data Weighout	.08, .09, .16, .26, .17	0, 0, 0, 1, 0	1, 0, 0, 1, 0	1.9, 0, 0, 2.0, 0	3.7, 0, 0, 3.9, 0	5.6, 0, 0, 5.9, 0	.97, 0, 0, .71, 0	2.3 (0.60) ^g
U.S. TOTAL	2008-2012									640 (0.17)
CANADA										
Bay of Fundy Sink Gillnet ^f	1997-2001	Can. Trips	unk		19, 5, 3, 5, 39		43, 38, 32, 28, 73		unk	43 ^f (unk)
Herring Weir ^{d,e}	08-12	Coop. Data	unk		0, 0, 1, 0, 0		0, 0, 1, 0, 0		NA	0.2 (unk)
CANADIAN TOTAL	2008-2012									43 (unk)
GRAND TOTAL										683 (unk)

NA = Not available.

- a. Observer data (Obs. Data) are used to measure bycatch rates; the U.S. data are collected by the Northeast Fisheries Science Center (NEFSC) Sea Sampling Program and At-Sea Monitoring Program; the Canadian data are collected by DFO. NEFSC collects Weighout (Weighout) landings data that are used as a measure of total effort for the U.S. gillnet fisheries. The Canadian DFO catch and effort statistical system collected the total number of trips fished by the Canadians (Can. Trips), which was the measure of total effort for the Canadian groundfish gillnet fishery. Mandatory vessel trip report (VTR) (Trip Logbook) data are used to determine the spatial distribution of fishing effort in the Northeast sink gillnet fishery. Observed mortalities from herring weirs are collected by a cooperative program between fishermen and Canadian biologists (Coop. Data).
- b. Observer coverage for the U.S. Northeast and mid-Atlantic coastal gillnet fisheries, is based on tons of fish landed. Northeast bottom trawl fishery coverages are ratios based on trips. Total observer coverage reported for bottom trawl gear and gillnet gear in the year 2010 includes only samples collected from traditional fisheries observer, but not the fishery monitors. Monitor trips were incorporated starting in 2011, the first full year of monitor coverage.
- c. Since 2002 in the Northeast gillnet fishery, harbor porpoises were taken on pingered strings within strata that required pingers but that stratum also had observed strings without pingers. For estimates made during 1998 and after, a weighted bycatch rate was applied to effort from both pingered and non-pingered hauls within a stratum. The weighted bycatch rate was:
- $$\sum_i \frac{\text{ping}_{non-pinger} \# \text{ porpoise}_i}{\text{sslandings}_i} \cdot \frac{\# \text{ hauls}_i}{\text{total} \# \text{ hauls}}$$
- There were 10, 33, 44, 0, 11, 0, 2, 8, 6, 2, 26, 2, 4, 12, 2, 9, 6, 11, 23, 11 and 30 observed harbor porpoise takes on pinger trips from 1992 to 2012, respectively, that were included in the observed mortality column. In addition, there were 9, 0, 2, 1, 1, 4, 0, 1, 7, 21, 33, 24, 7, 13, 20, 41, 11, and 31 observed harbor porpoise takes in 1995 to 2012, respectively, on trips dedicated to fish sampling versus dedicated to watching for marine mammals; these were also included in the observed mortality column.
- d. There were 255 licenses for herring weirs in the Canadian Bay of Fundy region.
- e. Data provided by H. Koopman pers. comm.
- f. The Canadian gillnet fishery was not observed during 2002 and afterwards, but the fishery is still active; thus, the current bycatch estimate for this fishery is assumed to be the average estimate using last five years that the fishery was observed in (1997-2001).
- g. Fishery related bycatch rates for years 2008-2012 were estimated using an annual stratified ratio-estimator.

- These estimates replace the 2008-2010 annual estimates reported in the 2013 stock assessment report that were generated using a different method.
- h. Thirteen harbor porpoises in the Northeast area and 10 in the mid-Atlantic area were incidentally caught as part of a 2009-2010 NEFSC gillnet hanging ratio study to examine the impact of hanging ratio on harbor porpoise bycatch in gillnets. These animals were included in the observed interactions and added to the total estimates, though these interactions and their associated fishing effort were not included in the estimation of the bycatch rate that was expanded to the rest of the fishery.
 - i. Serious injuries were evaluated for the 2008–2012 period using new guidelines and include both at-sea monitor and traditional observer data (Waring *et al.* 2014, 2015)

**Other Mortality
U.S.**

There is evidence that harbor porpoises were harvested by natives in Maine and Canada before the 1960s, and the meat was used for human consumption, oil, and fish bait (NMFS 1992). The extent of these past harvests is unknown, though it is believed to have been small. Up until the early 1980s, small kills by native hunters (Passamaquoddy Indians) were reported. In recent years it was believed to have nearly stopped (Polacheck 1989) until media reports in September 1997 depicted a Passamaquoddy tribe member dressing out a harbor porpoise. Further articles describing use of porpoise products for food and other purposes were timed to coincide with ongoing legal action in state court.

During 2008, 58 harbor porpoises were reported stranded on Atlantic U.S. beaches. Of these, four were reported as having signs of human interaction. One of these was classified as a fishery interaction.

During 2009, 65 harbor porpoises were reported stranded on Atlantic U.S. beaches. Of these, three stranding mortalities were reported as having signs of human interaction, all of which were fishery interactions.

During 2010, 82 harbor porpoises were reported stranded on Atlantic U.S. beaches. Of these, six stranding mortalities were reported as having signs of human interaction, three of which were reported to be fishery interactions.

During 2011, 164 harbor porpoises were reported stranded on Atlantic U.S. beaches. Of these, nine stranding mortalities were reported as having signs of human interaction, three of which were reported to be fishery interactions.

During 2012, 45 harbor porpoises were reported stranded on Atlantic U.S. beaches. Of these, four stranding mortalities were reported as having signs of human interaction, one of which was reported to be a fishery interaction.

Stranding data probably underestimate the extent of fishery-related mortality and serious injury because all of the marine mammals that die or are seriously injured may not wash ashore, nor will all of those that do wash ashore necessarily show signs of entanglement or other fishery-interaction. Finally, the level of technical expertise among stranding network personnel varies widely as does the ability to recognize signs of fishery interaction.

Table 4. Harbor Porpoise (*Phocoena phocoena phocoena*) reported strandings along the U.S. and Canadian Atlantic coast, 2008-2012.

Area	Year					Total
	2008	2009	2010	2011	2012	
Maine ^{a, f, h}	7	4	7	15	7	40
New Hampshire	0	0	5	1	3	9
Massachusetts ^{a, f, g, h}	25	19	28	102	25	199
Rhode Island ^b	1	1	0	4	0	6
New York ^{c, g, h}	3	9	1	11	3	27
New Jersey ^{e, f}	8	4	7	1	2	22
Pennsylvania	0	1	0	0	0	1
Delaware	0	0	2	0	0	2

Maryland	2	5	4	0	1	12
Virginia ^{e,g}	6	8	10	2	2	28
North Carolina ^d	6	14	18	28	2	68
TOTAL U.S.	58	65	82	164	45	414
Nova Scotia/Prince Edward Island ⁱ	6	6	5	13	6	36
Newfoundland and New Brunswick ^j	4	2	1	0	0	7
GRAND TOTAL	68	73	88	177	51	457

- a. In Massachusetts one animal was taken to a rehab facility in 2008. In 2011, 5 animals were released alive and one taken to rehab. One Maine animal taken to rehab in 2012.
- b. In Rhode Island in 2011, one animal classified as human interaction (HI) due to fluke amputation.
- c. One of the 2012 New York strandings classified as human interaction due to interaction with marine debris.
- d. In North Carolina one animal was immediately released in 2008.
- e. In 2009, 3 harbor porpoises were classified as fishery interactions, 2 in VA and a third in NJ.
- f. Six total HI cases in 2010; 2 in Massachusetts, 1 in Maine, 1 in North Carolina and 2 in New Jersey. One of the New Jersey records, one of the North Carolina records, and the Maine record were fishery interactions.
- g. Nine total HI cases in 2011; 5 in Massachusetts, 1 in Rhode Island, 2 in New York and 1 in Virginia. Two of these Massachusetts animals and the Virginia animal were fishery interactions.
- h. Four HI cases in 2012. One of these was a fishery interaction (Massachusetts).
- i. Data supplied by Nova Scotia Marine Animal Response Society (pers. comm.). One of the 2012 animals was trapped in a mackerel net.
- j. (Ledwell and Huntington 2009, 2010, 2011, 2012, 2013).

CANADA

The Nova Scotia Stranding Network documented whales and dolphins stranded between 1991 and 1996 on the coast of Nova Scotia (Hooker *et al.* 1997). Researchers with the Canadian Department of Fisheries and Oceans documented strandings on the beaches of Sable Island during 1970 to 1998 (Lucas and Hooker 2000). Sable Island is approximately 170 km southeast of mainland Nova Scotia. On the mainland of Nova Scotia, a total of 8 stranded harbor porpoises were recorded between 1991 and 1996: 1 in May 1991, 2 in 1993 (July and September), 1 in August 1994 (released alive), 1 in August 1994, and 3 in 1996 (March, April, and July (released alive)). On Sable Island, 8 stranded dead harbor porpoises were documented, most in January and February; 1 in May 1991, 1 in January 1992, 1 in January 1993, 3 in February 1997, 1 in May 1997, and 1 in June 1997. Two strandings during May-June 1997 were neonates (> 80 cm). The harbor porpoises that stranded in the winter (January-February) were on Sable Island, those in the spring (March to June) were in the Bay of Fundy (2 in Minas Basin and 1 near Yarmouth) and on Sable Island (2), and those in the summer (July to September) were scattered along the coast from the Bay of Fundy to Halifax.

Whales and dolphins stranded since 1997 on the coast of Nova Scotia were recorded by the Marine Animal Response Society and the Nova Scotia Stranding Network, including 3 harbor porpoises stranded in 1997 (1 in April, 1 in June and 1 in July), 2 stranded in June 1998, 1 in March 1999, 3 in 2000 (1 in February, 1 in June, and 1 in August); 2 in 2001 (1 in July and 1 in December), 5 in 2002 (3 in July (1 released alive), 1 in August, and 1 in September (released alive)), 3 in 2003 (2 in May (1 was released alive) and 1 in June (disentangled and released alive)), 4 in 2004 (1 in April, 1 in May, 1 in July (released alive) and 1 in November), 6 in 2005 (1 in April (released alive), 1 in May, 3 in June and 1 in July), 4 in 2006 (1 in June, 1 in August, 1 in September, and 1 in December), 4 in 2007, 6 in 2008, 6 in 2009 (2 released alive), 5 (1 released alive) in 2010, 13 (4 released alive) in 2011, and 6 in 2012; Table 3).

Five dead stranded harbor porpoises were reported in 2005 by the Newfoundland and Labrador Whale Release and Strandings Program, 1 in 2007 and 4 in 2008, 2 in 2009 (one dead entangled and one live release), 1 in 2010 and 0 in 2011 and 2012 (Ledwell and Huntington 2004; 2006; 2007; 2008; 2009; 2010, 2011, 2012, 2013; Table 3).

U.S. management measures taken to reduce bycatch

A ruling to reduce harbor porpoise bycatch in U.S. Atlantic gillnets was published in the Federal Register (63 FR 66464) on 02 December 1998 and became effective 01 January 1999. The Gulf of Maine portion of the Harbor Porpoise Take Reduction Plan (HPTRP) pertains to all fishing with sink gillnets and other gillnets capable of catching regulated groundfish in New England waters, from Maine through Rhode Island. For more information on this rule, please see <http://www.greateratlantic.fisheries.noaa.gov/protected/porptrp/>.

STATUS OF STOCK

This is not a strategic stock because average annual human-related mortality and serious injury does not exceed PBR. The total U.S. fishery-related mortality and serious injury for this stock is not less than 10% of the calculated PBR and, therefore, cannot be considered to be insignificant and approaching zero mortality and serious injury rate. The status of harbor porpoises, relative to OSP, in the U.S. Atlantic EEZ is unknown. Population trends for this species have not been investigated. On 7 January 1993, NMFS proposed listing the Gulf of Maine harbor porpoise as threatened under the Endangered Species Act (NMFS 1993). On 5 January 1999, NMFS determined the proposed listing was not warranted (NMFS 1999). On 2 August 2001, NMFS made available a review of the biological status of the Gulf of Maine/Bay of Fundy harbor porpoise population. The determination was made that listing under the Endangered Species Act (ESA) was not warranted, and this stock was removed from the ESA candidate species list (NMFS 2001).

REFERENCES CITED

- AIS, Inc. 2010. The effects of hanging ratio on marine mammal interactions and catch retention of commercially important finfish species. NOAA Contract No. EA133F-08-CN-0240 Final report: 28 pp. <http://nefsc.noaa.gov/publications/reports/EA133F08CN0240.pdf>.
- Andersen, M. S., K. A. Forney, T. V. N. Cole, T. Eagle, R. Angliss, K. Long, L. Barre, L. Van Atta, D. Borggaard, T. Rowles, B. Norberg, J. Whaley, and L. Engleby 2008. Differentiating serious and non-serious injury of marine mammals: Report of the serious injury technical workshop. NOAA Tech. Memo. NMFS-OPR-39.
- Angliss, R.P. and D.P. DeMaster. 1998. Differentiating serious and non-serious injury of marine mammals taken incidental to commercial fishing operations: Report of the serious injury workshop, 1-2 April 1997, Silver Spring, MD. NOAA Tech. Memo. NMFS-OPR-13. 48 pp.
- Barlow, J. and P. Boveng 1991. Modeling age-specific mortality for marine mammal populations. Mar. Mamm. Sci. 7: 50-65.
- Benjamins, S., J. Lawson and G. Stenson 2007. Recent harbour porpoise bycatch in gillnet fisheries in Newfoundland and Labrador, Canada. J. Cetacean Res. Manage. 9(3): 189-199.
- Bravington, M.V. and K.D. Bisack 1996. Estimates of harbour porpoise bycatch in the Gulf of Maine sink gillnet fishery, 1990-1993. Rep. Int. Whal. Comm. 46: 567-574.
- Caswell, H., S. Brault, A.J. Read and T.D. Smith 1998. Harbor porpoise and fisheries: An uncertainty analysis of incidental mortality. Ecol. Appl. 8(4): 1226-1238.
- CUD 1994. Estimating harbor porpoise bycatch in the Gulf of Maine sink gillnet fishery. Conservation and Utilization Division. Northeast Fish. Sci. Cent. Ref. Doc. 94-24.
- DFO 1998. Harbour porpoise bycatch in the lower Bay of Fundy gillnet fishery. DFO Maritimes Regional Fisheries Status Report 98/7E. Available from Department of Fisheries and Oceans, Resource management Branch, P.O. Box 550, Halifax, NS B3J 2S7, Canada.
- Gaskin, D.E. 1977. Harbour porpoise, *Phocoena phocoena* (L.), in the western approaches to the Bay of Fundy 1969-75. Rep. Int. Whal. Comm. 27: 487-492.
- Gaskin, D.E. 1984. The harbor porpoise *Phocoena phocoena* (L.): Regional populations, status, and information on direct and indirect catches. Rep. Int. Whal. Comm. 34: 569-586.
- Gaskin, D.E. 1992. The status of the harbour porpoise. Can. Field-Nat. 106: 36-54.
- Gilbert, J.R. 1987. Marine Mammal Interaction with New England Gillnet Fisheries. NMFS. NA84EAC00070. draft report 21 pp. Available at <http://nefsc.noaa.gov/publications/reports/NA84EAC00070.pdf>
- Gilbert, J.R. and K.M. Wynne 1985. Harbor seal populations and fisheries interactions with marine mammals in New England, 1984. NMFS. NA80FAC00029 and NA84EAC00070 15 pp. Available at <http://nefsc.noaa.gov/publications/reports/NA80FAC00029.pdf>
- Hatch, J.M. and C.D. Orphanides 2014. Estimates of cetacean and pinniped bycatch in the 2012 New England sink and mid-Atlantic gillnet fisheries. Northeast Fish Sci Cent Ref Doc. 14-02. 20 pp.
- Hiby, L. 1999. The objective identification of duplicate sightings in aerial survey for porpoise. Pages 179-189 in: G. W. Garner, S. C. Amstrup, J. L. Laake *et al.*, (eds.) Marine Mammal Survey and Assessment Methods.

- Balkema, Rotterdam.
- Hiltunen, K.H. 2006. Mixed-stock analysis of harbor porpoises (*Phocoena phocoena*) along the U.S. mid-Atlantic coast using microsatellite DNA markers. MS thesis. The College of Charleston, Charleston, SC. 92 pp.
- Hooker, S.K., R.W. Baird and M.A. Showell 1997. Cetacean strandings and bycatches in Nova Scotia, Eastern Canada, 1991-1996. Meeting document SC/49/O5 submitted to the 1997 International Whaling Commission Scientific Committee meeting in Bournemouth, UK.
- Johnston, D.W. 1995. Spatial and temporal differences in heavy metal concentrations in the tissues of harbour porpoises (*Phocoena phocoena* L.) from the western North Atlantic. M.S. thesis. University of Guelph, Guelph, Ontario, Canada. 152 pp.
- Kraus, S.D., J.H. Prescott and G.S. Stone 1983. Harbor porpoise, *Phocoena phocoena*, in the U.S. coastal waters off the Gulf of Maine: a survey to determine seasonal distribution and abundance. NMFS. NA82FAC00027 22 pp.
- Kraus, S.D., A.J. Read, A. Solow, K. Baldwin, T. Spradlin, E. Anderson and J. Williamson 1997. Acoustic alarms reduce porpoise mortality. *Nature* 388(6642): 525.
- Laake, J.L., J. Calambokidis, S.D. Osmek, and D.J. Rugh 1997. Probability of detecting harbor porpoise from aerial surveys: estimating $g(0)$. *J. Wildl. Manage.* 63-75.
- Laake, J.L., and D.L. Borchers 2004. Methods for incomplete detection at distance zero, In: *Advanced distance sampling*, edited by S. T. Buckland, D. R. Andersen, K. P. Burnham, J. L. Laake, and L. Thomas, pp. 108-189, Oxford University Press, New York.
- Lamb, A. 2000. Patterns of harbor porpoise mortality in two US Atlantic sink gillnet fisheries and changes in life history parameters. M.S. thesis. Boston University, Boston, MA.
- Lawson, J.W. and J.-F. Gosselin 2009. Distribution and preliminary abundance estimates for cetaceans seen during Canada's Marine Megafauna Survey - A component of the 2007 TNASS. *Can. Sci. Advisory Sec. Res. Doc.* 208/031. 33 pp. http://www.dfo-mpo.gc.ca/csas-sccs/publications/resdocs-docrech/2009/2009_031-eng.htm
- Lawson, J.W. and J.-F. Gosselin. 2011. Fully-corrected cetacean abundance estimates from the Canadian TNASS survey. Working Paper 10. National Marine Mammal Peer Review Meeting. Ottawa, Can. 28 pp.
- Ledwell, W. and J. Huntington 2004. Marine animal entrapments in fishing gear in Newfoundland and Labrador and a summary of the Whale Release and Strandings Program during 2004. Report to the Department of Fisheries and Oceans. St. John's, Newfoundland, Canada. 3 pp.
- Ledwell, W. and J. Huntington 2006. Whale, leatherback sea turtles. And basking shark entrapments in fishing gear in Newfoundland and Labrador and a summary of the Whale Release and Strandings Program during 2005. Report to the Department of Fisheries and Oceans. St. John's, Newfoundland, Canada. 18 pp.
- Ledwell, W. and J. Huntington 2007. Whale and leatherback sea turtle entrapment in fishing gear in Newfoundland and Labrador and a summary of the Whale Release and Strandings Program during 2006. Report to the Department of Fisheries and Oceans. St. John's, Newfoundland, Canada. 3 pp.
- Ledwell, W. and J. Huntington 2008. Incidental entrapments in fishing gear reported in 2007 in Newfoundland and Labrador and a summary of the Whale Release and Strandings Program. A report to the Department of Fisheries and Oceans, St. John's, Newfoundland, Canada. 20 pp.
- Ledwell, W. and J. Huntington 2009. Incidental entrapments in fishing gear and strandings reported to the whale release and strandings group in Newfoundland and Labrador and a summary of the Whale Release and Strandings Program during 2008. A report to the Department of Fisheries and Oceans Canada, St. John's, Newfoundland, Canada. 29 pp.
- Ledwell, W. and J. Huntington 2010. Incidental entrapments in fishing gear and strandings reported to the whale release and strandings group in Newfoundland and Labrador and a summary of the Whale Release and Strandings Group in Newfoundland and Labrador and a summary of the whale release and strandings program during 2009-2010. A report to the Department of Fisheries and Oceans Canada, St. John's, Newfoundland, Canada. 23 pp.
- Ledwell, W. and J. Huntington 2011. Whale, leatherback sea turtles. and basking sharks entrapped in fishing gear in Newfoundland and Labrador and a summary or the strandings, sightings and education work during 2010-2011. A preliminary report to Fisheries and Oceans Canada, St. John's, Newfoundland, Canada. 25 pp.
- Ledwell, W. and J. Huntington 2012. Incidental entrapments of cetaceans and leatherback sea turtles in fishing gear reported during 2011-2012 and a summary of the Whale Release and Stranding Group activities. A preliminary report to Fisheries and Oceans Canada, St. John's, Newfoundland, Canada. 17 pp.
- Ledwell, W. and J. Huntington 2012. Incidental entrapments in fishing gear and stranding reported to and responded to by the Whale Release and Strandings Group in Newfoundland and Labrador and a summary of the

- Whale Release and Strandings program during 2012. Report to Fisheries and Oceans Canada, St. John's, Newfoundland, Canada. 18 pp.
- Lesage, V., J. Keays, S. Turgeon and S. Hurtubise 2006. Bycatch of harbour porpoises (*Phocoena phocoena*) in gillnet fisheries of the Estuary and Gulf of St. Lawrence in 2000-2002. *J. Cetacean Res. Manage.* 8(1): 67-78.
- Lucas, Z.N. and S.K. Hooker 2000. Cetacean strandings on Sable Island, Nova Scotia, 1970-1998. *Can. Field-Nat.* 114(1): 46-61.
- Moore, J.E. and A.J. Read 2008. A Bayesian uncertainty analysis of cetacean demography and bycatch mortality using age-at-death data. *Ecol. Appl.* 18(8): 1914-1931.
- Neimanis, A.S., H.N. Koopman, A.J. Westgate, L.D. Murison and A.J. Read 2004. Entrapment of harbour porpoises (*Phocoena phocoena*) in herring weirs in the Bay of Fundy, Canada. *J. Cetacean Res. Manage.* 6(1): 7-17.
- Neimanis, A.S., A.J. Read, A.J. Westgate, H.N. Koopman, J.Y. Wang, L.D. Murison and D.E. Gaskin 1995. Entrapment of harbour porpoises (*Phocoena phocoena*) in herring weirs in the Bay of Fundy, Canada. International Whaling Commission, Working paper SC/47/SM18.
- NMFS 1992. Harbor porpoise in Eastern North America: Status and Research Needs. Results of a scientific workshop held May 5-8, 1992 at NEFSC, Woods Hole, MA, USA. Northeast Fish. Sci. Cent. Ref. Doc., 92-06. National Marine Fisheries Service. 28 pp.
- NMFS 1993. Proposed listing of Gulf of Maine population of harbor porpoises as threatened under the Endangered Species Act. Federal Register 58: 3108-3120.
- NMFS 1999. Listing of Gulf of Maine/Bay of Fundy population of harbor porpoise as threatened under the Endangered Species Act. Federal Register 64(2): 465-471.
- NMFS 2001. Status review of the Gulf of Maine/Bay of Fundy population of harbor porpoise under the Endangered Species Act (ESA). Federal Register 66(203): 53195-53197.
- NMFS 2006 NOAA Twin Otter aircraft circle-back abundance survey. Unpublished report. 26 pp. <http://www.nefsc.noaa.gov/psb/surveys/documents/CruiseReport.aerial2006.full.pdf>.
- NOAA. 2012. Federal Register 77:3233. National Policy for Distinguishing Serious From Non-Serious Injuries of Marine Mammals. Available from: <http://www.nmfs.noaa.gov/op/pds/documents/02/238/02-238-01.pdf>.
- Orphanides, C.D. and D.L. Palka. 2013. Analysis of harbor porpoise gillnet bycatch, compliance, and enforcement trends in the US northwestern Atlantic, January 1999 to May 2010. *Endang Species Res* 20: 251-269.
- Orphanides, C.D. and D.L. Palka. 2012. 2010-2011 HPTRP consequential bycatch and compliance rates. US Dept Commer, Northeast Fish Sci Center Ref Doc. 12-08; 25 p. Available at: <http://www.nefsc.noaa.gov/publications/crd/crd1208/>
- Orphanides, C. D. 2012. New England harbor porpoise bycatch rates during 2012-2012 associated with Consequence Closure Areas. US Dept Commer, Northeast Fish Sci Center Ref Doc. 12-19; 15 p. Available at: <http://www.nefsc.noaa.gov/publications/crd/crd1219/>
- Orphanides, C.D. 2013. Estimates of cetacean and pinniped bycatch during 2010 and 2011 in the New England Sink Gillnet fishery, Mid-Atlantic Gillnet fishery, and two NMFS gillnet experiments. Northeast Fish Sci Cent Ref Doc. 13-13 38 pp. Available at: <http://nefsc.noaa.gov/publications/crd/crd1313/>.
- Palka, D. 1995a. Influences on spatial patterns of Gulf of Maine harbor porpoises. Pages 69-75 in: A. S. Blix, L. Walloe and O. Ulltang, (eds.) *Whales, Seals, Fish and Man*. Elsevier Science. Amsterdam.
- Palka, D.L., A.J. Read, A.J. Westgate and D.W. Johnston 1996. Summary of current knowledge of harbour porpoises in US and Canadian Atlantic waters. *Rep. Int. Whal. Comm.* 46: 559-565.
- Palka, D.L. 2005. Aerial surveys in the northwest Atlantic: estimation of g(0), Proceedings of a Workshop on Estimation of g(0) in Line-Transect Surveys of Cetaceans. European Cetacean Society's 18th Annual Conference; Kolmården, Sweden; Mar. 28, 2004.
- Palka, D. 2000. Abundance of the Gulf of Maine/Bay of Fundy harbor porpoise based on shipboard and aerial surveys during 1999. Northeast Fish. Sci. Cent. Ref. Doc. 00-07. 29 pp. <http://www.nefsc.noaa.gov/psb/pubs/palkalabref00-07.pdf>
- Palka, D.L. 2012. Cetacean abundance estimates in US northwestern Atlantic Ocean waters from summer 2011 line transect survey. Northeast Fish. Sci. Cent. Ref. Doc. 12-29. 37 pp. <http://www.nefsc.noaa.gov/nefsc/publications/crd/crd1229/>
- Polacheck, T. 1989. Harbor porpoises and the gillnet fishery. *Oceanus* 32(1): 63-70.
- Polacheck, T., F.W. Wenzel and G. Early 1995. What do stranding data say about harbor porpoise (*Phocoena phocoena*)? *Rep. Int. Whal. Comm.* (Special Issue) 16: 169-180.
- Read, A.J., J.E. Craddock and D. Gannon 1994. Life history of harbour porpoises and pilot whales taken in commercial fishing operations off the northeastern United States. Final Report, Phase II. 50-EANE-2-

00082. final report, phase II.
- Read, A.J. and A.A. Hohn 1995. Life in the fast lane: the life history of harbour porpoises from the Gulf of Maine. *Mar. Mamm. Sci.* 11(4): 423-440.
- Read, A.J., J.R. Nicolas and J.E. Craddock 1996. Winter capture of a harbor porpoise in a pelagic drift net off North Carolina. *Fish. Bull.* 94(2): 381-383.
- Read, A.J. and A.J. Westgate 1997. Monitoring the movements of harbour porpoises (*Phocoena phocoena*) with satellite telemetry. *Ma. Biol.* 130: 315-22.
- Read, A.J. 2013. Development of conservation strategies to mitigate the bycatch of harbor porpoises in the Gulf of Maine. *Endang Species Res* 20:235-250.
- Rosel, P.E., S.C. France, J.Y. Wang and T.D. Kocher 1999a. Genetic structure of harbour porpoise *Phocoena phocoena* populations in the northwest Atlantic based on mitochondrial and nuclear markers. *Mol. Ecol.* 8: S41-S54.
- Rosel, P.E., R. Tiedemann and M. Walton 1999b. Genetic evidence for limited trans-Atlantic movements of the harbor porpoise *Phocoena phocoena*. *Mar. Biol.* 133: 583-591.
- Rossmann, M.C. and R.L. Merrick 1999. Harbor porpoise bycatch in the Northeast Multispecies Gillnet Fishery and the mid-Atlantic Coastal Gillnet Fishery in 1998 and during January-May 1999. *Northeast Fish. Sci. Cent. Ref. Doc.* 99-17. 36 pp. <http://www.nefsc.noaa.gov/nefsc/publications/crd/crd9917.pdf>
- Schofield, D.T., G. Early, F.W. Wenzel, K. Matassa, C. Perry, G. Beekman, B. Whitaker, E. Gebhard, W. Walton and M. Swingle 2008. Rehabilitation and homing behavior of a satellite-tracked harbor porpoise (*Phocoena phocoena*). *Aq. Mamm.* 34(1): 1-8.
- Smith, G.J.D., A.J. Read and D.E. Gaskin 1983. Incidental catch of harbor porpoises, *Phocoena phocoena* (L.), in herring weirs in Charlotte County, New Brunswick, Canada. *Fish. Bull.* 81(3): 660-662.
- Taylor, B.L., M. Martinez, T. Gerrodette, J. Barlow and Y.N. Hrovat. 2007. Lessons from monitoring trends in abundance in marine mammals. *Mar. Mamm. Sci.* 23(1): 157-175.
- Thomas L, J.L. Laake, E. Rexstad, S. Strindberg, F.F.C. Marques, S.T. Buckland, D.L. Borchers, D.R. Anderson, K.P. Burnham, M.L. Burt, S.L. Hedley, J.H. Pollard, J.R.B. Bishop and T.A. Marques. 2009. Distance 6.0. Release 2. [Internet]. University of St. Andrews (UK): Research Unit for Wildlife Population Assessment. Available from: <http://www.ruwpa.st-and.ac.uk/distance/>.
- Trippel, E.A. and T.D. Shepherd 2004. By-catch of harbour porpoise (*Phocoena phocoena*) in the Lower Bay of Fundy gillnet fishery from 1998-2001. Department of Fisheries and Oceans. Ottawa, Ontario. DFO Research Document 2004/2521 iv + 33 pp. http://www.fmap.ca/ramweb/papers-total/Trippel_Shepherd_2004.pdf
- Trippel, E.A., M.B. Strong, J.M. Terhune and J.D. Conway 1999. Mitigation of harbour porpoise (*Phocoena phocoena*) bycatch in the gillnet fishery in the lower Bay of Fundy. *Can. J. Fish. Aquat. Sci* 56: 113-123.
- Trippel, E.A., J.Y. Wang, M.B. Strong, L.S. Carter and J.D. Conway 1996. Incidental mortality of harbour porpoise (*Phocoena phocoena*) by the gill-net fishery in the lower Bay of Fundy. *Can. J. Fish. Aquat. Sci* 53: 1294-1300.
- Wade, P.R. and R.P. Angliss 1997. Guidelines for assessing marine mammal stocks: Report of the GAMMS Workshop April 3-5, 1996, Seattle, Washington. NOAA Tech. Memo. NMFS-OPR-12. 93 pp.
- Wang, J.Y., D.E. Gaskin and B.N. White 1996. Mitochondrial DNA analysis of harbour porpoise, *Phocoena phocoena*, subpopulations in North American waters. *Can. J. Fish. Aquat. Sci* 53: 1632-45.
- Waring, G.T., F.W. Wenzel M.C., E. Josephson, and M. Lyssikatos 2014. Serious injury determinations for small cetaceans and pinnipeds caught in commercial fisheries off the northeast U.S. coast, 2007–2011. *Northeast Fish. Sci. Cent. Ref. Doc.* 14-13. 26 p.
- Waring, G.T., M.C. Rossmann and F.W. Wenzel. 2015 Serious injury determinations for small cetaceans and seals caught in commercial fisheries off the northeast U.S. coast, 2012. *Northeast Fish. Sci. Cent. Ref. Doc.* 15-12. 19 p.
- Westgate, A.J., D.C.G. Muir, D.E. Gaskin and M.C.S. Kingsley 1997. Concentrations and accumulation patterns of organochlorine contaminants in the blubber of harbour porpoises, *Phocoena phocoena*, from the coast of Newfoundland, the Gulf of St. Lawrence and the Bay of Fundy/Gulf of Maine. *Envir. Pollut.* 95: 105-119.
- Westgate, A.J., A.J. Read, T.M. Cox, T.D. Schofield, B.R. Whitaker and K.E. Anderson 1998. Monitoring a rehabilitated harbor porpoise using satellite telemetry. *Mar. Mamm. Sci.* 14(3): 599-604.
- Westgate, A.J. and K.A. Tolley 1999. Geographical differences in organochlorine contaminants in harbour porpoises *Phocoena phocoena* from the western North Atlantic. *Mar. Ecol. Prog. Ser.* 177: 255-268.
- Woodley, T.H. and A.J. Read 1991. Potential rates of increase of a harbor porpoise (*Phocoena phocoena*) population subjected to incidental mortality in commercial fisheries. *Can. J. Fish. Aquat. Sci* 48: 2429-35.